

© Kirby Institute, UNSW Sydney

ISSN 1448-5915

This publication is available at: <http://www.kirby.unsw.edu.au>

Kirby Institute
UNSW Sydney
Sydney NSW 2052
Australia

Telephone: **02 9385 0900** Facsimile: **02 6100 2860** International prefix: **612** Email: info@kirby.unsw.edu.au

AUSTRALIAN NSP SURVEY
Prevalence of HIV, HCV and injecting and
sexual behaviour among NSP attendees

NATIONAL DATA REPORT
2018 - 2022

Report prepared by
Sue Heard, Farzaneh Zolala and Lisa Maher

Kirby Institute
UNSW Sydney
Sydney NSW 2052
Australia

July 2023

Any presentation of these data should include acknowledgement of the Australian Needle Syringe Program Survey, Kirby Institute, UNSW Sydney.

Citation: Heard, S; Zolala, F & Maher, L. (2023). Australian Needle Syringe Program Survey National Data Report 2018-2022: Prevalence of HIV, HCV and injecting and sexual behaviour among NSP attendees. Sydney: Kirby Institute, UNSW Sydney.

The Australian Needle Syringe Program Survey is funded by the Australian Government Department of Health and Aged Care. Kirby Institute, Faculty of Medicine & Health, UNSW Sydney.

ACKNOWLEDGEMENTS

We would like to acknowledge the many people who assist each year in the development and conduct of the Australian Needle Syringe Program Survey (ANSPS), particularly the clients, staff and managers at participating Needle Syringe Program (NSP) services. We also appreciate the dedication and vision of the founding members of the project and the late Dr Margaret MacDonald who was responsible for the development and conduct of the ANSPS from 1995 until 2003.

Ethical approvals for the survey are obtained from institutional ethics committees associated with the investigators and participating NSP sites. The Australian Needle Syringe Program Survey is funded by the Australian Government Department of Health and Aged Care.

Special thanks go to Mr Philip Cunningham, Senior Scientist and Operations Manager, Dr Beth Catlett, Post-Doctoral Research Scientist, Mr Mitchell Starr, Senior Hospital Scientist, Ms Thidarat Danwilai, Technical Officer, Mr Michael Palmer, Technical Officer and Mr Andrew Kelly, Technical Officer at the NSW State Reference Laboratory for HIV at St Vincent's Hospital and St Vincent's Centre for Applied Medical Research. We also appreciate the assistance provided by Ms Rachel McCleave from the Kirby Institute.

In 2022, the project received support and input from the following members of the ANSPS National Advisory Group: Mr Phillip Hull (ACT); Ms Kate Patten (NSW); Ms Anshul Kaul (NT); Ms Michelle Spudic (SA); Ms Myf Briggs (TAS); Mr Gary Morris and Ms Terrie Spall (VIC); Ms Jude Bevan (WA); Mr Adrian Gorringe (Australian Injecting and Illicit Drug Users League); and Ms Sue Heard, Dr Jenny Iversen and Professor Lisa Maher (Kirby Institute). We would particularly like to thank Mr Robert Kemp (QLD) for chairing the National Advisory Group (from 2011 to 2022) and to welcome Ms Jude Bevan as the incoming chair (2023).

In this report, we also wish to acknowledge and thank Dr Jenny Iversen who retired at the end of 2022 after a long and successful career in the harm reduction and research sectors, including 15 years at the Kirby Institute co-ordinating and eventually overseeing flagship surveillance projects such as the ANSPS. Dr Iversen's research on the epidemiology of hepatitis C infection in people who inject drugs has been recognised internationally, and her innovative work on HCV incidence and viraemic prevalence has informed global efforts to eliminate hepatitis C. The ANSPS has benefited greatly from her expertise and efforts and she will be missed.

ABBREVIATIONS

| | |
|------------|--|
| ACT | Australian Capital Territory |
| ANSPS | Australian Needle Syringe Program Survey |
| COVID-19 | Coronavirus disease 2019 |
| DAA | Direct acting antiviral |
| DBS | Dried blood spot |
| HCV | Hepatitis C virus |
| HIV | Human immunodeficiency virus |
| NSP | Needle syringe program |
| NSW | New South Wales |
| NT | Northern Territory |
| OAT | Opioid agonist therapy |
| PIEDs | Performance and image-enhancing drugs |
| PWID | People who inject drugs |
| PBS | Pharmaceutical Benefits Scheme |
| QLD | Queensland |
| RNA | Ribonucleic acid |
| SA | South Australia |
| SARS-CoV-2 | Severe Acute Respiratory Syndrome - Corona Virus 2 |
| TAS | Tasmania |
| VIC | Victoria |
| WA | Western Australia |

TABLE OF CONTENTS

Summary

| | |
|--------------------------------------|----|
| Key Findings | i |
| Introduction | 1 |
| COVID-19 | 1 |
| Demographic characteristics | 2 |
| Injection behaviour | 3 |
| Sexual behaviour..... | 5 |
| Drug treatment | 5 |
| HIV and HCV diagnostic testing | 6 |
| HIV antibody prevalence | 6 |
| HCV antibody prevalence..... | 8 |
| HCV RNA prevalence..... | 9 |
| HCV treatment | 11 |
| References..... | 13 |

Tables

| | |
|---------------------------------------|-----|
| 1. National | 14 |
| 2. Australian Capital Territory | 42 |
| 3. New South Wales..... | 62 |
| 4. Northern Territory | 90 |
| 5. Queensland | 110 |
| 6. South Australia..... | 138 |
| 7. Tasmania | 158 |
| 8. Victoria | 178 |
| 9. Western Australia | 206 |

Appendix

| | |
|------------------------------------|-----|
| A. Participating NSP services..... | 234 |
| B. Methodological notes | 236 |
| C. List of tables | 238 |

KEY FINDINGS

- The Australian NSP Survey (ANSPS) functions as a strategic early-warning system designed to monitor blood borne viral infections and injecting and sexual behaviour among people who inject drugs.
- In 2022, 53 Needle Syringe Programs (NSPs) throughout Australia participated in the ANSPS and 1,781 NSP attendees completed the survey. The response rate was 34%.
- The number of respondents in 2020, 2021 and 2022 was lower than in previous years due to the ongoing impacts of the COVID-19 pandemic and public health measures designed to reduce community transmission during ANSPS data collection periods.

COVID-19

- As a sentinel surveillance system with the capacity to respond to emerging issues, the 2020, 2021 and 2022 surveys included a module about COVID-19-related impacts, including vaccination uptake and, in 2022, questions designed to assess barriers to vaccine uptake.
- In 2022, the majority (78%) of respondents reported they had been tested for COVID-19 in the previous 12 months. One third (36%) of those reporting ever being tested for COVID-19 reported a positive result in the past 12 months and 6% reported a positive result more than a year ago.
- All respondents were eligible for COVID-19 vaccination (two primary doses plus booster) at the time of survey completion in October 2022, and the majority (82%) reported receiving at least one dose of a COVID-19 vaccine. This includes 46% reporting three or more doses and 29% reporting two doses.
- Among participants who reported having no doses of a COVID-19 vaccine (unvaccinated), one third (31%) indicated hesitancy due to concerns about vaccine safety or side effects. Among participants who reported having one or two doses of a COVID-19 vaccine, almost half (47%) reported the reason for not completing vaccination was that they hadn't got around to it yet.

Demographic characteristics

- Over the period 2018 to 2022, the median age of survey respondents was stable (range 42 to 45 years), with a decline in the proportion of young injectors (aged <25 years) from 4% in 2018 to 3% in 2022.
- The median time since first injection remained stable between 2018 and 2022 (range 22 to 24 years), and the proportion of new initiates (those injecting for less than three years) was also stable between 2018 and 2022 (range 4% to 6%).
- The proportion of respondents from an Aboriginal and Torres Strait Islander background increased from 19% in 2018 to 25% in 2022.
- The proportion of respondents reporting incarceration in the 12 months prior to survey completion decreased from 13% in 2018 to 11% in 2022.
- The proportion of respondents reporting sex work in the month prior to the survey remained low and stable at 5% to 6% over the last five years. Condom use at last paid sex was also stable over the five-year period 2018 to 2022 (range 44% to 70%).

Injection behaviour

- One in two respondents (51%) reported injecting daily or more frequently in all survey years between 2018 and 2022.
- The proportion of respondents reporting methamphetamine as the last drug injected was stable from 2018 to 2022 (range 47% to 51%). Methamphetamine was the most commonly reported drug last injected nationally in all years between 2018 and 2022.

- The proportion of respondents reporting heroin as the last drug injected declined from 26% in 2018 to 25% in 2022, while prevalence of pharmaceutical opioid injection declined from 7% in 2018 to 5% in 2022.
- The proportion of respondents who reported last injecting performance and image enhancing drugs (PIEDs) increased from 4% in 2018 to 7% in 2022.
- The proportion of respondents reporting injecting risk behaviours was stable across all three key indicators (receptive sharing of syringes, receptive sharing of drug preparation equipment and re-use of syringes) over the past five years 2018 to 2022.
- The prevalence of self-reported drug overdose in the previous 12 months declined from 19% in 2018 to 12% in 2022.

Drug treatment

- Engagement with drug treatment services remained high, with two thirds (67%) of respondents in 2022 reporting a lifetime history of drug treatment.
- Although there was a decline in the overall proportion of respondents who reported current engagement in opioid agonist therapy (OAT) between 2018 and 2022 (from 31% to 28%), engagement in OAT remained stable among respondents who reported last injecting an opioid (range 44% to 53%).

HIV antibody prevalence

- HIV antibody prevalence remained low and stable nationally, ranging from 1.5% to 2.5% over the five-year period 2018 to 2022.
- In all years 2018 to 2022 HIV antibody prevalence was higher among homosexual men, compared to bisexual or heterosexual men, and compared to women.

HCV antibody prevalence

- Hepatitis C virus (HCV) antibody prevalence declined nationally from 45% in 2018 to 32% in 2022. This is the sixth consecutive year that less than half of respondents had this marker indicating exposure to HCV, following two decades of HCV antibody prevalence $\geq 50\%$ (all years 1999 to 2016).
- Across all survey years 2018 to 2022, prevalence of HCV antibody was higher among older than among younger respondents and among those with longer injection histories.

HCV RNA prevalence

- The proportion of respondents with detectable HCV RNA (active infection) declined from 20% in 2018 to 12% in 2022. HCV RNA prevalence declined among both males (21% in 2018 to 13% in 2022) and females (19% in 2018 to 11% in 2022).

HCV treatment uptake

- Among respondents assessed as eligible for HCV treatment, the proportion who reported a lifetime history of HCV treatment increased, from 55% in 2018 to 68% in 2022.
- However, among respondents assessed as eligible for HCV treatment, the proportion who reported recent (last 12 months) treatment decreased over the 5-year period 2018 (39%) to 2022 (34%).
- There was also a decline in the last five years in the proportion of respondents who reported a recent HCV diagnostic test from 2018 (53%) to 2022 (49%).

INTRODUCTION

The Australian Needle Syringe Program Survey (ANSPS) provides serial point prevalence estimates of human immunodeficiency virus (HIV) and hepatitis C virus (HCV) antibody prevalence, HCV ribonucleic acid (RNA) prevalence and sexual and injecting risk behaviour among people who inject drugs (PWID) in Australia. Conducted annually over a one-two week period in October, all clients attending participating Needle Syringe Program (NSP) services are invited to complete a brief, anonymous questionnaire and to provide a capillary blood sample for HIV and HCV testing (see methodological notes [Appendix B] for further details on the ANSPS design and implementation). Commencing in 2015, HCV RNA testing has also been conducted among respondents with sufficient dried blood spot (DBS) samples. This report presents national and state/territory findings for the period 2018 to 2022.

COVID-19

In Australia, a range of public health measures have been implemented since COVID-19 was declared a global pandemic by the World Health Organisation in March 2020. As a sentinel surveillance system with the capacity to respond to emerging issues, the 2020, 2021 and 2022 ANSPS included a module about the impacts of COVID-19¹.

Following two years with participation by a reduced number of NSP services due to impacts of the COVID-19 pandemic, in 2022 53 sites participated in the 2022 ANSPS (compared to 37 sites in 2021 and 38 sites in 2020). In the 2022 ANSPS the sample size was 1,781 participants which is an increase compared to 2020 (n=1324) and 2021 (n=1474), however participation remained lower than in pre-COVID years (n=2,742 in 2018, n=2,609 in 2019). The annual response rate over the past five years ranged from 34% to 45% (Table 1.1.1). Of the 1,781 ANSPS respondents in 2022, the majority (72%) were recruited from NSP services (n=26) that participated in the ANSPS in each of the past five years (2018 to 2022). Appendix A provides details of participating NSP services over the period 2018 to 2022.

Among the overall 1,781 ANSPS respondents in 2022, 1,745 (98%) completed at least one question in the COVID-19 module. The following analysis excludes ANSPS respondents who did not report responses to individual 2022 COVID-19 module questions.

Almost eighty percent of respondents (78%, n=1,392) reported that they had a COVID-19 test in the last 12 months and 7% (n=129) reported having a COVID-19 test more than 12 months ago. Of those who reported ever having a COVID-19 test (n=1,521), 642 respondents (42%) reported a positive COVID-19 test result, with the majority of these in the past 12 months (Table 1.1.7).

In 2022, 82% of respondents (n=1,409) reported having at least one dose of a COVID-19 vaccine. This includes 46% (n=787) reporting three or more doses, 29% (n=496) reporting two doses and 7% (n=126) of participants reporting having received one dose of a COVID-19 vaccine.

Among respondents reporting at least one dose of COVID-19 vaccine, the type of vaccine was predominantly Pfizer with 65% (n=913) of respondents reporting they had received the Pfizer vaccine. Almost a third (29%, n=415) reported receiving AstraZeneca vaccine, 14% (n=197) reported they did not know the type of vaccine they received and 7% (n=101) reported receiving

the Moderna vaccine. Two fifths (39%, n=565) of participants received their last COVID-19 vaccination at a GP or medical centre, one fifth (21%, n=304) at a pharmacy and almost one fifth at a vaccine hub (18%, n=256). A minority of participants received their last vaccination in prison (6%, n=91), at an NSP (4%, n=52) or at an opioid treatment clinic (2%, n=30).

The 2022 ANSPS COVID-19 module also included questions on barriers to COVID-19 vaccine uptake and vaccine hesitancy. Among participants who reported not having any doses of a COVID-19 vaccine (n=304, 17%) and excluding those who did not report a reason, the main reasons given were concerns about vaccine safety (n=49, 17%), concerns about vaccine side effects (n=37, 14%), hadn't got around to it yet (n=37, 14%), personal choice (n=36, 13%), not perceived at risk of serious illness (n=30, 11%) and n=23 (8%) who reported that they didn't believe in the vaccine or thought COVID-19 was a hoax or conspiracy.

Of the participants who reported having one or two doses of a COVID-19 vaccine (n=622, 35%) and excluding those who did not report a reason, the main reasons provided for not receiving three or more doses of a COVID vaccine were that they hadn't got around to it yet (n=270, 47%), did not perceive themselves as being at risk of serious illness (n=61, 10%), or had concerns about vaccine side effects (n=54, 9%). A minority of participants reported personal choice (n=41, 7%), concerns about vaccine safety (n=40, 7%) and difficulty getting an appointment (n=28, 5%) as barriers to not being fully vaccinated.

Demographic characteristics

Approximately two thirds of ANSPS respondents were male across all years 2018 to 2022. In 2022, the majority (79%) of respondents identified as heterosexual, while 10% identified as bisexual and 5% as homosexual (Table 1.1.1). Women were significantly more likely to identify as bisexual than men in each of the past five years (18% vs 5% respectively in 2022, $p < 0.001$).

Over the period 2018 to 2022 the proportion of respondents from an Aboriginal and Torres Strait Islander background increased significantly from 19% to 25% (χ^2 trend $p < 0.001$)^{2,3}. In all years 2018 to 2022, the majority of respondents were born in Australia (range 86% to 88%), with the United Kingdom/Ireland (3%) and New Zealand (3%) the predominant countries of birth outside Australia in 2022. Most respondents also reported that their parents spoke English at home (range 93% to 95%) over the period 2018 to 2022.

Over the period 2018 to 2022, the median age at survey completion ranged from 42 to 45 years while the proportion of respondents aged <25 years (3% in 2022) decreased significantly (χ^2 trend $p = 0.001$, Table 1.1.1). The median time since first injection ranged from 22 to 24 years and the proportion of new initiates (those injecting for less than 3 years) was stable across the period 2018 to 2022, (range 4% to 6%, χ^2 trend $p = 0.980$). The median age at first drug injection was stable across the period 2018 to 2022, (19 years of age in all years). Among new initiates to injection drug use, the median age at first drug injection remained stable (range 29 to 34 years).

The proportion of respondents reporting incarceration in the 12 months prior to survey completion decreased significantly, from 13% to 11% over the five-year period 2018 to 2022 (χ^2 trend $p = 0.014$). Among respondents with a history of recent incarceration, approximately one in three reported

injecting while in prison and this proportion was stable over the period 2018 to 2022 (χ^2 trend $p=0.906$, Table 1.1.1).

Injection behaviour


Methamphetamine was the most commonly reported drug last injected nationally in all years 2018 to 2022 (Table 1.1.2). Reports of methamphetamine injection were stable over the five-year period 2018 to 2022 (range 47% to 51%, χ^2 trend $p=0.910$, Figure 1). Heroin was the second most commonly reported drug last injected nationally in 2022, with a significant decline in prevalence observed between 2018 (26%) and 2022 (25%, χ^2 trend $p=0.005$).

The third most commonly reported drug last injected nationally was performance and image enhancing drugs (PIEDs) which increased significantly over the five-year period, from 4% in 2018 to 7% in 2022 (χ^2 trend $p<0.001$, Figure 2). A significant increase in reports of PIEDs as the last drug injected was also observed in NSW (χ^2 trend $p=0.010$), QLD (χ^2 trend $p=0.001$) and WA (χ^2 trend $p=0.001$).

The prevalence of pharmaceutical opioids^{4,5} as the last drug injected (5% in 2022, including morphine, oxycodone and fentanyl) decreased over the period 2018 to 2022 (χ^2 trend $p=0.039$, Figure 2).

Methadone was reported as the last drug injected by less than ten percent of ANSPS respondents in all years 2018 to 2022 (range 6% to 4%) and prevalence remained stable over this period (χ^2 trend $p=0.617$). Buprenorphine and buprenorphine-naloxone injection were reported by a minority of survey respondents. The prevalence of both buprenorphine-naloxone injection (1% to 2%, χ^2 trend $p=0.246$), and buprenorphine injection (2% to 4%, χ^2 trend $p=0.095$) were stable over the period 2018 to 2022.


Figure 1. Proportion of respondents (%) reporting last injecting heroin, methamphetamine, and all other drugs, 2018–2022



A minority of survey respondents (1% across all survey years) reported cocaine as the last drug injected.

In 2022, for the first time, PIEDs was the most commonly reported drug last injected among new initiates (less than 3 years since first injection) with a significant increase observed between 2018 (18%) and 2022 (46%, χ^2 trend $p < 0.001$). In all years 2018 to 2021, methamphetamine was the most commonly reported last drug injected among new initiates (less than 3 years since first injection), with a significant decrease observed between 2018 (63%) and 2022 (38%, χ^2 trend $p = 0.001$).


Figure 2. Proportion of respondents (%) reporting last injecting pharmaceutical opioids, methadone, and performance and image enhancing drugs (PIEDs), 2018–2022



Daily or more frequent injection was reported by approximately half of respondents in each of the past five years (Table 1.1.2). Prevalence of at least one public injection (injection in a car, beach, park, street or squat) in the month prior to survey completion was stable between 2018 and 2022 (range 42% to 47%, χ^2 trend $p = 0.479$, Table 1.1.3). In 2022, one in eight respondents reported they had experienced a drug overdose in the previous 12 months ($n = 220$, 12%, Table 1.1.2), and reports of overdose declined significantly (χ^2 trend $p < 0.001$) over the period 2018 (19%) to 2022 (12%)⁶.

As shown in Figure 3, all three key injecting risk behaviour indicators were stable over the five-year period 2018 to 2022. The proportion of respondents who reported reuse of needles and syringes (including one's own) in the month preceding the survey was stable over the past 5 years (range 23% to 26%, χ^2 trend $p = 0.395$). Reports of receptive sharing of needles and syringes in the month prior to survey completion were also stable (range 16% to 19%, χ^2 trend $p = 0.428$) over the period 2018 to 2022, as were reports of receptive sharing of ancillary equipment (range 25% to 31%, χ^2 trend $p = 0.978$). As in previous years, spoons and water were the most commonly identified receptively shared ancillary items in 2022 (Table 1.1.3).

Figure 3. Proportion of respondents (%) reporting reuse and receptive sharing of needles and syringes and receptive sharing of ancillary equipment in the last month, 2018–2022



Sexual behaviour

In all years 2018 to 2022 approximately half of respondents reported sex with a regular partner in the month prior to survey completion (range 45% to 43%, Table 1.1.6). Of these, the majority (72% to 78%) reported that they never use condoms with their regular sex partner. A smaller proportion of respondents (14% to 16%) reported sex with other partners in the month preceding survey participation. Condom use was higher among this group, with 18% reporting always using condoms and 25% reporting some condom use with non-regular sex partners in 2022. The proportion of respondents reporting sex work in the month prior to the survey remained low and stable at 5% to 6% over the last five years (χ^2 trend $p=0.893$), and condom use at last paid sex was also stable, (range 44% to 70%, χ^2 trend $p=0.740$, Table 1.1.6). Of the respondents who reported sex work in the previous month in 2022 ($n=100$), 52% were women, 44% were men, and 3% identified their sex as other.

Drug treatment

Engagement with drug treatment services was high (Table 1.1.4) with two thirds (67%) of respondents in 2022 reporting a lifetime history of drug treatment. Nonetheless, there was a significant decline in the proportion of respondents reporting current OAT from 31% in 2018 to 28% in 2022 (χ^2 trend $p<0.001$). This may reflect the high proportion of respondents reporting methamphetamine as the last drug injected over this period (~50% in all years), with OAT not indicated for methamphetamine dependence. Among respondents who reported last injecting opioids, current OAT remained stable over the period 2018 to 2022 (range 44% to 53%, χ^2 trend $p=0.641$). In 2022, current methadone treatment was reported by 19% of respondents, with current buprenorphine treatment reported by 8% of respondents and current buprenorphine-naloxone treatment by 6% of respondents⁷.

HIV and HCV diagnostic testing

Prevalence of HIV diagnostic screening among ANSPS respondents over the period 2018 to 2022 was high, with 76% to 81% reporting a lifetime history of HIV diagnostic testing (Table 1.1.5). In 2022, 43% of respondents reported an HIV diagnostic test in the previous 12 months. Among the n=37 respondents who tested HIV antibody positive, the majority (78%, n=29) were aware of their HIV positive status. A minority (14%, n=5) of respondents who tested HIV positive reported that they were HIV negative, and 8% (n=3) did not report their HIV status.

Respondents also reported high prevalence of lifetime HCV diagnostic screening over the period 2018 to 2022 (range 78% to 82%). In 2022, 49% of respondents reported an HCV diagnostic test in the previous 12 months, however a significant decline in recent (last 12 months) HCV diagnostic testing was observed over the period 2018 (53%) to 2022 (49%, χ^2 trend $p < 0.001$).

HIV antibody prevalence

HIV antibody prevalence was low and stable nationally over the period 2018 to 2022 (range 1.5% to 2.5%, χ^2 trend $p = 0.853$, Table 1.2.1). HIV prevalence was low and stable in all jurisdictions (Figure 4). No women in the Australian Capital Territory, South Australia or Tasmania tested positive to HIV antibodies in any of the five years 2018 to 2022.

Nationally, HIV antibody prevalence was consistently higher among homosexual male respondents than among bisexual and heterosexual male or female respondents in all years, (Table 1.2.2 and Figure 5)⁸. In 2022, the median age of respondents living with HIV was 47.5 years (range 25 to 66 years) and the median time since first injection was 24 years (range 2 to 48 years). In 2022, comparable to previous years, 73% of respondents living with HIV reported they were on antiretroviral treatment.

Figure 4. HIV antibody prevalence (%) by jurisdiction, 2018–2022




Figure 5. HIV antibody prevalence (%) by gender and sexual identity, 2018–2022



Although respondents from an Aboriginal and Torres Strait Islander background were significantly more likely to test HIV antibody positive than respondents who were not from an Aboriginal and Torres Strait Islander background in 2018 and 2019 ($p < 0.05$), this difference was not statistically significant in 2020, 2021 or 2022 (Table 1.2.11). Further, prevalence of HIV infection among Aboriginal and Torres Strait Islander respondents remained stable between 2018 (3.6%) and 2022 (2.3%, χ^2 trend $p = 0.067$). Of the ten Aboriginal and Torres Strait Islander respondents who tested HIV antibody positive in 2022, half ($n = 5$) identified as homosexual ($n = 2$ bisexual, $n = 1$ heterosexual, $n = 1$ other term and $n = 1$ not reported).

HCV antibody prevalence

HCV antibody prevalence nationally declined significantly from 45% in 2018 to 32% in 2022 (χ^2 trend $p < 0.001$, Table 1.3.1). Over the five-year period 2018 to 2022, HCV antibody prevalence declined among both men (from 46% in 2018 to 31% in 2022, χ^2 trend $p < 0.001$) and women (from 43% in 2018 to 34% in 2022, χ^2 trend $p < 0.001$).

Figure 6. HCV antibody prevalence (%) by jurisdiction, 2018–2022






Figure 7. HCV antibody prevalence (%) by time since first injection, 2022



At the jurisdictional level (Figure 6), declines in HCV antibody prevalence were observed in most states and territories. Across all five years of the survey, HCV antibody prevalence was significantly higher among older respondents and those with longer injection histories (Table 1.3.3 and 1.3.4). Figure 7 illustrates the relationship between exposure to HCV and time since first injection among respondents in 2022.

In all survey years, prevalence of HCV antibody was also higher among respondents reporting heroin or other opioids as the last drug injected, compared to respondents who last injected methamphetamine (Figure 8). HCV antibody prevalence was also consistently higher among respondents who reported imprisonment in the year preceding survey completion compared to those who did not ($p < 0.010$ in all years 2018 to 2022) and among respondents who reported an Aboriginal and Torres Strait Islander background ($p \leq 0.001$ in all years 2018 to 2022).

Figure 8. HCV antibody prevalence (%) by drug last injected and gender, 2018–2022




HCV RNA prevalence

The ANSPS commenced HCV RNA testing in 2015, immediately before HCV direct acting antiviral (DAA) therapies were made available through the Australian Government’s Pharmaceutical Benefits Scheme (PBS) in March 2016⁹. The proportion of the ANSPS sample with sufficient DBS for HCV RNA testing has improved over time, with ~90% of respondents tested for HCV RNA in 2020, 2021 and 2022. In order to adjust for potential sample bias in the RNA tested sub-sample in previous years (2018-2019), weights were applied to adjust for gender (given higher rates of spontaneous clearance among women)¹⁰ and HCV antibody status. HCV RNA results with small sample sizes, particularly at the jurisdictional level, should be interpreted with caution.

As previously stated, over the last five years HCV antibody prevalence declined significantly (from 45% in 2018 to 32% in 2022, χ^2 trend $p < 0.001$). Lifetime HCV treatment increased significantly (from 55% in 2018 to 68% in 2022, χ^2 trend $p < 0.001$) and the proportion of respondents with


detectable HCV RNA decreased significantly, from 20% in 2018 to 12% in 2022 (χ^2 trend $p < 0.001$). As shown in Figure 9, HCV RNA prevalence declined significantly among both males (21% in 2018 to 13% in 2022, χ^2 trend $p < 0.001$) and females (19% in 2018 to 11% in 2022, χ^2 trend $p < 0.001$).

Figure 9. Proportion of respondents (%) with detectable HCV RNA* by gender and survey year




* Weighted for gender and HCV antibody status in 2018-2019

Figure 10. HCV viraemic status (%) by survey year



Results of HCV antibody and HCV RNA testing can be combined to provide an estimate of viraemic prevalence among respondents¹¹. As shown in Figure 10, the proportion of respondents with cleared infection (HCV antibody positive and HCV RNA negative) remained relatively stable (range 27% in 2019 to 20% in 2021 and 2022), while the proportion of respondents with no evidence of exposure to HCV (HCV antibody negative) increased from 55% in 2018 to 68% in 2022. Over the five-year period, 2018 to 2022, the proportion of respondents with active HCV infection declined by two-fifths (40%).


Figure 11. HCV viraemic status (%) nationally and by jurisdiction in 2022



HCV treatment


Among respondents who tested HCV antibody positive, and after excluding those who reported spontaneous HCV clearance, the proportion who reported a lifetime history of HCV treatment increased significantly from 55% in 2018 to 68% in 2022 (χ^2 trend $p=0.001$). Notwithstanding this overall increase in lifetime HCV treatment uptake, as shown in Figure 12, the significant increase occurred between 2018 (55%) and 2019 (64%, χ^2 trend $p=0.001$)^{11,12} with lifetime treatment uptake stable between 2019 (64%) and 2022 (68%, χ^2 trend $p=0.443$). These trends in lifetime treatment uptake were observed among both men and women. Conversely, among respondents who tested HCV antibody positive, and after excluding those who reported spontaneous or treatment-induced clearance more than 12 months previously, the proportion who reported recent (last 12 months) HCV treatment was stable between 2018 (39%) and 2019 (44%, χ^2 trend $p=0.088$) but declined significantly between 2019 and 2022 (34%, χ^2 trend $p<0.001$, Figure 12). These trends in recent treatment uptake were observed among both men and women¹³.

Figure 12. Proportion of respondents (%) reporting lifetime and recent* HCV treatment among HCV antibody positive respondents who did not report spontaneous clearance by survey year



* The denominator for recent HCV treatment excludes those who reported prior HCV treatment induced clearance.

Figure 13. Lifetime history of HCV treatment (%) nationally and by jurisdiction, 2018-2022



References

1. Iversen J, Wand H, Kemp R, Bevan J, Briggs M, Patten K, Heard S, Maher L. Uptake of COVID-19 vaccination among people who inject drugs. *Harm Reduction Journal* 2022;19 (59).
2. Doyle M, Maher L, Graham S, Wand H, Iversen J. Hepatitis C virus prevalence and associated risk factors among Indigenous Australians who inject drugs. *Australia New Zealand Journal of Public Health* 2018;42(1):52-6.
3. Graham S, Maher L, Wand H, Doyle M, Iversen J. Trends in hepatitis C antibody prevalence among Aboriginal and Torres Strait Islander people attending Australian Needle and Syringe Programs, 1996-2015. *International Journal of Drug Policy* 2017;47:69-76.
4. Iversen J, Dertadian G, Geddes L, Maher L. High risk injecting behaviour among people who inject pharmaceutical opioids in Australia. *International Journal of Drug Policy* 2017;42:1-6.
5. Geddes L, Iversen J, Memedovic S, Maher L. Intravenous fentanyl use among people who inject drugs in Australia. *Drug and Alcohol Review* 2018;37 Suppl 1:S314-S22.
6. Geddes L, Iversen J, Darke S, Dietze P, Maher L. Prevalence and correlates of multiple non-fatal opioid overdoses among people who inject drugs who utilise needle syringe programs in Australia. *International Journal of Drug Policy* 2021;96:103245
7. Geddes L, Iversen J, Wand H, Maher L. Incidence and factors associated with discontinuation of opioid agonist therapy among people who inject drugs in Australia. *Addiction* 2021;116(3):525-535.
8. Iversen J, Wand H, Topp L, Kaldor J, Maher L. Extremely low and sustained HIV incidence among people who inject drugs in a setting of harm reduction. *AIDS* 2014;28(2):275-8.
9. Iversen J, Dore G, Starr M, Catlett B, Cunningham P, Geddes L, Maher L. Estimating the Consensus Hepatitis C Cascade of Care among people who inject drugs in Australia: Pre and post availability of direct acting antiviral therapy. *International Journal of Drug Policy* 2020;83:102837.
10. Geddes L, Iversen J, Wand H, Esmaili A, Tsui J, Hellard M, Dore G, Grebely J, Dietze P, Prins M, Morris M, Shoukry N, Lloyd A, Lauer G, Cox A, Page K, Maher L. Sex discrepancies in the protective effect of opioid agonist therapy on incident hepatitis C infection. *Clinical Infectious Diseases* 2020;70(1):123-131.
11. Iversen J, Dore G, Catlett B, Cunningham P, Grebely J, Maher L. Association between rapid utilisation of direct hepatitis C antivirals and decline in the prevalence of viremia among people who inject drugs in Australia. *Journal of Hepatology* 2019;70(1):33-39.
12. Iversen J, Wand H, McManus H, Dore G, Maher L. Incidence of primary hepatitis C virus infection among people who inject drugs in Australia pre- and post-unrestricted availability of direct acting antiviral therapies. *Addiction* 2023;118(5):901-911.
13. Iversen J, Wand H, Chan PL, Le LV, Maher L. Systematic review of hepatitis C virus prevalence in the WHO Western Pacific Region. *Viruses* 2022;14(7):1548.

National

Table 1.1.1 Number (percentage) of respondents by demographic characteristics and survey year

| National | 2018 | 2019 | 2020* | 2021* | 2022 |
|---|-----------|-----------|-----------|-----------|-----------|
| Number of sites | 53 | 54 | 38 | 37 | 53 |
| N° surveyed | N=2742 | N=2609 | N=1324 | N=1474 | N=1781 |
| Response rate (%) | 45% | 45% | 35% | 45% | 34% |
| Gender (%) | | | | | |
| Male | 1828 (67) | 1722 (66) | 902 (68) | 933 (63) | 1156 (65) |
| Female | 899 (33) | 861 (33) | 413 (31) | 520 (35) | 586 (33) |
| Other | 13 (<1) | 18 (1) | 8 (1) | 19 (1) | 35 (2) |
| Not reported | 2 (<1) | 8 (<1) | 1 (<1) | 2 (<1) | 4 (<1) |
| Sexual identity (%) | | | | | |
| Heterosexual | 2133 (78) | 1969 (75) | 1045 (79) | 1204 (82) | 1409 (79) |
| Bisexual | 259 (9) | 262 (10) | 128 (10) | 149 (10) | 178 (10) |
| Gay/Lesbian | 122 (4) | 119 (5) | 50 (4) | 65 (4) | 87 (5) |
| Other | -- -- | -- -- | 35 (3) | 45 (3) | 93 (5) |
| Not reported | 228 (8) | 259 (10) | 66 (5) | 11 (1) | 14 (1) |
| Aboriginal and Torres Strait Islander origin (%) | | | | | |
| No | 2135 (78) | 1994 (76) | 982 (74) | 1031 (70) | 1312 (74) |
| Yes | 533 (19) | 568 (22) | 290 (22) | 373 (25) | 438 (25) |
| Not reported | 74 (3) | 47 (2) | 52 (4) | 70 (5) | 31 (2) |
| Region/country of birth (%) | | | | | |
| Australia | 2347 (86) | 2302 (88) | 1142 (86) | 1274 (86) | 1541 (87) |
| Other Oceania | 83 (3) | 67 (3) | 42 (3) | 43 (3) | 50 (3) |
| Asia | 31 (1) | 26 (1) | 18 (1) | 12 (1) | 26 (1) |
| UK & Ireland | 122 (4) | 87 (3) | 52 (4) | 57 (4) | 58 (3) |
| Other | 120 (4) | 102 (4) | 49 (4) | 48 (3) | 72 (4) |
| Not reported | 39 (1) | 25 (1) | 21 (2) | 40 (3) | 34 (2) |
| Main language spoken at home by parents (%) | | | | | |
| English | 2538 (93) | 2465 (94) | 1253 (95) | 1406 (95) | 1669 (94) |
| Non-English | 168 (6) | 128 (5) | 59 (4) | 51 (3) | 93 (5) |
| Not reported | 36 (1) | 16 (1) | 12 (1) | 17 (1) | 19 (1) |
| Age and time since first injection (years) | | | | | |
| <i>Median age</i> | 42 | 42 | 43 | 44 | 45 |
| <i>Age range</i> | 14-78 | 15-73 | 18-74 | 16-78 | 16-73 |
| Age group (%) | | | | | |
| <25 years | 116 (4) | 95 (4) | 44 (3) | 34 (2) | 52 (3) |
| 25+ years | 2617 (95) | 2502 (96) | 1279 (97) | 1431 (97) | 1711 (96) |
| Not reported | 9 (<1) | 12 (<1) | 1 (<1) | 9 (1) | 18 (1) |
| Median age first injection | | | | | |
| <i>Median age first injection</i> | 19 | 19 | 19 | 19 | 19 |
| <i>Age range</i> | 10-64 | 10-67 | 10-60 | 10-85 | 10-72 |
| <i>N° not reported</i> | 128 | 135 | 33 | 49 | 99 |
| Median yrs since first injection | | | | | |
| <i>Median yrs since first injection</i> | 22 | 22 | 22 | 23 | 24 |
| <i>Range</i> | <1-53 | <1-59 | <1-53 | <1-62 | <1-56 |
| Years since first injection | | | | | |
| <3 years | 137 (5) | 149 (6) | 82 (6) | 65 (4) | 97 (5) |
| 3+ years | 2475 (90) | 2321 (89) | 1208 (91) | 1353 (92) | 1584 (89) |
| Not reported | 130 (5) | 139 (5) | 34 (3) | 56 (4) | 100 (6) |
| Imprisonment last year (%) | | | | | |
| No | 2208 (81) | 2106 (81) | 1095 (83) | 1230 (83) | 1508 (85) |
| Yes | 346 (13) | 350 (13) | 158 (12) | 157 (11) | 198 (11) |
| Not reported | 188 (7) | 153 (6) | 71 (5) | 87 (6) | 75 (4) |
| N° in prison | | | | | |
| <i>N° in prison</i> | N=346 | N=350 | N=158 | N=157 | N=198 |
| <i>Injected in prison</i> | 116 (34) | 111 (32) | 48 (30) | 54 (34) | 66 (33) |

* Implementation of the 2020 and 2021 ANSPS was impacted by the COVID-19 pandemic, see page 1 for more information

Table 1.1.2 Number (percentage) of respondents by last drug injected, frequency of injecting and survey year

| National | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|---------------|---------------|---------------|---------------|---------------|
| N° surveyed | N=2742 | N=2609 | N=1324 | N=1474 | N=1781 |
| Drug last injected (%) | | | | | |
| Cocaine | 28 (1) | 20 (1) | 7 (1) | 8 (1) | 14 (1) |
| Methamphetamine | 1314 (48) | 1269 (49) | 619 (47) | 750 (51) | 840 (47) |
| Heroin | 714 (26) | 707 (27) | 286 (22) | 299 (20) | 451 (25) |
| Pharm. opioids | 204 (7) | 153 (6) | 109 (8) | 110 (7) | 87 (5) |
| Methadone | 112 (4) | 124 (5) | 77 (6) | 70 (5) | 75 (4) |
| Buprenorphine | 63 (2) | 43 (2) | 39 (3) | 53 (4) | 40 (2) |
| Buprenorphine/naloxone | 39 (1) | 23 (1) | 17 (1) | 33 (2) | 23 (1) |
| PIEDs | 100 (4) | 93 (4) | 84 (6) | 52 (4) | 118 (7) |
| More than one | 137 (5) | 119 (5) | 64 (5) | 77 (5) | 98 (6) |
| Other | 18 (1) | 37 (1) | 5 (<1) | 11 (1) | 17 (1) |
| Not reported | 13 (<1) | 21 (1) | 17 (1) | 11 (1) | 18 (1) |
| Ever injected opioids | | | | | |
| No | 627 (23) | 603 (23) | 377 (28) | 434 (29) | 472 (27) |
| Yes | 2078 (76) | 1989 (76) | 946 (71) | 1030 (70) | 1302 (73) |
| Not reported | 37 (1) | 17 (1) | 1 (<1) | 10 (1) | 7 (<1) |
| Frequency of injection last month (%) | | | | | |
| Not last month | 228 (8) | 235 (9) | 138 (10) | 186 (13) | 165 (9) |
| Less than weekly | 387 (14) | 371 (14) | 228 (17) | 293 (20) | 286 (16) |
| Weekly not daily | 669 (24) | 586 (22) | 300 (23) | 312 (21) | 393 (22) |
| Daily or more | 1396 (51) | 1376 (53) | 645 (49) | 654 (44) | 902 (51) |
| Not reported | 62 (2) | 41 (2) | 13 (1) | 29 (2) | 35 (2) |
| Present when someone injected for the first time in the last 12 months (%) | | | | | |
| No | 2157 (79) | 2056 (79) | 1088 (82) | 1207 (82) | 1459 (82) |
| Yes | 475 (17) | 448 (17) | 206 (16) | 234 (16) | 286 (16) |
| Not reported | 110 (4) | 105 (4) | 30 (2) | 33 (2) | 36 (2) |
| Overdosed in last 12 months | | | | | |
| No | 2106 (77) | 2060 (79) | 1120 (85) | 1265 (86) | 1525 (86) |
| Yes | 520 (19) | 438 (17) | 171 (13) | 166 (11) | 220 (12) |
| Not reported | 116 (4) | 111 (4) | 33 (2) | 43 (3) | 36 (2) |

Table 1.1.3 Number (percentage) of respondents by injecting behaviour in the last month and survey year

| National | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|-------------|-------------|-------------|-------------|-------------|
| N° injected last month | N=2452 | N=2333 | N=1173 | N=1259 | N=1581 |
| Places injected last month (%) | | | | | |
| All private | 1307 (53) | 1214 (52) | 673 (57) | 694 (55) | 841 (53) |
| Any public | 1129 (46) | 1105 (47) | 496 (42) | 559 (44) | 734 (46) |
| Not reported | 16 (1) | 14 (1) | 4 (<1) | 6 (<1) | 6 (<1) |
| Use of new and sterile needles and syringes last month (%) | | | | | |
| All injections | 1795 (73) | 1746 (75) | 891 (76) | 941 (75) | 1179 (75) |
| Most of the time | 502 (20) | 449 (19) | 220 (19) | 244 (19) | 312 (20) |
| Half of the time | 60 (2) | 53 (2) | 23 (2) | 27 (2) | 39 (2) |
| Some of the time | 44 (2) | 39 (2) | 19 (2) | 28 (2) | 23 (1) |
| Not last month | 20 (1) | 13 (1) | 4 (<1) | 10 (1) | 9 (1) |
| Not reported | 31 (1) | 33 (1) | 16 (1) | 9 (1) | 19 (1) |
| Re-used someone else's used needle & syringe last month (%) | | | | | |
| None | 1977 (81) | 1924 (82) | 980 (84) | 1030 (82) | 1269 (80) |
| Once | 164 (7) | 126 (5) | 66 (6) | 61 (5) | 93 (6) |
| Twice | 92 (4) | 113 (5) | 40 (3) | 61 (5) | 95 (6) |
| 3-5 times | 104 (4) | 82 (4) | 51 (4) | 44 (3) | 61 (4) |
| >5 times | 89 (4) | 57 (2) | 28 (2) | 54 (4) | 56 (4) |
| Not reported | 26 (1) | 31 (1) | 8 (1) | 9 (1) | 7 (<1) |
| Number of people needle & syringe was re-used after last month (%) | | | | | |
| None | 1977 (81) | 1924 (82) | 980 (84) | 1029 (82) | 1269 (80) |
| One | 194 (8) | 168 (7) | 73 (6) | 94 (7) | 125 (8) |
| Two | 36 (1) | 38 (2) | 20 (2) | 30 (2) | 38 (2) |
| Three to five | 34 (1) | 27 (1) | 14 (1) | 14 (1) | 21 (1) |
| More than five | 31 (1) | 17 (1) | 10 (1) | 14 (1) | 19 (1) |
| Don't know | 85 (3) | 79 (3) | 46 (4) | 50 (4) | 54 (3) |
| Not reported | 95 (4) | 80 (3) | 30 (3) | 28 (2) | 55 (3) |
| Relationship to people needle & syringe was used after last month (%) [more than one could be selected] | | | | | |
| Regular sex partner | 172 (7) | 147 (6) | 73 (6) | 77 (6) | 111 (7) |
| Casual sex partner | 21 (1) | 18 (1) | 14 (1) | 12 (1) | 13 (1) |
| Close friend | 131 (5) | 82 (4) | 48 (4) | 67 (5) | 86 (5) |
| Acquaintance | 55 (2) | 58 (2) | 30 (3) | 42 (3) | 40 (3) |
| Other | 38 (2) | 47 (2) | 19 (2) | 17 (1) | 28 (2) |
| Equipment used after someone else last month (%) [more than one could be selected] | | | | | |
| Spoon | 513 (21) | 482 (21) | 182 (16) | 252 (20) | 292 (18) |
| Water | 484 (20) | 451 (19) | 191 (16) | 284 (23) | 288 (18) |
| Filter | 243 (10) | 221 (9) | 71 (6) | 123 (10) | 122 (8) |
| Drug mix | 251 (10) | 234 (10) | 89 (8) | 127 (10) | 149 (9) |
| None | 1644 (67) | 1555 (67) | 836 (71) | 834 (66) | 1071 (68) |
| Injected by someone after that person injected themselves or others last month (%) | | | | | |
| No | 1980 (81) | 1939 (83) | 1005 (86) | 1030 (82) | 1276 (81) |
| Yes | 431 (18) | 369 (16) | 159 (14) | 219 (17) | 288 (18) |
| Not reported | 41 (2) | 25 (1) | 9 (1) | 10 (1) | 17 (1) |
| Source of needle acquisition | | | | | |
| Needle Syringe Program | 2140 (87) | 2032 (87) | 1001 (85) | 1071 (85) | 1341 (85) |
| Chemist/Pharmacy | 588 (24) | 572 (25) | 319 (27) | 327 (26) | 339 (21) |
| Personal sources | 306 (12) | 325 (14) | 130 (11) | 143 (11) | 190 (12) |
| Dispensing/Vending Machine | 400 (16) | 441 (19) | 172 (15) | 134 (11) | 234 (15) |
| Other sources | 100 (4) | 69 (3) | 37 (3) | 27 (2) | 43 (3) |

Table 1.1.4 Number (percentage) of respondents by treatment for drug use and survey year

| National | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|-----------|-----------|----------|-----------|-----------|
| N° surveyed | N=2742 | N=2609 | N=1324 | N=1474 | N=1781 |
| Ever any treatment/therapy for drug use (%) | | | | | |
| No | 810 (30) | 809 (31) | 460 (35) | 536 (36) | 580 (33) |
| Yes | 1901 (69) | 1789 (69) | 862 (65) | 937 (64) | 1198 (67) |
| Not reported | 31 (1) | 11 (<1) | 2 (<1) | 1 (<1) | 3 (<1) |
| History of methadone maintenance treatment (%) | | | | | |
| Currently | 583 (21) | 634 (24) | 235 (18) | 219 (15) | 345 (19) |
| Previously | 645 (24) | 522 (20) | 236 (18) | 281 (19) | 399 (22) |
| Never | 1411 (51) | 1395 (53) | 809 (61) | 921 (62) | 990 (56) |
| Not reported | 103 (4) | 58 (2) | 44 (3) | 53 (4) | 47 (3) |
| History of buprenorphine (Subutex®) treatment (%) | | | | | |
| Currently | 196 (7) | 169 (6) | 105 (8) | 119 (8) | 139 (8) |
| Previously | 703 (26) | 640 (25) | 278 (21) | 310 (21) | 422 (24) |
| Never | 1759 (64) | 1759 (67) | 903 (68) | 1001 (68) | 1175 (66) |
| Not reported | 84 (3) | 41 (2) | 38 (3) | 44 (3) | 45 (3) |
| History of buprenorphine-naloxone (Suboxone®) treatment (%) | | | | | |
| Currently | 212 (8) | 173 (7) | 70 (5) | 105 (7) | 101 (6) |
| Previously | 645 (24) | 596 (23) | 258 (19) | 298 (20) | 422 (24) |
| Never | 1799 (66) | 1795 (69) | 987 (75) | 1053 (71) | 1242 (70) |
| Not reported | 86 (3) | 45 (2) | 9 (1) | 18 (1) | 16 (1) |

Table 1.1.5 Number (percentage) of respondents by testing for HIV and HCV, HCV treatment by survey year

| National | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|-----------|-----------|-----------|-----------|-----------|
| N° surveyed | N=2742 | N=2609 | N=1324 | N=1474 | N=1781 |
| Previous HIV test (%) | | | | | |
| Yes, ever | 2211 (81) | 2105 (81) | 1042 (79) | 1116 (76) | 1360 (76) |
| Yes, last year | 1363 (50) | 1271 (49) | 601 (45) | 557 (38) | 768 (43) |
| >1 year ago | 848 (31) | 834 (32) | 441 (33) | 559 (38) | 592 (33) |
| Never tested | 390 (14) | 381 (15) | 232 (18) | 296 (20) | 357 (20) |
| Not reported | 141 (5) | 123 (5) | 50 (4) | 62 (4) | 64 (4) |
| Previous HCV test (%) | | | | | |
| Yes, ever | 2201 (80) | 2101 (81) | 1087 (82) | 1147 (78) | 1417 (80) |
| Yes, last year | 1455 (53) | 1409 (54) | 631 (48) | 686 (47) | 875 (49) |
| >1 year ago | 746 (27) | 692 (27) | 456 (34) | 462 (31) | 542 (30) |
| Never tested | 251 (9) | 257 (10) | 128 (10) | 194 (13) | 212 (12) |
| Not reported | 290 (11) | 251 (10) | 109 (8) | 132 (9) | 152 (9) |
| Lifetime treatment for HCV (%)[#] | | | | | |
| N° self-reported HCV diagnosis | N=738 | N=667 | N=289 | N=309 | N=334 |
| Antiviral treatment | 409 (55) | 428 (64) | 180 (62) | 191 (62) | 226 (68) |
| No antiviral treatment | 322 (44) | 237 (36) | 107 (37) | 116 (38) | 107 (32) |
| Not reported | 7 (1) | 2 (<1) | 2 (1) | 2 (1) | 1 (<1) |
| Treatment for HCV in past 12 months (%)^{**} | | | | | |
| N° self-reported HCV diagnosis | N=603 | N=502 | N=196 | N=200 | N=199 |
| Antiviral treatment | 236 (39) | 222 (44) | 62 (32) | 56 (28) | 68 (34) |
| No antiviral treatment | 360 (60) | 278 (55) | 132 (67) | 142 (71) | 130 (65) |
| Not reported | 7 (1) | 2 (<1) | 2 (1) | 2 (1) | 1 (1) |

[#] among people who tested HCV antibody positive and did not report spontaneous clearance

^{**} excludes people who reported treatment induced clearance more than 12 months previously

Table 1.1.6 Number (percentage) of respondents by sexual behaviour in the month prior to survey and survey year

| National | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|-------------|-------------|-------------|-------------|-------------|
| N° surveyed | N=2742 | N=2609 | N=1324 | N=1474 | N=1781 |
| Sex with a regular partner last month (%) | | | | | |
| No | 1374 (50) | 1345 (52) | 684 (52) | 758 (51) | 933 (52) |
| Yes | 1228 (45) | 1124 (43) | 584 (44) | 631 (43) | 774 (43) |
| Not reported | 140 (5) | 140 (5) | 56 (4) | 85 (6) | 74 (4) |
| Condom used with regular partner last month (%) | | | | | |
| N° surveyed | N=1228 | N=1124 | N=584 | N=631 | N=774 |
| Never | 924 (75) | 808 (72) | 450 (77) | 479 (76) | 604 (78) |
| Sometimes | 137 (11) | 150 (13) | 55 (9) | 68 (11) | 74 (10) |
| Every time | 95 (8) | 110 (10) | 51 (9) | 52 (8) | 61 (8) |
| Not reported | 72 (6) | 56 (5) | 28 (5) | 32 (5) | 35 (5) |
| Sex with other partner(s) last month (%) | | | | | |
| No | 2137 (78) | 2040 (78) | 1063 (80) | 1154 (78) | 1409 (79) |
| Yes | 420 (15) | 399 (15) | 190 (14) | 213 (14) | 283 (16) |
| Not reported | 185 (7) | 170 (7) | 71 (5) | 107 (7) | 89 (5) |
| Condom used with other partner(s) last month (%) | | | | | |
| N° surveyed | N=420 | N=399 | N=190 | N=213 | N=283 |
| Never | 179 (43) | 184 (46) | 90 (47) | 104 (49) | 152 (54) |
| Sometimes | 117 (28) | 100 (25) | 54 (28) | 47 (22) | 70 (25) |
| Every time | 105 (25) | 89 (22) | 41 (22) | 48 (23) | 52 (18) |
| Not reported | 19 (5) | 26 (7) | 5 (3) | 14 (7) | 9 (3) |
| Sex work last month (%) | | | | | |
| No | 2394 (87) | 2276 (87) | 1190 (90) | 1292 (88) | 1579 (89) |
| Yes | 154 (6) | 134 (5) | 61 (5) | 76 (5) | 100 (6) |
| Not reported | 194 (7) | 199 (8) | 73 (6) | 106 (7) | 102 (6) |
| Condom used at last sex work (%) | | | | | |
| Yes | 100 (65) | 78 (58) | 27 (44) | 43 (57) | 70 (70) |

Table 1.1.7 Number (percentage) of respondents by COVID-19 related impacts, 2020-2022

| National | 2020 | 2021 | 2022 |
|---|---------------|---------------|---------------|
| N° surveyed | N=1324 | N=1474 | N=1781 |
| COVID-19 test | | | |
| Yes | 337 (25) | 450 (31) | 1521 (85) |
| No | 802 (61) | 680 (46) | 213 (12) |
| Don't know | 60 (5) | 63 (4) | -- -- |
| Not reported | 125 (9) | 281 (19) | 47 (3) |
| Self-reported positive COVID-19 test result | | | |
| Yes | 4 (<1) | 16 (1) | 642 (42) |
| Yes, last 12 months | -- -- | -- -- | 546 (36) |
| Yes, more than a year ago | -- -- | -- -- | 96 (6) |
| COVID-19 vaccine uptake | | | |
| ≥1 dose | -- -- | 599 (41) | 1409 (79) |
| COVID-19 vaccine doses | | | |
| One dose | -- -- | 232 (16) | 126 (7) |
| Two doses | -- -- | 367 (25) | 496 (28) |
| Three or more doses | -- -- | -- -- | 787 (44) |
| No doses | -- -- | 635 (43) | 304 (17) |
| Not reported | -- -- | 240 (16) | 68 (4) |
| Vaccine type * | | | |
| | | | N=1409 |
| Astra Zeneca | -- -- | 139 (9) | 415 (29) |
| Pfizer | -- -- | 310 (21) | 913 (65) |
| Don't know | -- -- | 104 (7) | 197 (14) |
| Other | -- -- | -- -- | 121 (9) |
| Not reported | -- -- | 921 (62) | -- -- |
| * 2021 only one response permitted, 2022 multiple responses permitted | | | |
| Last vaccine location | | | |
| | | | N=1781 |
| GP or medical centre | -- -- | -- -- | 565 (32) |
| Pharmacy/ chemist | -- -- | -- -- | 304 (17) |
| Vaccine hub | -- -- | -- -- | 256 (14) |
| Opioid treatment/ methadone clinic | -- -- | -- -- | 30 (2) |
| NSP | -- -- | -- -- | 52 (3) |
| Prison | -- -- | -- -- | 91 (5) |
| Other | -- -- | -- -- | 40 (2) |
| Multiple response | -- -- | -- -- | 114 (6) |
| Not reported | -- -- | -- -- | 329 (18) |
| Vaccine hesitancy - if not had three or more doses | | | |
| Haven't got around to it | -- -- | -- -- | 344 (19) |
| Difficulty booking appointment | -- -- | -- -- | 37 (2) |
| Concerns about side effects | -- -- | -- -- | 97 (5) |
| Concerns about vaccine safety | -- -- | -- -- | 92 (5) |
| Not at risk of serious illness | -- -- | -- -- | 129 (7) |
| Other | -- -- | -- -- | 195 (11) |
| Multiple response | -- -- | -- -- | 85 (5) |
| Not reported | -- -- | -- -- | 802 (45) |

HIV antibody prevalence

Table 1.2.1 HIV antibody prevalence by gender and survey year

| National Survey year | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | 1784 | 37 (2.1) | 888 | 8 (0.9) | 2686 | 46 (1.7) |
| 2019 | 1675 | 51 (3.0) | 841 | 6 (0.7) | 2542 | 59 (2.3) |
| 2020 | 877 | 27 (3.1) | 406 | 3 (0.7) | 1292 | 32 (2.5) |
| 2021 | 902 | 16 (1.8) | 514 | 3 (0.6) | 1435 | 21 (1.5) |
| 2022 | 1132 | 28 (2.5) | 583 | 6 (1.0) | 1752 | 37 (2.1) |
| X ² p trend | | 0.949 | | 0.943 | | 0.853 |

Table 1.2.2 HIV antibody prevalence by sexual identity, gender and survey year

| National Sexual identity | Male | | Female | | Total | |
|-----------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Heterosexual | 1473 | 12 (0.8) | 611 | 6 (1.0) | 2086 | 18 (0.9) |
| Bisexual | 90 | 6 (6.7) | 162 | 2 (1.2) | 255 | 8 (3.1) |
| Homosexual | 80 | 16 (20.0) | 35 | 0 (0.0) | 120 | 17 (14.2) |
| p value | | <0.001 | | 0.773 | | <0.001 |
| 2019 | | | | | | |
| Heterosexual | 1355 | 13 (1.0) | 560 | 5 (0.9) | 1919 | 18 (0.9) |
| Bisexual | 92 | 13 (14.1) | 148 | 0 (0.0) | 253 | 14 (5.5) |
| Homosexual | 79 | 25 (31.7) | 36 | 0 (0.0) | 117 | 26 (22.2) |
| p value | | <0.001 | | 0.680 | | <0.001 |
| 2020 | | | | | | |
| Heterosexual | 746 | 5 (0.7) | 276 | 3 (1.1) | 1023 | 8 (0.8) |
| Bisexual | 43 | 4 (9.3) | 72 | 0 (0.0) | 120 | 5 (4.2) |
| Homosexual | 38 | 17 (44.7) | 10 | 0 (0.0) | 49 | 18 (36.7) |
| p value | | <0.001 | | 1.000 | | <0.001 |
| 2021 | | | | | | |
| Heterosexual | 793 | 4 (0.5) | 375 | 2 (0.5) | 1172 | 7 (0.6) |
| Bisexual | 49 | 0 (0.0) | 96 | 0 (0.0) | 148 | 0 (0.0) |
| Homosexual | 40 | 11 (27.5) | 16 | 1 (6.3) | 61 | 13 (21.3) |
| p value | | <0.001 | | 0.201 | | <0.001 |
| 2022 | | | | | | |
| Heterosexual | 973 | 3 (0.3) | 407 | 4 (1.0) | 1387 | 7 (0.5) |
| Bisexual | 61 | 2 (3.3) | 107 | 1 (0.9) | 177 | 5 (2.8) |
| Homosexual | 58 | 22 (37.9) | 24 | 0 (0.0) | 84 | 22 (26.2) |
| p value | | <0.001 | | 0.410 | | <0.001 |

Table 1.2.3 HIV antibody prevalence by age group, gender and survey year

| National Age group | Male | | Female | | Total | |
|-----------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| <25 years | 60 | 1 (1.7) | 49 | 0 (0.0) | 111 | 1 (0.9) |
| 25-34 years | 284 | 12 (4.2) | 194 | 2 (1.0) | 480 | 14 (2.9) |
| 35-44 years | 643 | 6 (0.9) | 338 | 1 (0.3) | 987 | 7 (0.7) |
| 45+ years | 791 | 18 (2.3) | 304 | 5 (1.6) | 1099 | 24 (2.2) |
| p value | | 0.013 | | 0.313 | | 0.005 |
| 2019 | | | | | | |
| <25 years | 51 | 0 (0.0) | 39 | 0 (0.0) | 91 | 0 (0.0) |
| 25-34 years | 254 | 8 (3.2) | 175 | 0 (0.0) | 436 | 9 (2.1) |
| 35-44 years | 605 | 11 (1.8) | 332 | 2 (0.6) | 943 | 13 (1.4) |
| 45+ years | 757 | 31 (4.1) | 294 | 4 (1.4) | 1062 | 36 (3.4) |
| p value | | 0.062 | | 0.457 | | 0.011 |
| 2020 | | | | | | |
| <25 years | 23 | 0 (0.0) | 18 | 1 (5.6) | 42 | 1 (2.4) |
| 25-34 years | 145 | 4 (2.8) | 78 | 1 (1.3) | 225 | 5 (2.2) |
| 35-44 years | 288 | 6 (2.1) | 160 | 0 (0.0) | 449 | 6 (1.3) |
| 45+ years | 420 | 17 (4.1) | 150 | 1 (0.7) | 575 | 20 (3.5) |
| p value | | 0.487 | | 0.054 | | 0.151 |
| 2021 | | | | | | |
| <25 years | 21 | 1 (4.8) | 12 | 0 (0.0) | 34 | 1 (2.9) |
| 25-34 years | 122 | 1 (0.8) | 83 | 0 (0.0) | 209 | 2 (1.0) |
| 35-44 years | 306 | 3 (1.0) | 194 | 2 (1.0) | 509 | 6 (1.2) |
| 45+ years | 449 | 11 (2.5) | 221 | 1 (0.5) | 675 | 12 (1.8) |
| p value | | 0.202 | | 0.785 | | 0.483 |
| 2022 | | | | | | |
| <25 years | 36 | 0 (0.0) | 12 | 0 (0.0) | 52 | 0 (0.0) |
| 25-34 years | 161 | 2 (1.2) | 84 | 0 (0.0) | 250 | 2 (0.8) |
| 35-44 years | 341 | 9 (2.6) | 213 | 3 (1.4) | 561 | 13 (2.3) |
| 45+ years | 583 | 16 (2.7) | 268 | 3 (1.1) | 871 | 21 (2.4) |
| p value | | 0.720 | | 0.880 | | 0.365 |

Table 1.2.4 HIV antibody prevalence by last drug injected, gender and survey year

| National Last drug injected | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Heroin | 460 | 8 (1.7) | 238 | 2 (0.8) | 701 | 10 (1.4) |
| Methamphetamine | 828 | 25 (3.0) | 452 | 6 (1.3) | 1287 | 32 (2.5) |
| Other opioids | 274 | 1 (0.4) | 137 | 0 (0.0) | 412 | 1 (0.2) |
| PIEDs | 90 | 0 (0.0) | 5 | 0 (0.0) | 96 | 0 (0.0) |
| Other drugs | 126 | 3 (2.4) | 53 | 0 (0.0) | 181 | 3 (1.7) |
| p value | | 0.035 | | 0.671 | | 0.012 |
| 2019 | | | | | | |
| Heroin | 429 | 9 (2.1) | 249 | 2 (0.8) | 689 | 12 (1.7) |
| Methamphetamine | 794 | 37 (4.7) | 427 | 3 (0.7) | 1231 | 41 (3.3) |
| Other opioids | 241 | 4 (1.7) | 96 | 0 (0.0) | 339 | 4 (1.2) |
| PIEDs | 88 | 0 (0.0) | 5 | 0 (0.0) | 93 | 0 (0.0) |
| Other drugs | 114 | 1 (0.9) | 55 | 1 (1.8) | 170 | 2 (1.2) |
| p value | | 0.008 | | 0.565 | | 0.030 |
| 2020 | | | | | | |
| Heroin | 191 | 2 (1.1) | 88 | 1 (1.1) | 280 | 3 (1.1) |
| Methamphetamine | 391 | 21 (5.4) | 207 | 2 (1.0) | 603 | 24 (4.0) |
| Other opioids | 161 | 2 (1.2) | 78 | 0 (0.0) | 241 | 2 (0.8) |
| PIEDs | 77 | 0 (0.0) | 3 | 0 (0.0) | 80 | 0 (0.0) |
| Other drugs | 50 | 1 (2.0) | 27 | 0 (0.0) | 78 | 2 (2.6) |
| p value | | 0.008 | | 1.000 | | 0.013 |
| 2021 | | | | | | |
| Heroin | 180 | 2 (1.1) | 106 | 1 (0.9) | 292 | 4 (1.4) |
| Methamphetamine | 419 | 11 (2.6) | 299 | 2 (0.7) | 725 | 14 (1.9) |
| Other opioids | 187 | 2 (1.1) | 72 | 0 (0.0) | 263 | 2 (0.8) |
| PIEDs | 51 | 0 (0.0) | 0 | 0 (0.0) | 51 | 0 (0.0) |
| Other drugs | 59 | 0 (0.0) | 33 | 0 (0.0) | 93 | 0 (0.0) |
| p value | | 0.515 | | 1.000 | | 0.564 |
| 2022 | | | | | | |
| Heroin | 262 | 1 (0.4) | 176 | 2 (1.1) | 443 | 4 (0.9) |
| Methamphetamine | 497 | 22 (4.4) | 308 | 4 (1.3) | 827 | 26 (3.1) |
| Other opioids | 163 | 1 (0.6) | 54 | 0 (0.0) | 223 | 1 (0.5) |
| PIEDs | 113 | 2 (1.8) | 4 | 0 (0.0) | 117 | 2 (1.7) |
| Other drugs | 90 | 2 (2.2) | 33 | 0 (0.0) | 126 | 3 (2.4) |
| p value | | 0.003 | | 1.000 | | 0.023 |

Table 1.2.5 HIV antibody prevalence by years of injection, gender and survey year

| National Years of injection | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| <3 years | 83 | 3 (3.6) | 46 | 0 (0.0) | 129 | 3 (2.3) |
| 3 to 10 years | 235 | 9 (3.8) | 148 | 0 (0.0) | 384 | 9 (2.3) |
| 11+ years | 1466 | 25 (1.7) | 694 | 8 (1.2) | 2173 | 34 (1.6) |
| p value | | 0.046 | | 0.584 | | 0.420 |
| 2019 | | | | | | |
| <3 years | 97 | 3 (3.1) | 49 | 0 (0.0) | 146 | 3 (2.1) |
| 3 to 10 years | 210 | 12 (5.7) | 145 | 1 (0.7) | 361 | 14 (3.9) |
| 11+ years | 1278 | 35 (2.7) | 608 | 5 (0.8) | 1091 | 41 (2.2) |
| p value | | 0.076 | | 1.000 | | 0.165 |
| 2020 | | | | | | |
| <3 years | 56 | 1 (1.8) | 26 | 0 (0.0) | 82 | 1 (1.2) |
| 3 to 10 years | 134 | 8 (6.0) | 61 | 2 (3.3) | 197 | 10 (5.1) |
| 11+ years | 665 | 18 (2.7) | 309 | 1 (0.3) | 981 | 21 (2.1) |
| p value | | 0.131 | | 0.123 | | 0.055 |
| 2021 | | | | | | |
| <3 years | 43 | 0 (0.0) | 19 | 0 (0.0) | 64 | 0 (0.0) |
| 3 to 10 years | 108 | 6 (5.6) | 76 | 0 (0.0) | 185 | 7 (3.8) |
| 11+ years | 719 | 9 (1.3) | 397 | 3 (0.8) | 1131 | 13 (1.2) |
| p value | | 0.015 | | 1.000 | | 0.035 |
| 2022 | | | | | | |
| <3 years | 69 | 1 (1.5) | 23 | 0 (0.0) | 96 | 1 (1.0) |
| 3 to 10 years | 127 | 2 (1.6) | 77 | 0 (0.0) | 207 | 2 (1.0) |
| 11+ years | 866 | 23 (2.7) | 456 | 6 (1.3) | 1349 | 32 (2.4) |
| p value | | 0.857 | | 0.690 | | 0.478 |

Table 1.2.6 HIV antibody prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| National Receptively shared syringe last month | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| No receptive sharing | 1309 | 29 (2.2) | 637 | 6 (0.9) | 1951 | 35 (1.8) |
| Receptive sharing | 293 | 5 (1.7) | 148 | 0 (0.0) | 444 | 5 (1.1) |
| p value | | 0.390 | | 0.601 | | 0.414 |
| 2019 | | | | | | |
| No receptive sharing | 1235 | 41 (3.3) | 625 | 3 (0.5) | 1877 | 46 (2.5) |
| Receptive sharing | 247 | 6 (2.4) | 115 | 2 (1.7) | 366 | 8 (2.2) |
| p value | | 0.466 | | 0.174 | | 0.762 |
| 2020 | | | | | | |
| No receptive sharing | 663 | 20 (3.0) | 284 | 3 (1.1) | 955 | 25 (2.6) |
| Receptive sharing | 107 | 2 (1.9) | 75 | 0 (0.0) | 182 | 2 (1.1) |
| p value | | 0.756 | | 0.494 | | 0.293 |
| 2021 | | | | | | |
| No receptive sharing | 638 | 14 (2.2) | 355 | 2 (0.6) | 1003 | 18 (1.8) |
| Receptive sharing | 135 | 1 (0.7) | 75 | 1 (1.3) | 215 | 2 (0.9) |
| p value | | 0.490 | | 0.438 | | 0.555 |
| 2022 | | | | | | |
| No receptive sharing | 814 | 22 (2.7) | 408 | 2 (0.5) | 1249 | 26 (2.1) |
| Receptive sharing | 189 | 3 (1.6) | 107 | 1 (0.9) | 303 | 4 (1.3) |
| p value | | 0.603 | | 0.504 | | 0.491 |

Table 1.2.7 HIV antibody prevalence by frequency of drug injection last month, gender and survey year

| National Frequency of injection | Male | | Female | | Total | |
|------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Less than daily | 717 | 13 (1.8) | 324 | 4 (1.2) | 1044 | 17 (1.6) |
| Daily or more | 899 | 21 (2.3) | 472 | 3 (0.6) | 1376 | 24 (1.7) |
| Not last month | 137 | 3 (2.2) | 82 | 1 (1.2) | 223 | 5 (2.2) |
| p value | | 0.760 | | 0.466 | | 0.817 |
| 2019 | | | | | | |
| Less than daily | 643 | 26 (4.0) | 291 | 2 (0.7) | 941 | 28 (3.0) |
| Daily or more | 856 | 21 (2.5) | 461 | 4 (0.9) | 1331 | 27 (2.0) |
| Not last month | 149 | 3 (2.0) | 78 | 0 (0.0) | 230 | 3 (1.3) |
| p value | | 0.172 | | 1.000 | | 0.204 |
| 2020 | | | | | | |
| Less than daily | 350 | 11 (3.1) | 159 | 2 (1.3) | 514 | 13 (2.5) |
| Daily or more | 425 | 11 (2.6) | 202 | 1 (0.5) | 630 | 14 (2.2) |
| Not last month | 93 | 5 (5.4) | 42 | 0 (0.0) | 136 | 5 (3.7) |
| p value | | 0.373 | | 0.702 | | 0.615 |
| 2021 | | | | | | |
| Less than daily | 373 | 10 (2.7) | 209 | 0 (0.0) | 588 | 10 (1.7) |
| Daily or more | 402 | 5 (1.2) | 228 | 3 (1.3) | 639 | 10 (1.6) |
| Not last month | 111 | 0 (0.0) | 67 | 0 (0.0) | 180 | 0 (0.0) |
| p value | | 0.118 | | 0.361 | | 0.199 |
| 2022 | | | | | | |
| Less than daily | 438 | 14 (3.2) | 214 | 1 (0.5) | 669 | 15 (2.2) |
| Daily or more | 570 | 11 (1.9) | 303 | 2 (0.7) | 890 | 15 (1.7) |
| Not last month | 105 | 3 (2.9) | 51 | 3 (5.9) | 158 | 6 (3.8) |
| p value | | 0.391 | | 0.020 | | 0.220 |

Table 1.2.8 HIV antibody prevalence by imprisonment last year, gender and survey year

| National Imprisonment last year | Male | | Female | | Total | |
|------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| No imprisonment | 1421 | 30 (2.1) | 748 | 7 (0.9) | 2179 | 38 (1.7) |
| Imprisonment | 249 | 7 (2.8) | 92 | 0 (0.0) | 343 | 7 (2.0) |
| p value | | 0.489 | | 1.000 | | 0.699 |
| 2019 | | | | | | |
| No imprisonment | 1322 | 43 (3.3) | 708 | 5 (0.7) | 2049 | 50 (2.4) |
| Imprisonment | 250 | 3 (1.2) | 89 | 1 (1.1) | 344 | 4 (1.2) |
| p value | | 0.099 | | 0.510 | | 0.170 |
| 2020 | | | | | | |
| No imprisonment | 711 | 23 (3.2) | 349 | 3 (0.9) | 1069 | 28 (2.6) |
| Imprisonment | 118 | 4 (3.4) | 38 | 0 (0.0) | 156 | 4 (2.6) |
| p value | | 1.000 | | 1.000 | | 0.614 |
| 2021 | | | | | | |
| No imprisonment | 733 | 13 (1.8) | 447 | 3 (0.7) | 1196 | 18 (1.5) |
| Imprisonment | 114 | 1 (0.9) | 39 | 0 (0.0) | 154 | 1 (0.7) |
| p value | | 0.707 | | 1.000 | | 0.714 |
| 2022 | | | | | | |
| No imprisonment | 938 | 28 (3.0) | 514 | 3 (0.6) | 1484 | 34 (2.3) |
| Imprisonment | 147 | 0 (0.0) | 48 | 2 (4.2) | 197 | 2 (1.0) |
| p value | | 0.024 | | 0.060 | | 0.427 |

Table 1.2.9 HIV antibody prevalence by condom use at last sex with regular partner, gender and survey year

| National Condom use at last sex | Male | | Female | | Total | |
|------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| No condom use | 545 | 11 (2.0) | 369 | 1 (0.3) | 917 | 12 (1.3) |
| Condom use | 154 | 7 (4.6) | 67 | 0 (0.0) | 226 | 7 (3.1) |
| p value | | 0.080 | | 1.000 | | 0.060 |
| 2019 | | | | | | |
| No condom use | 469 | 18 (3.8) | 317 | 0 (0.0) | 790 | 18 (2.3) |
| Condom use | 159 | 1 (0.6) | 89 | 3 (3.4) | 253 | 4 (1.6) |
| p value | | 0.056 | | 0.010 | | 0.621 |
| 2020 | | | | | | |
| No condom use | 278 | 13 (4.7) | 159 | 1 (0.6) | 439 | 15 (3.4) |
| Condom use | 75 | 3 (4.0) | 29 | 0 (0.0) | 104 | 3 (2.9) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2021 | | | | | | |
| No condom use | 267 | 4 (1.5) | 200 | 0 (0.0) | 473 | 4 (0.9) |
| Condom use | 65 | 0 (0.0) | 48 | 0 (0.0) | 116 | 1 (0.9) |
| p value | | 1.000 | | -- | | 1.000 |
| 2022 | | | | | | |
| No condom use | 353 | 10 (2.8) | 235 | 2 (0.9) | 598 | 13 (2.2) |
| Condom use | 83 | 1 (1.2) | 45 | 1 (2.2) | 133 | 3 (2.3) |
| p value | | 0.698 | | 0.410 | | 1.000 |

Table 1.2.10 HIV antibody prevalence by sex work last month, gender and survey year

| National Sex work last month | Male | | Female | | Total | |
|---------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| No sex work | 1614 | 31 (1.9) | 740 | 6 (0.8) | 2363 | 38 (1.6) |
| Sex work | 65 | 2 (3.1) | 81 | 0 (0.0) | 150 | 2 (1.3) |
| p value | | 0.368 | | 1.000 | | 1.000 |
| 2019 | | | | | | |
| No sex work | 1490 | 44 (3.0) | 717 | 5 (0.7) | 2226 | 51 (2.3) |
| Sex work | 60 | 4 (6.7) | 64 | 1 (1.6) | 129 | 5 (3.9) |
| p value | | 0.111 | | 0.402 | | 0.251 |
| 2020 | | | | | | |
| No sex work | 807 | 23 (2.9) | 352 | 2 (0.6) | 1166 | 26 (2.2) |
| Sex work | 26 | 4 (15.4) | 30 | 1 (3.3) | 58 | 6 (10.3) |
| p value | | 0.008 | | 0.218 | | <0.001 |
| 2021 | | | | | | |
| No sex work | 820 | 13 (1.6) | 425 | 3 (0.7) | 1259 | 16 (1.3) |
| Sex work | 27 | 1 (3.7) | 44 | 0 (0.0) | 74 | 3 (4.1) |
| p value | | 0.367 | | 1.000 | | 0.084 |
| 2022 | | | | | | |
| No sex work | 1018 | 22 (2.2) | 504 | 6 (1.2) | 1553 | 31 (2.0) |
| Sex work | 44 | 5 (11.4) | 52 | 0 (0.0) | 100 | 5 (5.0) |
| p value | | <0.001 | | 1.000 | | 0.046 |

Table 1.2.11 HIV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| National Aboriginal and Torres Strait Islander origin | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Non Indigenous | 1439 | 20 (1.4) | 662 | 3 (0.5) | 2111 | 23 (1.1) |
| Indigenous | 311 | 15 (4.8) | 209 | 3 (1.4) | 523 | 19 (3.6) |
| p value | | <0.001 | | 0.153 | | <0.001 |
| 2019 | | | | | | |
| Non Indigenous | 1314 | 37 (2.8) | 615 | 1 (0.2) | 1947 | 39 (2.0) |
| Indigenous | 331 | 14 (4.2) | 215 | 5 (2.3) | 552 | 20 (3.6) |
| p value | | 0.185 | | 0.005 | | 0.027 |
| 2020 | | | | | | |
| Non Indigenous | 675 | 20 (3.0) | 278 | 2 (0.7) | 958 | 23 (2.4) |
| Indigenous | 168 | 6 (3.6) | 115 | 1 (0.9) | 285 | 8 (2.8) |
| p value | | 0.683 | | 1.000 | | 0.699 |
| 2021 | | | | | | |
| Non Indigenous | 650 | 13 (2.0) | 347 | 1 (0.3) | 1008 | 15 (1.5) |
| Indigenous | 207 | 3 (1.5) | 145 | 2 (1.4) | 359 | 6 (1.7) |
| p value | | 0.773 | | 0.209 | | 0.804 |
| 2022 | | | | | | |
| Non Indigenous | 864 | 22 (2.6) | 399 | 4 (1.0) | 1287 | 27 (2.1) |
| Indigenous | 248 | 6 (2.4) | 175 | 2 (1.1) | 434 | 10 (2.3) |
| p value | | 0.910 | | 1.000 | | 0.798 |

Table 1.2.12 HIV antibody prevalence by main language spoken at home by parents, gender and survey year

| National Main language spoken at home by parents | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| English speaking | 1653 | 36 (2.2) | 846 | 4 (0.5) | 2508 | 41 (1.6) |
| Non-English speaking | 120 | 1 (0.8) | 39 | 2 (5.1) | 164 | 3 (1.8) |
| p value | | 0.511 | | 0.025 | | 0.750 |
| 2019 | | | | | | |
| English speaking | 1566 | 48 (3.1) | 817 | 5 (0.6) | 2405 | 55 (2.3) |
| Non-English speaking | 100 | 3 (3.0) | 22 | 1 (4.6) | 125 | 4 (3.2) |
| p value | | 1.000 | | 0.148 | | 0.534 |
| 2020 | | | | | | |
| English speaking | 824 | 26 (3.2) | 392 | 3 (0.8) | 1225 | 31 (2.5) |
| Non-English speaking | 48 | 1 (2.1) | 10 | 0 (0.0) | 58 | 1 (1.7) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2021 | | | | | | |
| English speaking | 856 | 15 (1.8) | 495 | 3 (0.6) | 1368 | 20 (1.5) |
| Non-English speaking | 35 | 1 (2.9) | 14 | 0 (0.0) | 51 | 1 (2.0) |
| p value | | 0.476 | | 1.000 | | 0.539 |
| 2022 | | | | | | |
| English speaking | 1052 | 26 (2.5) | 556 | 6 (1.1) | 1643 | 35 (2.1) |
| Non-English speaking | 68 | 2 (2.9) | 22 | 0 (0.0) | 91 | 2 (2.2) |
| p value | | 0.685 | | 1.000 | | 1.000 |

Table 1.2.13 HIV antibody prevalence by region/country of birth, gender and survey year

| National Region/country of birth | Male | | Female | | Total | |
|-------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Australia | 1518 | 33 (2.2) | 793 | 6 (0.8) | 2320 | 39 (1.7) |
| Other Oceania | 57 | 1 (1.8) | 23 | 0 (0.0) | 82 | 2 (2.4) |
| Asia | 26 | 0 (0.0) | 4 | 0 (0.0) | 30 | 0 (0.0) |
| UK & Ireland | 83 | 2 (2.4) | 38 | 0 (0.0) | 121 | 2 (1.7) |
| Other | 90 | 1 (1.1) | 24 | 0 (0.0) | 117 | 1 (0.9) |
| p value | | 0.964 | | 1.000 | | 0.936 |
| 2019 | | | | | | |
| Australia | 1467 | 45 (3.1) | 763 | 6 (0.8) | 2248 | 53 (2.4) |
| Other Oceania | 38 | 1 (2.6) | 23 | 0 (0.0) | 66 | 1 (1.5) |
| Asia | 22 | 3 (13.6) | 3 | 0 (0.0) | 25 | 3 (12.0) |
| UK & Ireland | 55 | 0 (0.0) | 28 | 0 (0.0) | 83 | 0 (0.0) |
| Other | 77 | 2 (2.6) | 22 | 0 (0.0) | 100 | 2 (2.0) |
| p value | | 0.085 | | 1.000 | | 0.055 |
| 2020 | | | | | | |
| Australia | 740 | 22 (3.0) | 365 | 3 (0.8) | 1113 | 26 (2.3) |
| Other Oceania | 33 | 1 (3.0) | 9 | 0 (0.0) | 42 | 1 (2.4) |
| Asia | 16 | 1 (6.3) | 2 | 0 (0.0) | 18 | 1 (5.6) |
| UK & Ireland | 34 | 1 (2.9) | 17 | 0 (0.0) | 51 | 1 (2.0) |
| Other | 39 | 1 (2.6) | 10 | 0 (0.0) | 49 | 1 (2.0) |
| p value | | 0.722 | | 1.000 | | 0.652 |
| 2021 | | | | | | |
| Australia | 776 | 12 (1.6) | 449 | 3 (0.7) | 1241 | 16 (1.3) |
| Other Oceania | 22 | 0 (0.0) | 18 | 0 (0.0) | 43 | 1 (2.3) |
| Asia | 7 | 0 (0.0) | 5 | 0 (0.0) | 12 | 0 (0.0) |
| UK & Ireland | 39 | 0 (0.0) | 15 | 0 (0.0) | 54 | 0 (0.0) |
| Other | 32 | 3 (9.4) | 15 | 0 (0.0) | 47 | 3 (6.4) |
| p value | | 0.080 | | 1.000 | | 0.087 |
| 2022 | | | | | | |
| Australia | 968 | 22 (2.3) | 517 | 6 (1.2) | 1518 | 30 (2.0) |
| Other Oceania | 36 | 2 (5.6) | 13 | 0 (0.0) | 50 | 2 (4.0) |
| Asia | 12 | 0 (0.0) | 11 | 0 (0.0) | 23 | 0 (0.0) |
| UK & Ireland | 35 | 1 (2.9) | 22 | 0 (0.0) | 58 | 1 (1.7) |
| Other | 54 | 3 (5.6) | 14 | 0 (0.0) | 69 | 3 (4.4) |
| p value | | 0.241 | | 1.000 | | 0.352 |

HCV antibody prevalence

Table 1.3.1 HCV antibody prevalence by gender and survey year

| National Survey year | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | 1770 | 817 (46) | 884 | 376 (43) | 2668 | 1200 (45) |
| 2019 | 1670 | 781 (47) | 835 | 360 (43) | 2531 | 1151 (45) |
| 2020 | 877 | 342 (39) | 405 | 163 (40) | 1291 | 507 (39) |
| 2021 | 902 | 343 (38) | 514 | 173 (34) | 1435 | 522 (36) |
| 2022 | 1115 | 350 (31) | 573 | 194 (34) | 1724 | 554 (32) |
| X ² p trend | | <0.001 | | <0.001 | | <0.001 |

Table 1.3.2 HCV antibody prevalence by sexual identity, gender and survey year

| National Sexual identity | Male | | Female | | Total | |
|-----------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heterosexual | 1462 | 690 (47) | 610 | 264 (43) | 2074 | 955 (46) |
| Bisexual | 89 | 34 (38) | 161 | 64 (40) | 253 | 101 (40) |
| Homosexual | 80 | 28 (35) | 35 | 15 (43) | 120 | 44 (37) |
| p value | | 0.032 | | 0.723 | | 0.031 |
| 2019 | | | | | | |
| Heterosexual | 1351 | 645 (48) | 554 | 239 (43) | 1909 | 887 (46) |
| Bisexual | 91 | 38 (42) | 149 | 61 (41) | 253 | 104 (41) |
| Homosexual | 79 | 15 (19) | 36 | 15 (42) | 117 | 30 (26) |
| p value | | <0.001 | | 0.894 | | <0.001 |
| 2020 | | | | | | |
| Heterosexual | 746 | 297 (40) | 275 | 106 (39) | 1022 | 404 (40) |
| Bisexual | 43 | 16 (37) | 72 | 40 (56) | 120 | 56 (47) |
| Homosexual | 38 | 6 (16) | 10 | 3 (30) | 49 | 9 (18) |
| p value | | 0.024 | | 0.027 | | 0.004 |
| 2021 | | | | | | |
| Heterosexual | 791 | 316 (40) | 375 | 118 (31) | 1170 | 436 (37) |
| Bisexual | 49 | 16 (33) | 96 | 38 (40) | 148 | 54 (36) |
| Homosexual | 42 | 6 (14) | 16 | 7 (44) | 63 | 13 (21) |
| p value | | 0.004 | | 0.216 | | 0.156 |
| 2022 | | | | | | |
| Heterosexual | 956 | 316 (33) | 398 | 141 (35) | 1361 | 463 (34) |
| Bisexual | 60 | 17 (28) | 106 | 26 (25) | 175 | 43 (25) |
| Homosexual | 59 | 6 (10) | 24 | 6 (25) | 85 | 13 (15) |
| p value | | 0.005 | | 0.018 | | 0.002 |

Table 1.3.3 HCV antibody prevalence by age group, gender and survey year

| National Age group | Male | | Female | | Total | |
|-----------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <25 years | 58 | 10 (17) | 49 | 10 (20) | 109 | 21 (19) |
| 25-34 years | 283 | 96 (34) | 192 | 75 (39) | 477 | 172 (36) |
| 35-44 years | 638 | 296 (46) | 337 | 150 (45) | 981 | 449 (46) |
| 45+ years | 785 | 411 (52) | 303 | 140 (46) | 1092 | 553 (51) |
| p value | | <0.001 | | 0.005 | | <0.001 |
| 2019 | | | | | | |
| <25 years | 51 | 13 (25) | 40 | 8 (20) | 92 | 21 (23) |
| 25-34 years | 254 | 66 (26) | 175 | 74 (42) | 436 | 141 (32) |
| 35-44 years | 601 | 275 (46) | 331 | 137 (41) | 938 | 416 (44) |
| 45+ years | 756 | 424 (56) | 288 | 141 (49) | 1055 | 569 (54) |
| p value | | <0.001 | | 0.004 | | <0.001 |
| 2020 | | | | | | |
| <25 years | 23 | 3 (13) | 18 | 7 (39) | 42 | 10 (24) |
| 25-34 years | 145 | 30 (21) | 78 | 26 (33) | 225 | 56 (25) |
| 35-44 years | 289 | 118 (41) | 159 | 70 (44) | 449 | 188 (42) |
| 45+ years | 419 | 190 (45) | 150 | 60 (40) | 574 | 252 (44) |
| p value | | <0.001 | | 0.473 | | <0.001 |
| 2021 | | | | | | |
| <25 years | 21 | 4 (19) | 12 | 3 (25) | 34 | 7 (21) |
| 25-34 years | 122 | 30 (25) | 83 | 25 (30) | 209 | 56 (27) |
| 35-44 years | 307 | 126 (41) | 194 | 60 (31) | 510 | 189 (37) |
| 45+ years | 448 | 183 (41) | 221 | 83 (38) | 674 | 268 (40) |
| p value | | 0.001 | | 0.410 | | 0.002 |
| 2022 | | | | | | |
| <25 years | 36 | 5 (14) | 11 | 2 (18) | 50 | 7 (14) |
| 25-34 years | 162 | 27 (17) | 83 | 19 (23) | 250 | 46 (18) |
| 35-44 years | 337 | 102 (30) | 210 | 67 (32) | 554 | 172 (31) |
| 45+ years | 569 | 212 (37) | 263 | 104 (40) | 852 | 322 (38) |
| p value | | <0.001 | | 0.020 | | <0.001 |

Table 1.3.4 HCV antibody prevalence by years since first injection, gender and survey year

| National Years of injection | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <3 years | 83 | 9 (11) | 46 | 2 (4) | 129 | 11 (9) |
| 3 to 10 years | 233 | 51 (22) | 146 | 47 (32) | 380 | 99 (26) |
| 11+ years | 1373 | 713 (52) | 659 | 312 (47) | 2043 | 1031 (50) |
| p value | | <0.001 | | <0.001 | | <0.001 |
| 2019 | | | | | | |
| <3 years | 96 | 8 (8) | 50 | 0 (0) | 146 | 8 (5) |
| 3 to 10 years | 210 | 46 (22) | 145 | 48 (33) | 361 | 96 (27) |
| 11+ years | 1273 | 683 (54) | 601 | 294 (49) | 1889 | 982 (52) |
| p value | | <0.001 | | <0.001 | | <0.001 |
| 2020 | | | | | | |
| <3 years | 56 | 5 (9) | 26 | 0 (0) | 82 | 5 (6) |
| 3 to 10 years | 135 | 21 (16) | 61 | 22 (36) | 198 | 43 (22) |
| 11+ years | 663 | 305 (46) | 308 | 136 (44) | 978 | 443 (45) |
| p value | | <0.001 | | <0.001 | | <0.001 |
| 2021 | | | | | | |
| <3 years | 43 | 2 (5) | 19 | 3 (16) | 64 | 5 (8) |
| 3 to 10 years | 108 | 19 (18) | 76 | 16 (21) | 185 | 35 (19) |
| 11+ years | 719 | 308 (43) | 397 | 147 (37) | 1131 | 460 (41) |
| p value | | <0.001 | | 0.005 | | <0.001 |
| 2022 | | | | | | |
| <3 years | 69 | 3 (4) | 22 | 1 (5) | 94 | 4 (4) |
| 3 to 10 years | 127 | 16 (13) | 76 | 12 (16) | 206 | 29 (14) |
| 11+ years | 850 | 310 (36) | 448 | 170 (38) | 1325 | 488 (37) |
| p value | | <0.001 | | <0.001 | | <0.001 |

Table 1.3.5 HCV antibody prevalence by re-use of someone else's used needle and syringe last month, years since first injection and survey year

| National Receptively shared syringe last month | <3 years | | 3+ years | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| No receptive sharing | 86 | 6 (7) | 1787 | 813 (46) | 1935 | 849 (44) |
| Receptive sharing | 23 | 4 (17) | 391 | 219 (56) | 442 | 244 (55) |
| p value | | 0.214 | | <0.001 | | <0.001 |
| 2019 | | | | | | |
| No receptive sharing | 113 | 5 (4) | 1683 | 803 (48) | 1870 | 853 (46) |
| Receptive sharing | 15 | 1 (7) | 318 | 170 (53) | 362 | 185 (51) |
| p value | | 0.534 | | 0.060 | | 0.055 |
| 2020 | | | | | | |
| No receptive sharing | 65 | 4 (6) | 872 | 352 (40) | 953 | 364 (38) |
| Receptive sharing | 5 | 0 (0) | 171 | 88 (51) | 182 | 93 (51) |
| p value | | 1.000 | | 0.007 | | 0.001 |
| 2021 | | | | | | |
| No receptive sharing | 44 | 3 (7) | 924 | 343 (37) | 1003 | 358 (36) |
| Receptive sharing | 5 | 2 (40) | 202 | 100 (50) | 214 | 106 (50) |
| p value | | 0.075 | | 0.001 | | <0.001 |
| 2022 | | | | | | |
| No receptive sharing | 73 | 2 (3) | 1100 | 347 (32) | 1228 | 370 (30) |
| Receptive sharing | 6 | 2 (33) | 273 | 121 (44) | 299 | 130 (43) |
| p value | | 0.027 | | <0.001 | | <0.001 |

Table 1.3.6 HCV antibody prevalence by last drug injected, years since first injection and survey year

| National Last drug injected | <3 years | | 3+ years | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heroin | 12 | 2 (17) | 650 | 381 (59) | 685 | 399 (58) |
| Methamphetamine | 82 | 7 (9) | 1127 | 433 (38) | 1260 | 463 (37) |
| Other opioids | 3 | 1 (33) | 385 | 212 (55) | 404 | 223 (55) |
| PIEDs | 20 | 0 (0) | 68 | 1 (1) | 95 | 1 (1) |
| Other drugs | 11 | 1 (9) | 149 | 81 (55) | 172 | 88 (51) |
| p value | | 0.266 | | <0.001 | | <0.001 |
| 2019 | | | | | | |
| Heroin | 14 | 2 (14) | 632 | 415 (66) | 677 | 441 (65) |
| Methamphetamine | 90 | 4 (4) | 1058 | 385 (36) | 1210 | 415 (34) |
| Other opioids | 7 | 1 (14) | 319 | 171 (54) | 334 | 178 (53) |
| PIEDs | 27 | 0 (0) | 59 | 1 (2) | 91 | 1 (1) |
| Other drugs | 5 | 1 (20) | 148 | 89 (60) | 168 | 97 (58) |
| p value | | 0.147 | | <0.001 | | <0.001 |
| 2020 | | | | | | |
| Heroin | 6 | 1 (17) | 267 | 160 (60) | 278 | 164 (59) |
| Methamphetamine | 42 | 4 (10) | 546 | 162 (30) | 599 | 173 (29) |
| Other opioids | 5 | 0 (0) | 230 | 116 (50) | 238 | 119 (50) |
| PIEDs | 23 | 0 (0) | 49 | 2 (4) | 79 | 2 (3) |
| Other drugs | 4 | 0 (0) | 69 | 40 (58) | 76 | 42 (55) |
| p value | | 0.404 | | <0.001 | | <0.001 |
| 2021 | | | | | | |
| Heroin | 4 | 1 (25) | 277 | 134 (48) | 289 | 139 (48) |
| Methamphetamine | 39 | 1 (3) | 649 | 184 (28) | 710 | 193 (27) |
| Other opioids | 8 | 2 (25) | 239 | 119 (50) | 257 | 125 (49) |
| PIEDs | 8 | 0 (0) | 41 | 4 (10) | 51 | 4 (8) |
| Other drugs | 2 | 0 (0) | 81 | 42 (52) | 91 | 45 (49) |
| p value | | 0.079 | | <0.001 | | <0.001 |
| 2022 | | | | | | |
| Heroin | 4 | 0 (0) | 406 | 185 (46) | 424 | 191 (45) |
| Methamphetamine | 35 | 3 (9) | 725 | 195 (27) | 806 | 216 (27) |
| Other opioids | 7 | 1 (14) | 196 | 92 (47) | 214 | 97 (45) |
| PIEDs | 43 | 0 (0) | 63 | 2 (3) | 115 | 3 (3) |
| Other drugs | 2 | 0 (0) | 107 | 35 (33) | 117 | 36 (31) |
| p value | | 0.255 | | <0.001 | | <0.001 |

Table 1.3.7 HCV antibody prevalence by frequency of drug injection last month, years since first injection and survey year

| National Frequency of injection last month | <3 years | | 3+ years | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| Less than daily | 60 | 6 (10) | 948 | 432 (46) | 1040 | 453 (44) |
| Daily or more | 50 | 4 (8) | 1253 | 611 (49) | 1362 | 652 (48) |
| Not last month | 18 | 1 (6) | 185 | 68 (37) | 223 | 74 (33) |
| p value | | 0.914 | | 0.007 | | <0.001 |
| 2019 | | | | | | |
| Less than daily | 65 | 4 (6) | 835 | 365 (44) | 935 | 389 (41) |
| Daily or more | 66 | 3 (5) | 1190 | 617 (52) | 1326 | 660 (50) |
| Not last month | 15 | 1 (7) | 195 | 82 (42) | 231 | 86 (37) |
| p value | | 0.772 | | <0.001 | | <0.001 |
| 2020 | | | | | | |
| Less than daily | 38 | 2 (5) | 469 | 170 (36) | 516 | 176 (34) |
| Daily or more | 32 | 2 (6) | 581 | 273 (47) | 626 | 284 (45) |
| Not last month | 12 | 1 (8) | 116 | 38 (33) | 137 | 41 (30) |
| p value | | 1.000 | | <0.001 | | <0.001 |
| 2021 | | | | | | |
| Less than daily | 32 | 1 (3) | 541 | 160 (30) | 589 | 167 (28) |
| Daily or more | 18 | 4 (22) | 592 | 286 (48) | 637 | 301 (47) |
| Not last month | 13 | 0 (0) | 160 | 38 (24) | 181 | 41 (23) |
| p value | | 0.041 | | <0.001 | | <0.001 |
| 2022 | | | | | | |
| Less than daily | 55 | 1 (2) | 575 | 177 (31) | 657 | 187 (28) |
| Daily or more | 24 | 3 (13) | 802 | 293 (37) | 877 | 315 (36) |
| Not last month | 13 | 0 (0) | 129 | 42 (33) | 155 | 44 (28) |
| p value | | 0.106 | | 0.079 | | 0.005 |

Table 1.3.8 HCV antibody prevalence by imprisonment last year, years since first injection and survey year

| National Imprisonment last year | <3 years | | 3+ years | | Total | |
|------------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| No imprisonment | 109 | 8 (7) | 1979 | 889 (45) | 2166 | 932 (43) |
| Imprisonment | 16 | 1 (6) | 300 | 172 (57) | 338 | 186 (55) |
| p value | | 1.000 | | <0.001 | | <0.001 |
| 2019 | | | | | | |
| No imprisonment | 129 | 7 (5) | 1815 | 827 (46) | 2041 | 878 (43) |
| Imprisonment | 12 | 0 (0) | 301 | 178 (59) | 342 | 194 (57) |
| p value | | 1.000 | | <0.001 | | <0.001 |
| 2020 | | | | | | |
| No imprisonment | 75 | 4 (5) | 969 | 381 (39) | 1068 | 395 (37) |
| Imprisonment | 7 | 1 (14) | 141 | 68 (48) | 156 | 75 (48) |
| p value | | 0.367 | | 0.044 | | 0.008 |
| 2021 | | | | | | |
| No imprisonment | 55 | 4 (7) | 1100 | 395 (36) | 1196 | 413 (35) |
| Imprisonment | 4 | 0 (0) | 142 | 66 (46) | 154 | 70 (45) |
| p value | | 1.000 | | 0.014 | | 0.008 |
| 2022 | | | | | | |
| No imprisonment | 85 | 2 (2) | 1298 | 424 (33) | 1459 | 450 (31) |
| Imprisonment | 7 | 2 (29) | 171 | 65 (38) | 195 | 74 (38) |
| p value | | 0.028 | | 0.163 | | 0.045 |

Table 1.3.9 HCV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| National Aboriginal and Torres Strait Islander origin | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Non Indigenous | 1431 | 625 (44) | 660 | 270 (41) | 2101 | 899 (43) |
| Indigenous | 305 | 175 (57) | 208 | 99 (48) | 516 | 276 (53) |
| p value | | <0.001 | | 0.089 | | <0.001 |
| 2019 | | | | | | |
| Non Indigenous | 1309 | 584 (45) | 610 | 253 (41) | 1937 | 844 (44) |
| Indigenous | 331 | 180 (54) | 214 | 102 (48) | 551 | 284 (52) |
| p value | | 0.001 | | 0.116 | | 0.001 |
| 2020 | | | | | | |
| Non Indigenous | 676 | 234 (35) | 277 | 101 (36) | 958 | 337 (35) |
| Indigenous | 167 | 91 (54) | 115 | 54 (47) | 284 | 145 (51) |
| p value | | <0.001 | | 0.053 | | <0.001 |
| 2021 | | | | | | |
| Non Indigenous | 650 | 214 (33) | 347 | 108 (31) | 1008 | 324 (32) |
| Indigenous | 207 | 108 (52) | 145 | 58 (40) | 359 | 169 (47) |
| p value | | <0.001 | | 0.058 | | <0.001 |
| 2022 | | | | | | |
| Non Indigenous | 849 | 251 (30) | 390 | 130 (33) | 1262 | 387 (31) |
| Indigenous | 246 | 92 (37) | 174 | 59 (34) | 431 | 154 (36) |
| p value | | 0.020 | | 0.894 | | 0.052 |

Table 1.3.10 HCV antibody prevalence by main language spoken at home by parents, gender and survey year

| National Main language spoken at home by parents | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| English speaking | 1640 | 752 (46) | 843 | 360 (43) | 2492 | 1116 (45) |
| Non-English speaking | 118 | 61 (52) | 39 | 16 (41) | 162 | 80 (49) |
| p value | | 0.219 | | 0.836 | | 0.254 |
| 2019 | | | | | | |
| English speaking | 1561 | 726 (47) | 811 | 347 (43) | 2394 | 1081 (45) |
| Non-English speaking | 100 | 53 (53) | 22 | 12 (55) | 125 | 67 (54) |
| p value | | 0.207 | | 0.272 | | 0.065 |
| 2020 | | | | | | |
| English speaking | 824 | 319 (39) | 391 | 157 (40) | 1224 | 478 (39) |
| Non-English speaking | 48 | 22 (46) | 10 | 6 (60) | 58 | 28 (48) |
| p value | | 0.326 | | 0.328 | | 0.160 |
| 2021 | | | | | | |
| English speaking | 856 | 326 (38) | 495 | 163 (33) | 1368 | 494 (36) |
| Non-English speaking | 35 | 14 (40) | 14 | 7 (50) | 51 | 22 (43) |
| p value | | 0.819 | | 0.182 | | 0.306 |
| 2022 | | | | | | |
| English speaking | 1038 | 318 (31) | 547 | 188 (34) | 1619 | 516 (32) |
| Non-English speaking | 66 | 28 (42) | 21 | 5 (24) | 88 | 33 (38) |
| p value | | 0.045 | | 0.316 | | 0.271 |

Table 1.3.11 HCV antibody prevalence by region/country of birth, gender and survey year

| National Region/Country of birth | Male | | Female | | Total | |
|-------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Australia | 1504 | 689 (46) | 790 | 342 (43) | 2303 | 1035 (45) |
| Other Oceania | 57 | 23 (40) | 23 | 6 (26) | 82 | 29 (35) |
| Asia | 26 | 17 (65) | 4 | 4 (100) | 30 | 21 (70) |
| UK & Ireland | 82 | 39 (48) | 38 | 14 (37) | 120 | 53 (44) |
| Other | 90 | 44 (49) | 24 | 8 (33) | 117 | 55 (47) |
| p value | | 0.282 | | 0.052 | | 0.028 |
| 2019 | | | | | | |
| Australia | 1463 | 684 (47) | 757 | 325 (43) | 2238 | 1015 (45) |
| Other Oceania | 38 | 19 (50) | 23 | 11 (48) | 66 | 32 (48) |
| Asia | 23 | 7 (30) | 3 | 1 (33) | 26 | 8 (31) |
| UK & Ireland | 55 | 31 (56) | 28 | 14 (50) | 83 | 45 (54) |
| Other | 75 | 33 (44) | 22 | 7 (32) | 98 | 40 (41) |
| p value | | 0.299 | | 0.741 | | 0.204 |
| 2020 | | | | | | |
| Australia | 740 | 292 (39) | 365 | 149 (41) | 1113 | 443 (40) |
| Other Oceania | 33 | 10 (30) | 9 | 2 (22) | 42 | 12 (29) |
| Asia | 16 | 5 (31) | 2 | 1 (50) | 18 | 6 (33) |
| UK & Ireland | 34 | 10 (29) | 16 | 5 (31) | 50 | 15 (30) |
| Other | 39 | 22 (56) | 10 | 5 (50) | 49 | 27 (55) |
| p value | | 0.102 | | 0.679 | | 0.054 |
| 2021 | | | | | | |
| Australia | 775 | 292 (38) | 449 | 147 (33) | 1240 | 445 (36) |
| Other Oceania | 22 | 12 (55) | 18 | 8 (44) | 43 | 20 (47) |
| Asia | 7 | 1 (14) | 5 | 3 (60) | 12 | 4 (33) |
| UK & Ireland | 39 | 12 (31) | 15 | 3 (20) | 54 | 15 (28) |
| Other | 33 | 14 (42) | 15 | 7 (47) | 48 | 21 (44) |
| p value | | 0.270 | | 0.264 | | 0.293 |
| 2022 | | | | | | |
| Australia | 951 | 293 (31) | 508 | 173 (34) | 1491 | 475 (32) |
| Other Oceania | 36 | 16 (44) | 13 | 4 (31) | 50 | 21 (42) |
| Asia | 12 | 2 (17) | 11 | 3 (27) | 23 | 5 (22) |
| UK & Ireland | 35 | 7 (20) | 22 | 7 (32) | 58 | 14 (24) |
| Other | 54 | 19 (35) | 13 | 2 (15) | 68 | 21 (31) |
| p value | | 0.169 | | 0.739 | | 0.280 |

HCV RNA prevalence

Table 1.4.1 HCV RNA prevalence by gender and survey year

| National Survey year | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | 932 | 195 (21) | 458 | 89 (19) | 1397 | 284 (20) |
| 2019* | 1412 | 264 (19) | 706 | 111 (16) | 2140 | 378 (18) |
| 2020 | 808 | 136 (17) | 361 | 57 (16) | 1175 | 193 (16) |
| 2021 | 835 | 148 (18) | 474 | 62 (13) | 1328 | 214 (16) |
| 2022 | 1065 | 137 (13) | 552 | 58 (11) | 1653 | 200 (12) |
| X ² p trend | | <0.001 | | <0.001 | | <0.001 |

* 2018, 2019: Weighted for gender and HCV antibody status

Totals include respondents where gender was reported as other or not reported

Table 1.4.2 HCV RNA prevalence by sexual identity, gender and survey year

| National Sexual identity | Male | | Female | | Total | |
|-----------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heterosexual | 764 | 166 (22) | 309 | 65 (21) | 1074 | 231 (22) |
| Bisexual | 44 | 7 (16) | 85 | 13 (15) | 131 | 21 (16) |
| Homosexual | 44 | 5 (11) | 21 | 3 (14) | 68 | 8 (12) |
| p value | | 0.155 | | 0.439 | | 0.062 |
| 2019* | | | | | | |
| Heterosexual | 1148 | 215 (19) | 460 | 69 (15) | 1611 | 285 (18) |
| Bisexual | 77 | 14 (18) | 132 | 20 (15) | 220 | 36 (16) |
| Homosexual | 70 | 3 (4) | 34 | 6 (18) | 106 | 9 (8) |
| p value | | 0.008 | | 0.921 | | 0.043 |
| 2020 | | | | | | |
| Heterosexual | 684 | 117 (17) | 244 | 43 (18) | 929 | 160 (17) |
| Bisexual | 41 | 7 (17) | 65 | 10 (15) | 109 | 17 (16) |
| Homosexual | 36 | 2 (6) | 10 | 0 (0) | 46 | 2 (4) |
| p value | | 0.191 | | 0.326 | | 0.069 |
| 2021 | | | | | | |
| Heterosexual | 736 | 136 (18) | 345 | 41 (12) | 1085 | 179 (16) |
| Bisexual | 43 | 8 (19) | 90 | 11 (12) | 136 | 19 (14) |
| Homosexual | 37 | 1 (3) | 16 | 4 (25) | 58 | 5 (9) |
| p value | | 0.049 | | 0.298 | | 0.226 |
| 2022 | | | | | | |
| Heterosexual | 914 | 122 (13) | 386 | 47 (12) | 1307 | 171 (13) |
| Bisexual | 58 | 7 (12) | 100 | 5 (5) | 166 | 12 (7) |
| Homosexual | 54 | 3 (6) | 21 | 1 (5) | 77 | 5 (6) |
| p value | | 0.314 | | 0.311 | | 0.102 |

Table 1.4.3 HCV RNA prevalence by age group, gender and survey year

| National Age group | Male | | Female | | Total | |
|-----------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <25 years | 33 | 6 (18) | 25 | 6 (24) | 59 | 12 (20) |
| 25-34 years | 138 | 22 (16) | 102 | 23 (23) | 241 | 45 (19) |
| 35-44 years | 345 | 81 (23) | 172 | 34 (20) | 521 | 116 (22) |
| 45+ years | 410 | 84 (20) | 157 | 26 (17) | 570 | 110 (19) |
| p value | | 0.571 | | 0.633 | | 0.686 |
| 2019* | | | | | | |
| <25 years | 44 | 5 (11) | 39 | 7 (18) | 84 | 12 (14) |
| 25-34 years | 220 | 27 (12) | 146 | 24 (16) | 372 | 52 (14) |
| 35-44 years | 503 | 102 (20) | 277 | 46 (17) | 785 | 148 (19) |
| 45+ years | 639 | 129 (20) | 244 | 34 (14) | 891 | 165 (19) |
| p value | | 0.028 | | 0.773 | | 0.143 |
| 2020 | | | | | | |
| <25 years | 20 | 1 (5) | 19 | 4 (21) | 40 | 5 (13) |
| 25-34 years | 129 | 12 (9) | 71 | 12 (17) | 201 | 24 (12) |
| 35-44 years | 271 | 63 (23) | 139 | 27 (19) | 411 | 90 (22) |
| 45+ years | 387 | 60 (16) | 132 | 14 (11) | 522 | 74 (14) |
| p value | | 0.001 | | 0.211 | | 0.003 |
| 2021 | | | | | | |
| <25 years | 21 | 2 (10) | 12 | 2 (17) | 34 | 4 (12) |
| 25-34 years | 113 | 21 (19) | 78 | 10 (13) | 195 | 33 (17) |
| 35-44 years | 281 | 55 (20) | 174 | 18 (10) | 464 | 74 (16) |
| 45+ years | 416 | 70 (17) | 206 | 31 (15) | 627 | 102 (16) |
| p value | | 0.593 | | 0.573 | | 0.899 |
| 2022 | | | | | | |
| <25 years | 35 | 3 (9) | 12 | 1 (8) | 51 | 4 (8) |
| 25-34 years | 158 | 14 (9) | 79 | 8 (10) | 242 | 22 (9) |
| 35-44 years | 326 | 39 (12) | 201 | 18 (9) | 534 | 59 (11) |
| 45+ years | 535 | 80 (15) | 254 | 31 (12) | 808 | 114 (14) |
| p value | | 0.181 | | 0.731 | | 0.103 |

Table 1.4.4 HCV RNA prevalence by years since first injection, gender and survey year

| National Years of injection | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <3 years | 37 | 3 (8) | 23 | 3 (13) | 60 | 6 (10) |
| 3 to 10 years | 118 | 20 (17) | 66 | 15 (23) | 184 | 35 (19) |
| 11+ years | 733 | 159 (22) | 351 | 67 (19) | 1090 | 227 (21) |
| p value | | 0.163 | | 0.537 | | 0.164 |
| 2019* | | | | | | |
| <3 years | 87 | 7 (8) | 43 | 1 (2) | 130 | 8 (6) |
| 3 to 10 years | 174 | 22 (13) | 126 | 28 (22) | 305 | 51 (17) |
| 11+ years | 1073 | 220 (21) | 509 | 76 (15) | 1594 | 297 (19) |
| p value | | 0.001 | | 0.007 | | 0.001 |
| 2020 | | | | | | |
| <3 years | 51 | 2 (4) | 22 | 0 (0) | 73 | 2 (3) |
| 3 to 10 years | 125 | 12 (10) | 57 | 12 (21) | 183 | 24 (13) |
| 11+ years | 610 | 117 (19) | 273 | 44 (16) | 888 | 161 (18) |
| p value | | 0.001 | | 0.071 | | 0.001 |
| 2021 | | | | | | |
| <3 years | 42 | 1 (2) | 17 | 1 (6) | 61 | 2 (3) |
| 3 to 10 years | 103 | 21 (20) | 67 | 9 (13) | 171 | 31 (18) |
| 11+ years | 660 | 122 (18) | 368 | 48 (13) | 1043 | 172 (16) |
| p value | | 0.024 | | 0.680 | | 0.018 |
| 2022 | | | | | | |
| <3 years | 67 | 2 (3) | 23 | 0 (0) | 94 | 2 (2) |
| 3 to 10 years | 126 | 12 (10) | 75 | 8 (11) | 204 | 20 (10) |
| 11+ years | 807 | 115 (14) | 427 | 48 (11) | 1260 | 168 (13) |
| p value | | 0.009 | | 0.271 | | 0.001 |

Table 1.4.5 HCV RNA prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| National Receptively shared syringe last month | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| No receptive sharing | 685 | 133 (19) | 325 | 60 (18) | 1012 | 193 (19) |
| Receptive sharing | 143 | 48 (34) | 69 | 18 (26) | 214 | 66 (31) |
| p value | | <0.001 | | 0.164 | | <0.001 |
| 2019* | | | | | | |
| No receptive sharing | 1055 | 205 (19) | 527 | 81 (15) | 1597 | 288 (18) |
| Receptive sharing | 196 | 43 (22) | 99 | 21 (21) | 300 | 65 (22) |
| p value | | 0.456 | | 0.169 | | 0.152 |
| 2020 | | | | | | |
| No receptive sharing | 610 | 97 (16) | 246 | 39 (16) | 861 | 136 (16) |
| Receptive sharing | 94 | 33 (35) | 72 | 14 (19) | 166 | 47 (28) |
| p value | | <0.001 | | 0.472 | | <0.001 |
| 2021 | | | | | | |
| No receptive sharing | 588 | 101 (17) | 325 | 39 (12) | 923 | 142 (15) |
| Receptive sharing | 124 | 36 (29) | 72 | 15 (21) | 201 | 52 (26) |
| p value | | 0.002 | | 0.048 | | <0.001 |
| 2022 | | | | | | |
| No receptive sharing | 769 | 95 (12) | 385 | 30 (8) | 1180 | 127 (11) |
| Receptive sharing | 175 | 34 (19) | 101 | 22 (22) | 283 | 59 (21) |
| p value | | 0.014 | | <0.001 | | <0.001 |

Table 1.4.6 HCV RNA prevalence by last drug injected, gender and survey year

| National Last drug injected | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heroin | 294 | 59 (20) | 139 | 26 (19) | 435 | 85 (20) |
| Methamphetamine | 363 | 87 (24) | 205 | 35 (17) | 572 | 122 (21) |
| Other opioids | 142 | 28 (20) | 71 | 17 (24) | 214 | 45 (21) |
| PIEDs | 50 | 1 (2) | 1 | 0 (0) | 51 | 1 (2) |
| Other drugs | 64 | 16 (25) | 33 | 9 (27) | 98 | 25 (26) |
| p value | | 0.038 | | 0.531 | | 0.026 |
| 2019* | | | | | | |
| Heroin | 364 | 81 (22) | 205 | 41 (20) | 578 | 122 (21) |
| Methamphetamine | 659 | 122 (19) | 351 | 46 (13) | 1019 | 171 (17) |
| Other opioids | 202 | 39 (19) | 87 | 9 (10) | 291 | 49 (17) |
| PIEDs | 73 | 1 (1) | 5 | 0 (0) | 78 | 1 (1) |
| Other drugs | 90 | 18 (20) | 48 | 13 (27) | 139 | 30 (22) |
| p value | | 0.001 | | 0.019 | | <0.001 |
| 2020 | | | | | | |
| Heroin | 177 | 34 (19) | 85 | 14 (16) | 263 | 48 (18) |
| Methamphetamine | 354 | 56 (16) | 171 | 24 (14) | 529 | 80 (15) |
| Other opioids | 145 | 31 (21) | 74 | 8 (11) | 220 | 39 (18) |
| PIEDs | 71 | 0 (0) | 3 | 0 (0) | 74 | 0 (0) |
| Other drugs | 46 | 12 (26) | 25 | 11 (44) | 71 | 23 (32) |
| p value | | <0.001 | | 0.002 | | <0.001 |
| 2021 | | | | | | |
| Heroin | 164 | 29 (18) | 95 | 14 (15) | 265 | 45 (17) |
| Methamphetamine | 382 | 72 (19) | 267 | 28 (10) | 656 | 100 (15) |
| Other opioids | 173 | 32 (18) | 67 | 11 (16) | 244 | 44 (18) |
| PIEDs | 48 | 3 (6) | 0 | 0 (0) | 48 | 3 (6) |
| Other drugs | 50 | 7 (14) | 32 | 7 (22) | 82 | 14 (17) |
| p value | | 0.264 | | 0.198 | | 0.320 |
| 2022 | | | | | | |
| Heroin | 228 | 28 (12) | 158 | 22 (14) | 391 | 50 (13) |
| Methamphetamine | 470 | 73 (16) | 292 | 25 (9) | 783 | 100 (13) |
| Other opioids | 158 | 21 (13) | 52 | 8 (15) | 216 | 32 (15) |
| PIEDs | 111 | 0 (0) | 3 | 0 (0) | 114 | 0 (0) |
| Other drugs | 77 | 12 (16) | 25 | 2 (8) | 104 | 14 (13) |
| p value | | <0.001 | | 0.304 | | 0.001 |

Table 1.4.7 HCV RNA prevalence by frequency of drug injection last month, gender and survey year

| National Frequency of injection last month | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| Less than daily | 368 | 71 (19) | 173 | 35 (20) | 542 | 106 (20) |
| Daily or more | 468 | 110 (24) | 227 | 44 (19) | 696 | 154 (22) |
| Not last month | 81 | 12 (15) | 52 | 8 (15) | 134 | 20 (15) |
| p value | | 0.144 | | 0.733 | | 0.141 |
| 2019* | | | | | | |
| Less than daily | 559 | 93 (17) | 246 | 40 (16) | 810 | 133 (16) |
| Daily or more | 706 | 157 (22) | 390 | 63 (16) | 1108 | 223 (20) |
| Not last month | 131 | 12 (9) | 64 | 7 (11) | 197 | 19 (10) |
| p value | | <0.001 | | 0.518 | | <0.001 |
| 2020 | | | | | | |
| Less than daily | 326 | 49 (15) | 141 | 22 (16) | 471 | 71 (15) |
| Daily or more | 383 | 81 (21) | 179 | 31 (17) | 563 | 112 (20) |
| Not last month | 90 | 4 (4) | 38 | 4 (11) | 129 | 8 (6) |
| p value | | <0.001 | | 0.578 | | <0.001 |
| 2021 | | | | | | |
| Less than daily | 352 | 56 (16) | 196 | 20 (10) | 554 | 76 (14) |
| Daily or more | 362 | 81 (22) | 206 | 35 (17) | 577 | 119 (21) |
| Not last month | 106 | 6 (6) | 62 | 5 (8) | 170 | 11 (6) |
| p value | | <0.001 | | 0.060 | | <0.001 |
| 2022 | | | | | | |
| Less than daily | 424 | 47 (11) | 206 | 18 (9) | 647 | 67 (10) |
| Daily or more | 525 | 82 (16) | 281 | 34 (12) | 822 | 119 (14) |
| Not last month | 100 | 5 (5) | 50 | 5 (10) | 152 | 10 (7) |
| p value | | 0.006 | | 0.487 | | 0.005 |

Table 1.4.8 HCV RNA prevalence by imprisonment last year, gender and survey year

| National Imprisonment last year | Male | | Female | | Total | |
|------------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| No imprisonment | 732 | 147 (20) | 388 | 76 (20) | 1125 | 223 (20) |
| Imprisonment | 134 | 32 (24) | 41 | 9 (22) | 176 | 41 (23) |
| p value | | 0.326 | | 0.674 | | 0.287 |
| 2019* | | | | | | |
| No imprisonment | 1107 | 186 (17) | 594 | 88 (15) | 1718 | 276 (16) |
| Imprisonment | 215 | 57 (27) | 76 | 17 (22) | 295 | 75 (25) |
| p value | | <0.001 | | 0.101 | | <0.001 |
| 2020 | | | | | | |
| No imprisonment | 658 | 102 (16) | 309 | 48 (16) | 973 | 150 (15) |
| Imprisonment | 107 | 26 (24) | 35 | 7 (20) | 142 | 33 (23) |
| p value | | 0.024 | | 0.494 | | 0.019 |
| 2021 | | | | | | |
| No imprisonment | 680 | 110 (16) | 412 | 50 (12) | 1108 | 163 (15) |
| Imprisonment | 107 | 23 (21) | 36 | 6 (17) | 144 | 29 (20) |
| p value | | 0.172 | | 0.431 | | 0.089 |
| 2022 | | | | | | |
| No imprisonment | 882 | 105 (12) | 486 | 50 (10) | 1399 | 160 (11) |
| Imprisonment | 141 | 24 (17) | 46 | 7 (15) | 189 | 31 (16) |
| p value | | 0.089 | | 0.302 | | 0.049 |

Table 1.4.9 HCV RNA prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| National Aboriginal and Torres Strait Islander origin | Male | | Female | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| Non Indigenous | 750 | 144 (19) | 339 | 66 (19) | 1094 | 210 (19) |
| Indigenous | 163 | 47 (29) | 107 | 20 (19) | 272 | 67 (25) |
| p value | | 0.020 | | 0.880 | | 0.073 |
| 2019* | | | | | | |
| Non Indigenous | 1117 | 185 (17) | 514 | 71 (14) | 1646 | 258 (16) |
| Indigenous | 273 | 74 (27) | 182 | 39 (21) | 461 | 114 (25) |
| p value | | <0.001 | | 0.020 | | <0.001 |
| 2020 | | | | | | |
| Non Indigenous | 617 | 90 (15) | 241 | 35 (15) | 862 | 125 (15) |
| Indigenous | 158 | 38 (24) | 108 | 20 (19) | 267 | 58 (22) |
| p value | | 0.004 | | 0.344 | | 0.005 |
| 2021 | | | | | | |
| Non Indigenous | 596 | 81 (14) | 315 | 40 (13) | 922 | 123 (13) |
| Indigenous | 197 | 55 (28) | 140 | 21 (15) | 344 | 78 (23) |
| p value | | <0.001 | | 0.506 | | <0.001 |
| 2022 | | | | | | |
| Non Indigenous | 809 | 101 (12) | 375 | 41 (11) | 1207 | 144 (12) |
| Indigenous | 238 | 34 (14) | 168 | 15 (9) | 417 | 51 (12) |
| p value | | 0.466 | | 0.478 | | 0.871 |

Table 1.4.10 HCV RNA prevalence by main language spoken at home by parents, gender and survey year

| National Main language spoken at home by parents | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| English speaking | 859 | 178 (21) | 432 | 85 (20) | 1296 | 263 (20) |
| Non-English speaking | 66 | 17 (26) | 23 | 4 (17) | 90 | 21 (23) |
| p value | | 0.364 | | 0.956 | | 0.484 |
| 2019* | | | | | | |
| English speaking | 1322 | 248 (19) | 685 | 105 (15) | 2026 | 355 (18) |
| Non-English speaking | 85 | 17 (20) | 19 | 6 (32) | 107 | 23 (21) |
| p value | | 0.858 | | 0.056 | | 0.242 |
| 2020 | | | | | | |
| English speaking | 758 | 127 (17) | 351 | 54 (15) | 1115 | 181 (16) |
| Non-English speaking | 46 | 9 (20) | 9 | 3 (33) | 55 | 12 (22) |
| p value | | 0.622 | | 0.145 | | 0.276 |
| 2021 | | | | | | |
| English speaking | 794 | 140 (18) | 459 | 56 (12) | 1270 | 199 (16) |
| Non-English speaking | 32 | 7 (22) | 13 | 4 (31) | 47 | 12 (26) |
| p value | | 0.538 | | 0.047 | | 0.070 |
| 2022 | | | | | | |
| English speaking | 993 | 124 (12) | 527 | 55 (10) | 1555 | 184 (12) |
| Non-English speaking | 61 | 12 (20) | 20 | 2 (10) | 81 | 14 (17) |
| p value | | 0.104 | | 1.000 | | 0.143 |

Table 1.4.11 HCV RNA prevalence by region/country of birth, gender and survey year

| National Region/Country of birth | Male | | Female | | Total | |
|-------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Australia | 806 | 166 (21) | 400 | 80 (20) | 1211 | 247 (20) |
| Other Oceania | 26 | 3 (12) | 12 | 2 (17) | 39 | 5 (13) |
| Asia | 9 | 5 (56) | 1 | 0 (0) | 10 | 5 (50) |
| UK & Ireland | 35 | 5 (14) | 20 | 5 (25) | 55 | 10 (18) |
| Other | 49 | 14 (29) | 18 | 2 (11) | 69 | 16 (23) |
| p value | | 0.014 | | 0.650 | | 0.060 |
| 2019* | | | | | | |
| Australia | 1244 | 238 (19) | 636 | 101 (16) | 1895 | 340 (18) |
| Other Oceania | 32 | 4 (13) | 19 | 4 (21) | 55 | 10 (18) |
| Asia | 16 | 3 (19) | 3 | 0 (0) | 19 | 3 (16) |
| UK & Ireland | 44 | 8 (18) | 26 | 4 (15) | 70 | 13 (19) |
| Other | 63 | 10 (16) | 21 | 1 (5) | 85 | 11 (13) |
| p value | | 0.813 | | 0.575 | | 0.782 |
| 2020 | | | | | | |
| Australia | 681 | 109 (16) | 328 | 53 (16) | 1015 | 162 (16) |
| Other Oceania | 31 | 5 (16) | 8 | 1 (13) | 39 | 6 (15) |
| Asia | 16 | 0 (0) | 2 | 0 (0) | 18 | 0 (0) |
| UK & Ireland | 29 | 7 (24) | 13 | 2 (15) | 42 | 9 (21) |
| Other | 37 | 13 (35) | 9 | 1 (11) | 46 | 14 (30) |
| p value | | 0.018 | | 0.976 | | 0.048 |
| 2021 | | | | | | |
| Australia | 718 | 126 (18) | 416 | 52 (13) | 1150 | 181 (16) |
| Other Oceania | 20 | 7 (35) | 15 | 4 (27) | 38 | 12 (32) |
| Asia | 6 | 1 (17) | 5 | 2 (40) | 11 | 3 (27) |
| UK & Ireland | 36 | 6 (17) | 14 | 1 (7) | 50 | 7 (14) |
| Other | 30 | 5 (17) | 14 | 2 (14) | 44 | 7 (16) |
| p value | | 0.391 | | 0.187 | | 0.093 |
| 2022 | | | | | | |
| Australia | 912 | 111 (12) | 489 | 52 (11) | 1434 | 167 (12) |
| Other Oceania | 34 | 6 (18) | 12 | 1 (8) | 47 | 8 (17) |
| Asia | 12 | 2 (17) | 10 | 1 (10) | 22 | 3 (14) |
| UK & Ireland | 32 | 3 (9) | 22 | 2 (9) | 55 | 5 (9) |
| Other | 51 | 9 (18) | 13 | 1 (8) | 64 | 10 (16) |
| p value | | 0.512 | | 1.000 | | 0.559 |

Australian Capital Territory

Table 2.1.1 Number (percentage) of respondents by demographic characteristics and survey year

| Australian Capital Territory | 2018 | 2019 | 2020* | 2021* | 2022 |
|---|----------|----------|----------|---------|-----------|
| Number of sites | 1 | 1 | 1 | 1 | 1 |
| N° surveyed | N=127 | N=128 | N=104 | N=100 | N=100 |
| Response rate (%) | 93% | 71% | 67% | 68% | 67% |
| Gender (%) | | | | | |
| Male | 90 (71) | 86 (67) | 70 (67) | 68 (68) | 67 (67) |
| Female | 35 (28) | 42 (33) | 33 (32) | 32 (32) | 32 (32) |
| Other | 2 (2) | 0 (0) | 1 (1) | 0 (0) | 1 (1) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Sexual identity (%) | | | | | |
| Heterosexual | 100 (79) | 93 (73) | 84 (81) | 87 (87) | 81 (81) |
| Bisexual | 13 (10) | 13 (10) | 11 (11) | 7 (7) | 11 (11) |
| Gay/Lesbian | 3 (2) | 4 (3) | 2 (2) | 4 (4) | 1 (1) |
| Other | -- -- | -- -- | 3 (3) | 1 (1) | 7 (7) |
| Not reported | 11 (9) | 18 (14) | 4 (4) | 1 (1) | 0 (0) |
| Aboriginal and Torres Strait Islander origin (%) | | | | | |
| No | 98 (77) | 106 (83) | 82 (79) | 68 (68) | 71 (71) |
| Yes | 27 (21) | 21 (16) | 19 (18) | 29 (29) | 27 (27) |
| Not reported | 2 (2) | 1 (1) | 3 (3) | 3 (3) | 2 (2) |
| Region/country of birth (%) | | | | | |
| Australia | 110 (87) | 113 (88) | 93 (89) | 91 (91) | 85 (85) |
| Other Oceania | 1 (1) | 6 (5) | 3 (3) | 2 (2) | 5 (5) |
| Asia | 1 (1) | 0 (0) | 1 (1) | 0 (0) | 2 (2) |
| UK & Ireland | 6 (5) | 4 (3) | 3 (3) | 3 (3) | 4 (4) |
| Other | 7 (6) | 5 (4) | 2 (2) | 4 (4) | 1 (1) |
| Not reported | 2 (2) | 0 (0) | 2 (2) | 0 (0) | 3 (3) |
| Main language spoken at home by parents (%) | | | | | |
| English | 114 (90) | 119 (93) | 99 (95) | 95 (95) | 94 (94) |
| Non-English | 11 (9) | 9 (7) | 5 (5) | 5 (5) | 4 (4) |
| Not reported | 2 (2) | 0 (0) | 0 (0) | 0 (0) | 2 (2) |
| Age and time since first injection (years) | | | | | |
| <i>Median age</i> | 41 | 42 | 44 | 42.5 | 45 |
| <i>Age range</i> | 20-67 | 18-66 | 23-63 | 21-65 | 25-62 |
| Age group (%) | | | | | |
| <25 years | 4 (3) | 5 (4) | 1 (1) | 2 (2) | 0 (0) |
| 25+ years | 123 (97) | 123 (96) | 103 (99) | 98 (98) | 100 (100) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Median age first injection | | | | | |
| <i>Median age first injection</i> | 18 | 18 | 18 | 19 | 19 |
| <i>Age range</i> | 10-50 | 11-43 | 11-53 | 12-65 | 10-60 |
| <i>N° not reported</i> | 5 | 3 | 3 | 4 | 3 |
| Median yrs since first injection | | | | | |
| <i>Median yrs since first injection</i> | 21 | 24 | 24 | 23 | 23 |
| <i>Range</i> | <1-46 | <1-46 | <1-47 | <1-43 | <1-42 |
| Years since first injection | | | | | |
| <3 years | 3 (2) | 8 (6) | 4 (4) | 10 (10) | 8 (8) |
| 3+ years | 119 (94) | 117 (91) | 97 (93) | 86 (86) | 89 (89) |
| Not reported | 5 (4) | 3 (2) | 3 (3) | 4 (4) | 3 (3) |
| Imprisonment last year (%) | | | | | |
| No | 99 (78) | 104 (81) | 84 (81) | 81 (81) | 89 (89) |
| Yes | 11 (9) | 19 (15) | 15 (14) | 12 (12) | 9 (9) |
| Not reported | 17 (13) | 5 (4) | 5 (5) | 7 (7) | 2 (2) |
| N° in prison | | | | | |
| <i>N° in prison</i> | N=11 | N=19 | N=15 | N=12 | N=9 |
| Injected in prison | 5 (45) | 5 (26) | 3 (20) | 8 (67) | 3 (33) |

* Implementation of the 2020 and 2021 ANSPS was impacted by the COVID-19 pandemic, see page 1 for more information

Table 2.1.2 Number (percentage) of respondents by last drug injected, frequency of injecting and survey year

| Australian Capital Territory | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|----------|----------|---------|---------|---------|
| N° surveyed | N=127 | N=128 | N=104 | N=100 | N=100 |
| Drug last injected (%) | | | | | |
| Cocaine | 1 (1) | 0 (0) | 2 (2) | 1 (1) | 0 (0) |
| Methamphetamine | 60 (47) | 58 (45) | 39 (38) | 40 (40) | 42 (42) |
| Heroin | 51 (40) | 59 (46) | 50 (48) | 44 (44) | 44 (44) |
| Pharm. opioids | 4 (3) | 2 (2) | 3 (3) | 0 (0) | 0 (0) |
| Methadone | 3 (2) | 5 (4) | 4 (4) | 6 (6) | 2 (2) |
| Buprenorphine | 0 (0) | 0 (0) | 1 (1) | 0 (0) | 0 (0) |
| Buprenorphine/naloxone | 1 (1) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| PIEDs | 0 (0) | 0 (0) | 1 (1) | 1 (1) | 1 (1) |
| More than one | 5 (4) | 4 (3) | 3 (3) | 7 (7) | 7 (7) |
| Other | 1 (1) | 0 (0) | 0 (0) | 0 (0) | 3 (3) |
| Not reported | 1 (1) | 0 (0) | 1 (1) | 1 (1) | 1 (1) |
| Ever injected opioids | | | | | |
| No | 15 (12) | 8 (6) | 11 (11) | 15 (15) | 14 (14) |
| Yes | 110 (87) | 120 (94) | 93 (89) | 84 (84) | 86 (86) |
| Not reported | 2 (2) | 0 (0) | 0 (0) | 1 (1) | 0 (0) |
| Frequency of injection last month (%) | | | | | |
| Not last month | 11 (9) | 10 (8) | 7 (7) | 7 (7) | 3 (3) |
| Less than weekly | 17 (13) | 18 (14) | 14 (13) | 16 (16) | 17 (17) |
| Weekly not daily | 30 (24) | 32 (25) | 24 (23) | 17 (17) | 24 (24) |
| Daily or more | 66 (52) | 66 (52) | 59 (57) | 58 (58) | 54 (54) |
| Not reported | 3 (2) | 2 (2) | 0 (0) | 2 (2) | 2 (2) |
| Present when someone injected for the first time in the last 12 months (%) | | | | | |
| No | 103 (81) | 111 (87) | 89 (86) | 83 (83) | 90 (90) |
| Yes | 20 (16) | 15 (12) | 11 (11) | 16 (16) | 9 (9) |
| Not reported | 4 (3) | 2 (2) | 4 (4) | 1 (1) | 1 (1) |
| Overdosed in last 12 months | | | | | |
| No | 93 (73) | 101 (79) | 77 (74) | 79 (79) | 80 (80) |
| Yes | 30 (24) | 26 (20) | 23 (22) | 19 (19) | 18 (18) |
| Not reported | 4 (3) | 1 (1) | 4 (4) | 2 (2) | 2 (2) |

Table 2.1.3 Number (percentage) of respondents by injecting behaviour in the last month and survey year

| Australian Capital Territory | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|----------|----------|---------|---------|---------|
| N° injected last month | N=113 | N=116 | N=97 | N=91 | N=95 |
| Places injected last month (%) | | | | | |
| All private | 50 (44) | 53 (46) | 39 (40) | 56 (62) | 52 (55) |
| Any public | 63 (56) | 61 (53) | 58 (60) | 35 (38) | 43 (45) |
| Not reported | 0 (0) | 2 (2) | 0 (0) | 0 (0) | 0 (0) |
| Use of new and sterile needles and syringes last month (%) | | | | | |
| All injections | 76 (67) | 86 (74) | 73 (75) | 70 (77) | 74 (78) |
| Most of the time | 30 (27) | 25 (22) | 18 (19) | 20 (22) | 17 (18) |
| Half of the time | 4 (4) | 2 (2) | 3 (3) | 0 (0) | 3 (3) |
| Some of the time | 2 (2) | 0 (0) | 0 (0) | 1 (1) | 1 (1) |
| Not last month | 0 (0) | 1 (1) | 1 (1) | 0 (0) | 0 (0) |
| Not reported | 1 (1) | 2 (2) | 2 (2) | 0 (0) | 0 (0) |
| Re-used someone else's used needle & syringe last month (%) | | | | | |
| None | 83 (73) | 99 (85) | 79 (81) | 71 (78) | 80 (84) |
| Once | 8 (7) | 7 (6) | 8 (8) | 4 (4) | 1 (1) |
| Twice | 9 (8) | 3 (3) | 1 (1) | 6 (7) | 4 (4) |
| 3-5 times | 6 (5) | 4 (3) | 7 (7) | 1 (1) | 5 (5) |
| >5 times | 6 (5) | 2 (2) | 1 (1) | 8 (9) | 5 (5) |
| Not reported | 1 (1) | 1 (1) | 1 (1) | 1 (1) | 0 (0) |
| Number of people needle & syringe was re-used after last month (%) | | | | | |
| None | 83 (73) | 99 (85) | 79 (81) | 71 (78) | 80 (84) |
| One | 11 (10) | 4 (3) | 8 (8) | 7 (8) | 5 (5) |
| Two | 1 (1) | 3 (3) | 1 (1) | 3 (3) | 3 (3) |
| Three to five | 2 (2) | 1 (1) | 0 (0) | 0 (0) | 2 (2) |
| More than five | 2 (2) | 0 (0) | 0 (0) | 2 (2) | 0 (0) |
| Don't know | 10 (9) | 5 (4) | 8 (8) | 6 (7) | 3 (3) |
| Not reported | 4 (4) | 4 (3) | 1 (1) | 2 (2) | 2 (2) |
| Relationship to people needle & syringe was used after last month (%) [more than one could be selected] | | | | | |
| Regular sex partner | 10 (9) | 5 (4) | 3 (3) | 8 (9) | 7 (7) |
| Casual sex partner | 0 (0) | 1 (1) | 1 (1) | 1 (1) | 0 (0) |
| Close friend | 5 (4) | 4 (3) | 9 (9) | 4 (4) | 1 (1) |
| Acquaintance | 5 (4) | 2 (2) | 4 (4) | 2 (2) | 2 (2) |
| Other | 1 (1) | 3 (3) | 0 (0) | 1 (1) | 3 (3) |
| Equipment used after someone else last month (%) [more than one could be selected] | | | | | |
| Spoon | 33 (29) | 27 (23) | 24 (25) | 29 (32) | 29 (31) |
| Water | 21 (19) | 14 (12) | 12 (12) | 21 (23) | 14 (15) |
| Filter | 11 (10) | 7 (6) | 9 (9) | 12 (13) | 9 (9) |
| Drug mix | 14 (12) | 9 (8) | 8 (8) | 11 (12) | 16 (17) |
| None | 74 (65) | 79 (68) | 65 (67) | 58 (64) | 63 (66) |
| Injected by someone after that person injected themselves or others last month (%) | | | | | |
| No | 90 (80) | 102 (88) | 86 (89) | 74 (81) | 82 (86) |
| Yes | 23 (20) | 13 (11) | 11 (11) | 16 (18) | 13 (14) |
| Not reported | 0 (0) | 1 (1) | 0 (0) | 1 (1) | 0 (0) |
| Source of needle acquisition | | | | | |
| Needle Syringe Program | 101 (89) | 108 (93) | 86 (89) | 75 (82) | 84 (88) |
| Chemist/Pharmacy | 37 (33) | 33 (28) | 26 (27) | 25 (27) | 23 (24) |
| Personal sources | 24 (21) | 17 (15) | 17 (18) | 14 (15) | 19 (20) |
| Dispensing/Vending Machine | 29 (26) | 32 (28) | 17 (18) | 9 (10) | 16 (17) |
| Other sources | 5 (4) | 3 (3) | 3 (3) | 3 (3) | 2 (2) |

Table 2.1.4 Number (percentage) of respondents by treatment for drug use and survey year

| Australian Capital Territory | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|---------|----------|---------|---------|---------|
| N° surveyed | N=127 | N=128 | N=104 | N=100 | N=100 |
| Ever any treatment/therapy for drug use (%) | | | | | |
| No | 33 (26) | 26 (20) | 21 (20) | 21 (21) | 28 (28) |
| Yes | 93 (73) | 101 (80) | 83 (80) | 79 (79) | 72 (72) |
| Not reported | 1 (<1) | 1 (<1) | 0 (0) | 0 (0) | 0 (0) |
| History of methadone maintenance treatment (%) | | | | | |
| Currently | 33 (26) | 49 (38) | 33 (32) | 39 (39) | 37 (37) |
| Previously | 29 (23) | 32 (25) | 23 (22) | 26 (26) | 18 (18) |
| Never | 58 (46) | 43 (34) | 45 (43) | 33 (33) | 44 (44) |
| Not reported | 7 (6) | 4 (3) | 3 (3) | 2 (2) | 1 (1) |
| History of buprenorphine (Subutex®) treatment (%) | | | | | |
| Currently | 7 (6) | 5 (4) | 7 (7) | 5 (5) | 9 (9) |
| Previously | 31 (24) | 29 (23) | 35 (34) | 38 (38) | 28 (28) |
| Never | 86 (68) | 91 (71) | 60 (58) | 56 (56) | 63 (63) |
| Not reported | 3 (2) | 3 (2) | 2 (2) | 1 (1) | 0 (0) |
| History of buprenorphine-naloxone (Suboxone®) treatment (%) | | | | | |
| Currently | 8 (6) | 9 (7) | 3 (3) | 5 (5) | 5 (5) |
| Previously | 33 (26) | 28 (22) | 29 (28) | 34 (34) | 31 (31) |
| Never | 83 (65) | 88 (69) | 72 (69) | 60 (60) | 63 (63) |
| Not reported | 3 (2) | 3 (2) | 0 (0) | 1 (1) | 1 (1) |

Table 2.1.5 Number (percentage) of respondents by testing for HIV and HCV, HCV treatment by survey year

| Australian Capital Territory | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|----------|----------|---------|---------|---------|
| N° surveyed | N=127 | N=128 | N=104 | N=100 | N=100 |
| Previous HIV test (%) | | | | | |
| Yes, ever | 102 (80) | 113 (88) | 85 (82) | 82 (82) | 83 (83) |
| Yes, last year | 62 (49) | 70 (55) | 49 (47) | 44 (44) | 51 (51) |
| >1 year ago | 40 (32) | 43 (34) | 36 (35) | 38 (38) | 32 (32) |
| Never tested | 15 (12) | 12 (9) | 14 (13) | 16 (16) | 17 (17) |
| Not reported | 10 (8) | 3 (2) | 5 (5) | 2 (2) | 0 (0) |
| Previous HCV test (%) | | | | | |
| Yes, ever | 104 (82) | 116 (91) | 87 (84) | 80 (80) | 85 (85) |
| Yes, last year | 68 (54) | 80 (63) | 48 (46) | 46 (46) | 58 (58) |
| >1 year ago | 36 (28) | 36 (28) | 39 (38) | 34 (34) | 27 (27) |
| Never tested | 11 (9) | 4 (3) | 8 (8) | 7 (7) | 11 (11) |
| Not reported | 12 (9) | 8 (6) | 9 (9) | 13 (13) | 4 (4) |
| Lifetime treatment for HCV (%)# | | | | | |
| N° self-reported HCV diagnosis | N=46 | N=50 | N=29 | N=24 | N=23 |
| Antiviral treatment | 21 (46) | 33 (66) | 16 (55) | 11 (46) | 16 (70) |
| No antiviral treatment | 25 (54) | 17 (34) | 12 (41) | 13 (54) | 7 (30) |
| Not reported | 0 (0) | 0 (0) | 1 (3) | 0 (0) | 0 (0) |
| Treatment for HCV in past 12 months (%)** | | | | | |
| N° self-reported HCV diagnosis | N=40 | N=37 | N=18 | N=18 | N=15 |
| Antiviral treatment | 14 (35) | 17 (46) | 2 (11) | 4 (22) | 6 (40) |
| No antiviral treatment | 26 (65) | 20 (54) | 15 (83) | 14 (78) | 9 (60) |
| Not reported | 0 (0) | 0 (0) | 1 (6) | 0 (0) | 0 (0) |

among people who tested HCV antibody positive and did not report spontaneous clearance

* excludes people who reported treatment induced clearance more than 12 months previously

Table 2.1.6 Number (percentage) of respondents by sexual behaviour in the month prior to survey and survey year

| Australian Capital Territory | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|----------|----------|---------|---------|---------|
| N° surveyed | N=127 | N=128 | N=104 | N=100 | N=100 |
| Sex with a regular partner last month (%) | | | | | |
| No | 75 (59) | 78 (61) | 69 (66) | 62 (62) | 59 (59) |
| Yes | 43 (34) | 46 (36) | 31 (30) | 35 (35) | 41 (41) |
| Not reported | 9 (7) | 4 (3) | 4 (4) | 3 (3) | 0 (0) |
| Condom used with regular partner last month (%) | | | | | |
| N° surveyed | N=43 | N=46 | N=31 | N=35 | N=41 |
| Never | 34 (79) | 38 (83) | 25 (81) | 23 (66) | 32 (78) |
| Sometimes | 4 (9) | 3 (7) | 4 (13) | 1 (3) | 5 (12) |
| Every time | 2 (5) | 3 (7) | 1 (3) | 6 (17) | 3 (7) |
| Not reported | 3 (7) | 2 (4) | 1 (3) | 5 (14) | 1 (2) |
| Sex with other partner(s) last month (%) | | | | | |
| No | 103 (81) | 115 (90) | 95 (91) | 86 (86) | 91 (91) |
| Yes | 12 (9) | 11 (9) | 5 (5) | 10 (10) | 5 (5) |
| Not reported | 12 (9) | 2 (2) | 4 (4) | 4 (4) | 4 (4) |
| Condom used with other partner(s) last month (%) | | | | | |
| N° surveyed | N=12 | N=11 | N=5 | N=10 | N=5 |
| Never | 7 (58) | 7 (64) | 2 (40) | 5 (50) | 3 (60) |
| Sometimes | 0 (0) | 1 (9) | 3 (60) | 2 (20) | 0 (0) |
| Every time | 4 (33) | 2 (18) | 0 (0) | 3 (30) | 2 (40) |
| Not reported | 1 (8) | 1 (9) | 0 (0) | 0 (0) | 0 (0) |
| Sex work last month (%) | | | | | |
| No | 108 (85) | 117 (91) | 97 (93) | 93 (93) | 98 (98) |
| Yes | 8 (6) | 5 (4) | 4 (4) | 5 (5) | 2 (2) |
| Not reported | 11 (9) | 6 (5) | 3 (3) | 2 (2) | 0 (0) |
| Condom used at last sex work (%) | | | | | |
| Yes | 7 (88) | 4 (80) | 1 (25) | 1 (20) | 2 (100) |

Table 2.1.7 Number (percentage) of respondents by COVID-19 related impacts, 2020-2022

| Australian Capital Territory | 2020 | 2021 | 2022 |
|---|---------|---------|---------|
| N° surveyed | N=104 | N=100 | N=100 |
| COVID-19 test | | | |
| Yes | 27 (26) | 56 (56) | 89 (89) |
| No | 71 (68) | 23 (23) | 10 (10) |
| Don't know | 5 (5) | 5 (5) | -- -- |
| Not reported | 1 (1) | 16 (16) | 1 (1) |
| Self-reported positive COVID-19 test result | | | |
| Yes | 0 (0) | 10 (10) | 33 (37) |
| Yes, last 12 months | -- -- | -- -- | 28 (31) |
| Yes, more than a year ago | -- -- | -- -- | 5 (6) |
| COVID-19 vaccine uptake | | | |
| ≥1 dose | -- -- | 82 (82) | 85 (85) |
| COVID-19 vaccine doses | | | |
| One dose | -- -- | 11 (11) | 5 (5) |
| Two doses | -- -- | 71 (71) | 27 (27) |
| Three or more doses | -- -- | -- -- | 53 (53) |
| No doses | -- -- | 12 (12) | 14 (14) |
| Not reported | -- -- | 6 (6) | 1 (1) |
| Vaccine type * | | | N=85 |
| Astra Zeneca | -- -- | 17 (17) | 30 (35) |
| Pfizer | -- -- | 46 (46) | 46 (54) |
| Don't know | -- -- | 4 (4) | 16 (19) |
| Other | -- -- | -- -- | 14 (16) |
| Not reported | -- -- | 33 (33) | -- -- |
| * 2021 only one response permitted, 2022 multiple responses permitted | | | |
| Last vaccine location | | | N=100 |
| GP or medical centre | -- -- | -- -- | 46 (46) |
| Pharmacy/ chemist | -- -- | -- -- | 14 (14) |
| Vaccine hub | -- -- | -- -- | 7 (7) |
| Opioid treatment/ methadone clinic | -- -- | -- -- | 0 (0) |
| NSP | -- -- | -- -- | 5 (5) |
| Prison | -- -- | -- -- | 4 (4) |
| Other | -- -- | -- -- | 4 (4) |
| Multiple response | -- -- | -- -- | 9 (9) |
| Not reported | -- -- | -- -- | 11 (11) |
| Vaccine hesitancy - if not had three or more doses | | | |
| Haven't got around to it | -- -- | -- -- | 19 (19) |
| Difficulty booking appointment | -- -- | -- -- | 2 (2) |
| Concerns about side effects | -- -- | -- -- | 4 (4) |
| Concerns about vaccine safety | -- -- | -- -- | 3 (3) |
| Not at risk of serious illness | -- -- | -- -- | 11 (11) |
| Other | -- -- | -- -- | 18 (18) |
| Multiple response | -- -- | -- -- | 4 (4) |
| Not reported | -- -- | -- -- | 39 (39) |

HIV antibody prevalence

Table 2.2.1 HIV antibody prevalence by gender and survey year

| Aust. Capital Territory Survey year | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | 90 | 0 (0.0) | 35 | 0 (0.0) | 127 | 0 (0.0) |
| 2019 | 84 | 0 (0.0) | 42 | 0 (0.0) | 126 | 0 (0.0) |
| 2020 | 69 | 0 (0.0) | 33 | 0 (0.0) | 103 | 0 (0.0) |
| 2021 | 67 | 0 (0.0) | 32 | 0 (0.0) | 99 | 0 (0.0) |
| 2022 | 66 | 1 (1.5) | 32 | 0 (0.0) | 99 | 1 (1.0) |
| X ² p trend | | 0.127 | | -- | | 0.130 |

HCV antibody prevalence

Table 2.3.1 HCV antibody prevalence by gender and survey year

| Aust. Capital Territory Survey year | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | 90 | 51 (57) | 35 | 22 (63) | 127 | 73 (57) |
| 2019 | 84 | 38 (45) | 42 | 23 (55) | 126 | 61 (48) |
| 2020 | 69 | 32 (46) | 33 | 19 (58) | 103 | 51 (50) |
| 2021 | 67 | 27 (40) | 32 | 9 (28) | 99 | 36 (36) |
| 2022 | 67 | 25 (37) | 32 | 10 (31) | 100 | 35 (35) |
| X ² p trend | | 0.137 | | 0.001 | | 0.001 |

Table 2.3.2 HCV antibody prevalence by sexual identity, gender and survey year

| Aust. Capital Territory Sexual identity | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heterosexual | 77 | 43 (56) | 22 | 14 (64) | 100 | 57 (57) |
| Bisexual | 6 | 2 (33) | 7 | 4 (57) | 13 | 6 (46) |
| Homosexual | 2 | 1 (50) | 0 | 0 (0) | 3 | 1 (33) |
| p value | | 0.701 | | 1.000 | | 0.594 |
| 2019 | | | | | | |
| Heterosexual | 64 | 28 (44) | 27 | 15 (56) | 91 | 43 (47) |
| Bisexual | 5 | 3 (60) | 8 | 5 (63) | 13 | 8 (62) |
| Homosexual | 3 | 3 (100) | 1 | 0 (0) | 4 | 3 (75) |
| p value | | 0.158 | | 0.495 | | 0.401 |
| 2020 | | | | | | |
| Heterosexual | 63 | 28 (44) | 21 | 12 (57) | 84 | 40 (48) |
| Bisexual | 5 | 3 (60) | 5 | 3 (60) | 10 | 6 (60) |
| Homosexual | 0 | 0 (0) | 2 | 1 (50) | 2 | 1 (50) |
| p value | | 0.653 | | 0.580 | | 0.381 |
| 2021 | | | | | | |
| Heterosexual | 62 | 25 (40) | 24 | 7 (29) | 86 | 32 (37) |
| Bisexual | 3 | 2 (67) | 4 | 2 (50) | 7 | 4 (57) |
| Homosexual | 2 | 0 (0) | 2 | 0 (0) | 4 | 0 (0) |
| p value | | 0.490 | | 0.717 | | 0.260 |
| 2022 | | | | | | |
| Heterosexual | 60 | 23 (38) | 21 | 7 (33) | 81 | 30 (37) |
| Bisexual | 3 | 0 (0) | 8 | 2 (25) | 11 | 2 (18) |
| Homosexual | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | 0.124 | | 0.557 | | 0.187 |

Table 2.3.3 HCV antibody prevalence by age group, gender and survey year

| Aust. Capital Territory Age group | Male | | Female | | Total | |
|--------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <25 years | 2 | 0 (0) | 2 | 1 (50) | 4 | 1 (25) |
| 25-34 years | 13 | 7 (54) | 9 | 6 (67) | 22 | 13 (59) |
| 35-44 years | 40 | 28 (70) | 14 | 8 (57) | 56 | 36 (64) |
| 45+ years | 35 | 16 (46) | 10 | 7 (70) | 45 | 23 (51) |
| p value | | 0.056 | | 0.952 | | 0.321 |
| 2019 | | | | | | |
| <25 years | 1 | 0 (0) | 4 | 2 (50) | 5 | 2 (40) |
| 25-34 years | 9 | 1 (11) | 10 | 6 (60) | 19 | 7 (37) |
| 35-44 years | 33 | 16 (48) | 14 | 6 (43) | 47 | 22 (47) |
| 45+ years | 41 | 21 (51) | 14 | 9 (64) | 55 | 30 (55) |
| p value | | 0.097 | | 0.777 | | 0.602 |
| 2020 | | | | | | |
| <25 years | 0 | 0 (0) | 1 | 1 (100) | 1 | 1 (100) |
| 25-34 years | 9 | 2 (22) | 5 | 4 (80) | 15 | 6 (40) |
| 35-44 years | 23 | 8 (35) | 15 | 8 (53) | 38 | 16 (42) |
| 45+ years | 37 | 22 (59) | 12 | 6 (50) | 49 | 28 (57) |
| p value | | 0.059 | | 0.668 | | 0.298 |
| 2021 | | | | | | |
| <25 years | 0 | 0 (0) | 2 | 1 (50) | 2 | 1 (50) |
| 25-34 years | 15 | 5 (33) | 5 | 1 (20) | 20 | 6 (30) |
| 35-44 years | 23 | 11 (48) | 11 | 4 (36) | 34 | 15 (44) |
| 45+ years | 29 | 11 (38) | 14 | 3 (21) | 43 | 14 (33) |
| p value | | 0.634 | | 0.764 | | 0.579 |
| 2022 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 8 | 1 (13) | 9 | 1 (11) | 18 | 2 (11) |
| 35-44 years | 23 | 11 (48) | 8 | 2 (25) | 31 | 13 (42) |
| 45+ years | 36 | 13 (36) | 15 | 7 (47) | 51 | 20 (39) |
| p value | | 0.232 | | 0.204 | | 0.053 |

Table 2.3.4 HCV antibody prevalence by years since first injection, gender and survey year

| Aust. Capital Territory Years of injection | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <3 years | 3 | 2 (67) | 0 | 0 (0) | 3 | 2 (67) |
| 3 to 10 years | 9 | 2 (22) | 7 | 4 (57) | 16 | 6 (38) |
| 11+ years | 74 | 43 (58) | 27 | 17 (63) | 103 | 60 (58) |
| p value | | 0.106 | | 1.000 | | 0.286 |
| 2019 | | | | | | |
| <3 years | 3 | 0 (0) | 5 | 0 (0) | 8 | 0 (0) |
| 3 to 10 years | 6 | 1 (17) | 6 | 5 (83) | 12 | 6 (50) |
| 11+ years | 73 | 36 (49) | 30 | 17 (57) | 103 | 53 (51) |
| p value | | 0.108 | | 0.016 | | 0.014 |
| 2020 | | | | | | |
| <3 years | 3 | 0 (0) | 1 | 0 (0) | 4 | 0 (0) |
| 3 to 10 years | 6 | 1 (17) | 2 | 2 (100) | 8 | 3 (38) |
| 11+ years | 59 | 31 (53) | 28 | 15 (54) | 88 | 46 (52) |
| p value | | 0.069 | | 0.337 | | 0.122 |
| 2021 | | | | | | |
| <3 years | 5 | 1 (20) | 5 | 2 (40) | 10 | 3 (30) |
| 3 to 10 years | 4 | 1 (25) | 4 | 1 (25) | 8 | 2 (25) |
| 11+ years | 57 | 24 (42) | 20 | 6 (30) | 77 | 30 (39) |
| p value | | 0.649 | | 1.000 | | 0.785 |
| 2022 | | | | | | |
| <3 years | 4 | 0 (0) | 4 | 0 (0) | 8 | 0 (0) |
| 3 to 10 years | 6 | 0 (0) | 6 | 3 (50) | 12 | 3 (25) |
| 11+ years | 57 | 25 (44) | 20 | 6 (30) | 77 | 31 (40) |
| p value | | 0.031 | | 0.344 | | 0.046 |

Table 2.3.5 HCV antibody prevalence by re-use of someone else's used needle and syringe last month, years since first injection and survey year

| Aust. Capital Territory Receptively shared syringe last month | <3 years | | 3+ years | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| No receptive sharing | 2 | 1 (50) | 79 | 46 (58) | 83 | 49 (59) |
| Receptive sharing | 1 | 1 (100) | 27 | 15 (56) | 29 | 17 (59) |
| p value | | 1.000 | | 0.808 | | 0.969 |
| 2019 | | | | | | |
| No receptive sharing | 8 | 0 (0) | 87 | 45 (52) | 97 | 46 (47) |
| Receptive sharing | 0 | 0 (0) | 15 | 8 (53) | 16 | 9 (56) |
| p value | | -- | | 0.908 | | 0.513 |
| 2020 | | | | | | |
| No receptive sharing | 4 | 0 (0) | 73 | 36 (49) | 78 | 36 (46) |
| Receptive sharing | 0 | 0 (0) | 15 | 9 (60) | 17 | 11 (65) |
| p value | | -- | | 0.451 | | 0.166 |
| 2021 | | | | | | |
| No receptive sharing | 7 | 2 (29) | 60 | 24 (40) | 70 | 27 (39) |
| Receptive sharing | 2 | 1 (50) | 17 | 8 (47) | 19 | 9 (47) |
| p value | | 1.000 | | 0.602 | | 0.488 |
| 2022 | | | | | | |
| No receptive sharing | 7 | 0 (0) | 70 | 24 (34) | 80 | 25 (31) |
| Receptive sharing | 0 | 0 (0) | 15 | 8 (53) | 15 | 8 (53) |
| p value | | -- | | 0.167 | | 0.099 |

Table 2.3.6 HCV antibody prevalence by last drug injected, years since first injection and survey year

| Aust. Capital Territory Last drug injected | <3 years | | 3+ years | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heroin | 0 | 0 (0) | 49 | 28 (57) | 51 | 30 (59) |
| Methamphetamine | 2 | 1 (50) | 54 | 28 (52) | 58 | 31 (53) |
| Other opioids | 0 | 0 (0) | 7 | 5 (71) | 7 | 5 (71) |
| Other drugs | 1 | 1 (100) | 6 | 5 (83) | 7 | 6 (86) |
| p value | | 0.386 | | 0.411 | | 0.358 |
| 2019 | | | | | | |
| Heroin | 2 | 0 (0) | 53 | 32 (60) | 57 | 33 (58) |
| Methamphetamine | 5 | 0 (0) | 50 | 19 (38) | 56 | 20 (36) |
| Other opioids | 1 | 0 (0) | 6 | 3 (50) | 7 | 3 (43) |
| Other drugs | 0 | 0 (0) | 4 | 3 (75) | 4 | 3 (75) |
| p value | | -- | | 0.104 | | 0.076 |
| 2020 | | | | | | |
| Heroin | 1 | 0 (0) | 46 | 28 (61) | 50 | 30 (60) |
| Methamphetamine | 3 | 0 (0) | 35 | 16 (46) | 38 | 16 (42) |
| Other opioids | 0 | 0 (0) | 8 | 3 (38) | 8 | 3 (38) |
| Other drugs | 0 | 0 (0) | 7 | 2 (29) | 7 | 2 (29) |
| p value | | -- | | 0.243 | | 0.193 |
| 2021 | | | | | | |
| Heroin | 2 | 0 (0) | 39 | 14 (36) | 43 | 15 (35) |
| Methamphetamine | 5 | 0 (0) | 32 | 12 (38) | 39 | 12 (31) |
| Other opioids | 2 | 2 (100) | 3 | 1 (33) | 5 | 3 (60) |
| Other drugs | 0 | 0 (0) | 9 | 5 (56) | 9 | 5 (56) |
| p value | | 0.011 | | 0.738 | | 0.361 |
| 2022 | | | | | | |
| Heroin | 1 | 0 (0) | 42 | 22 (52) | 44 | 22 (50) |
| Methamphetamine | 6 | 0 (0) | 33 | 7 (21) | 41 | 8 (20) |
| Other opioids | 0 | 0 (0) | 2 | 1 (50) | 2 | 1 (50) |
| Other drugs | 1 | 0 (0) | 9 | 2 (22) | 10 | 2 (20) |
| p value | | -- | | 0.033 | | 0.019 |

Table 2.3.7 HCV antibody prevalence by frequency of drug injection last month, years since first injection and survey year

| Aust. Capital Territory Frequency of injection last month | <3 years | | 3+ years | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| Less than daily | 3 | 2 (67) | 41 | 22 (54) | 47 | 27 (57) |
| Daily or more | 0 | 0 (0) | 66 | 40 (61) | 66 | 40 (61) |
| Not last month | 0 | 0 (0) | 9 | 4 (44) | 11 | 6 (55) |
| p value | | -- | | 0.541 | | 0.902 |
| 2019 | | | | | | |
| Less than daily | 6 | 0 (0) | 42 | 22 (52) | 50 | 24 (48) |
| Daily or more | 2 | 0 (0) | 61 | 31 (51) | 64 | 31 (48) |
| Not last month | 0 | 0 (0) | 10 | 4 (40) | 10 | 4 (40) |
| p value | | -- | | 0.850 | | 0.926 |
| 2020 | | | | | | |
| Less than daily | 1 | 0 (0) | 36 | 11 (31) | 37 | 11 (30) |
| Daily or more | 3 | 0 (0) | 53 | 35 (66) | 59 | 37 (63) |
| Not last month | 0 | 0 (0) | 7 | 3 (43) | 7 | 3 (43) |
| p value | | -- | | 0.003 | | 0.005 |
| 2021 | | | | | | |
| Less than daily | 4 | 1 (25) | 27 | 5 (19) | 32 | 6 (19) |
| Daily or more | 5 | 2 (40) | 51 | 27 (53) | 58 | 30 (52) |
| Not last month | 1 | 0 (0) | 5 | 0 (0) | 7 | 0 (0) |
| p value | | 1.000 | | 0.001 | | 0.001 |
| 2022 | | | | | | |
| Less than daily | 4 | 0 (0) | 35 | 15 (43) | 41 | 16 (39) |
| Daily or more | 3 | 0 (0) | 50 | 17 (34) | 54 | 17 (31) |
| Not last month | 1 | 0 (0) | 2 | 1 (50) | 3 | 1 (33) |
| p value | | -- | | 0.627 | | 0.782 |

Table 2.3.8 HCV antibody prevalence by imprisonment last year, years since first injection and survey year

| Aust. Capital Territory Imprisonment last year | <3 years | | 3+ years | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| No imprisonment | 2 | 1 (50) | 95 | 50 (53) | 99 | 53 (54) |
| Imprisonment | 0 | 0 (0) | 10 | 6 (60) | 11 | 7 (64) |
| p value | | -- | | 0.748 | | 0.751 |
| 2019 | | | | | | |
| No imprisonment | 6 | 0 (0) | 95 | 50 (53) | 102 | 51 (50) |
| Imprisonment | 2 | 0 (0) | 15 | 6 (40) | 19 | 7 (37) |
| p value | | -- | | 0.363 | | 0.292 |
| 2020 | | | | | | |
| No imprisonment | 3 | 0 (0) | 80 | 38 (48) | 83 | 38 (46) |
| Imprisonment | 1 | 0 (0) | 12 | 7 (58) | 15 | 9 (60) |
| p value | | -- | | 0.310 | | 0.310 |
| 2021 | | | | | | |
| No imprisonment | 8 | 3 (38) | 68 | 25 (37) | 80 | 29 (36) |
| Imprisonment | 1 | 0 (0) | 11 | 3 (27) | 12 | 3 (25) |
| p value | | 1.000 | | 0.738 | | 0.532 |
| 2022 | | | | | | |
| No imprisonment | 8 | 0 (0) | 78 | 30 (38) | 89 | 31 (35) |
| Imprisonment | 0 | 0 (0) | 9 | 3 (33) | 9 | 3 (33) |
| p value | | -- | | 1.000 | | 1.000 |

Table 2.3.9 HCV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Aust. Capital Territory Aboriginal and Torres Strait Islander origin | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Non Indigenous | 74 | 41 (55) | 22 | 12 (55) | 98 | 53 (54) |
| Indigenous | 15 | 9 (60) | 12 | 9 (75) | 27 | 18 (67) |
| p value | | 0.744 | | 0.292 | | 0.242 |
| 2019 | | | | | | |
| Non Indigenous | 73 | 34 (47) | 31 | 16 (52) | 104 | 50 (48) |
| Indigenous | 10 | 4 (40) | 11 | 7 (64) | 21 | 11 (52) |
| p value | | 0.748 | | 0.726 | | 0.719 |
| 2020 | | | | | | |
| Non Indigenous | 57 | 22 (39) | 23 | 11 (48) | 81 | 33 (41) |
| Indigenous | 10 | 8 (80) | 9 | 7 (78) | 19 | 15 (79) |
| p value | | 0.034 | | 0.235 | | 0.004 |
| 2021 | | | | | | |
| Non Indigenous | 49 | 18 (37) | 19 | 5 (26) | 68 | 23 (34) |
| Indigenous | 16 | 9 (56) | 12 | 3 (25) | 28 | 12 (43) |
| p value | | 0.169 | | 1.000 | | 0.403 |
| 2022 | | | | | | |
| Non Indigenous | 53 | 21 (40) | 18 | 6 (33) | 71 | 27 (38) |
| Indigenous | 13 | 3 (23) | 13 | 3 (23) | 27 | 6 (22) |
| p value | | 0.345 | | 0.696 | | 0.139 |

Table 2.3.10 HCV antibody prevalence by main language spoken at home by parents, gender and survey year

| Aust. Capital Territory Main language spoken at home by parents | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| English speaking | 81 | 47 (58) | 31 | 20 (65) | 114 | 67 (59) |
| Non-English speaking | 7 | 3 (43) | 4 | 2 (50) | 11 | 5 (45) |
| p value | | 0.459 | | 0.618 | | 0.393 |
| 2019 | | | | | | |
| English speaking | 78 | 38 (49) | 39 | 20 (51) | 117 | 58 (50) |
| Non-English speaking | 6 | 0 (0) | 3 | 3 (100) | 9 | 3 (33) |
| p value | | 0.030 | | 0.102 | | 0.494 |
| 2020 | | | | | | |
| English speaking | 64 | 29 (45) | 33 | 19 (58) | 98 | 48 (49) |
| Non-English speaking | 5 | 3 (60) | 0 | 0 (0) | 5 | 3 (60) |
| p value | | 0.657 | | -- | | 0.678 |
| 2021 | | | | | | |
| English speaking | 65 | 26 (40) | 29 | 7 (24) | 94 | 33 (35) |
| Non-English speaking | 2 | 1 (50) | 3 | 2 (67) | 5 | 3 (60) |
| p value | | 1.000 | | 0.184 | | 0.350 |
| 2022 | | | | | | |
| English speaking | 63 | 22 (35) | 31 | 10 (32) | 94 | 32 (34) |
| Non-English speaking | 3 | 2 (67) | 1 | 0 (0) | 4 | 2 (50) |
| p value | | 0.548 | | 1.000 | | 0.608 |

Table 2.3.11 HCV antibody prevalence by region/country of birth, gender and survey year

| Aust. Capital Territory | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Australia | 76 | 45 (59) | 32 | 21 (66) | 110 | 66 (60) |
| Other Oceania | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| Asia | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| UK & Ireland | 5 | 2 (40) | 1 | 0 (0) | 6 | 2 (33) |
| Other | 6 | 3 (50) | 1 | 0 (0) | 7 | 3 (43) |
| p value | | 0.472 | | 0.139 | | 0.212 |
| 2019 | | | | | | |
| Australia | 72 | 33 (46) | 39 | 21 (54) | 111 | 54 (49) |
| Other Oceania | 3 | 2 (67) | 3 | 2 (67) | 6 | 4 (67) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 4 | 2 (50) | 0 | 0 (0) | 4 | 2 (50) |
| Other | 5 | 1 (20) | 0 | 0 (0) | 5 | 1 (20) |
| p value | | 0.618 | | 1.000 | | 0.520 |
| 2020 | | | | | | |
| Australia | 60 | 27 (45) | 31 | 18 (58) | 92 | 45 (49) |
| Other Oceania | 3 | 1 (33) | 0 | 0 (0) | 3 | 1 (33) |
| Asia | 1 | 1 (100) | 0 | 0 (0) | 1 | 1 (100) |
| UK & Ireland | 1 | 1 (100) | 2 | 1 (50) | 3 | 2 (67) |
| Other | 2 | 2 (100) | 0 | 0 (0) | 2 | 2 (100) |
| p value | | 0.315 | | 1.000 | | 0.636 |
| 2021 | | | | | | |
| Australia | 62 | 23 (37) | 28 | 7 (25) | 90 | 30 (33) |
| Other Oceania | 2 | 2 (100) | 0 | 0 (0) | 2 | 2 (100) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 2 | 1 (50) | 1 | 0 (0) | 3 | 1 (33) |
| Other | 1 | 1 (100) | 3 | 2 (67) | 4 | 3 (75) |
| p value | | 0.128 | | 0.421 | | 0.070 |
| 2022 | | | | | | |
| Australia | 56 | 19 (34) | 28 | 9 (32) | 85 | 28 (33) |
| Other Oceania | 5 | 3 (60) | 0 | 0 (0) | 5 | 3 (60) |
| Asia | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) |
| UK & Ireland | 2 | 0 (0) | 2 | 0 (0) | 4 | 0 (0) |
| Other | 1 | 1 (100) | 0 | 0 (0) | 1 | 1 (100) |
| p value | | 0.402 | | 1.000 | | 0.177 |

HCV RNA prevalence

Table 2.4.1 HCV RNA prevalence by gender and survey year

| Australian Capital Territory | Male | | Female | | Total | |
|------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Survey year | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | 67 | 17 (25) | 26 | 9 (35) | 95 | 26 (27) |
| 2019* | 77 | 16 (21) | 38 | 10 (26) | 115 | 25 (22) |
| 2020 | 60 | 13 (22) | 30 | 5 (17) | 91 | 18 (20) |
| 2021 | 62 | 14 (23) | 32 | 6 (19) | 94 | 20 (21) |
| 2022 | 65 | 7 (11) | 31 | 3 (10) | 97 | 10 (10) |
| X ² p trend | | 0.080 | | 0.017 | | 0.007 |

* 2018, 2019: Weighted for gender and HCV antibody status

Totals include respondents where gender was reported as other or not reported

Table 2.4.2 HCV RNA prevalence by sexual identity, gender and survey year

| Australian Capital Territory | Male | | Female | | Total | |
|------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Sexual identity | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heterosexual | 62 | 14 (23) | 17 | 5 (29) | 80 | 20 (25) |
| Bisexual | 2 | 1 (50) | 6 | 3 (50) | 8 | 4 (50) |
| Homosexual | 1 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| p value | | 0.608 | | 0.598 | | 0.315 |
| 2019* | | | | | | |
| Heterosexual | 61 | 12 (20) | 25 | 6 (24) | 85 | 17 (20) |
| Bisexual | 4 | 1 (25) | 8 | 2 (25) | 12 | 3 (25) |
| Homosexual | 3 | 1 (33) | 1 | 0 (0) | 4 | 1 (25) |
| p value | | 0.823 | | 0.856 | | 0.924 |
| 2020 | | | | | | |
| Heterosexual | 55 | 12 (22) | 19 | 2 (11) | 74 | 14 (19) |
| Bisexual | 4 | 1 (25) | 4 | 2 (50) | 8 | 3 (38) |
| Homosexual | 0 | 0 (0) | 2 | 0 (0) | 2 | 0 (0) |
| p value | | 0.882 | | 0.120 | | 0.356 |
| 2021 | | | | | | |
| Heterosexual | 57 | 13 (23) | 24 | 4 (17) | 81 | 17 (21) |
| Bisexual | 3 | 1 (33) | 4 | 2 (50) | 7 | 3 (43) |
| Homosexual | 2 | 0 (0) | 2 | 0 (0) | 4 | 0 (0) |
| p value | | 0.676 | | 0.233 | | 0.226 |
| 2022 | | | | | | |
| Heterosexual | 58 | 6 (10) | 20 | 2 (10) | 78 | 8 (10) |
| Bisexual | 3 | 0 (0) | 8 | 1 (13) | 11 | 1 (9) |
| Homosexual | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | 0.394 | | 1.000 | | 0.533 |

Table 2.4.3 HCV RNA prevalence by age group, gender and survey year

| Australian Capital Territory | Male | | Female | | Total | |
|------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Age group | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <25 years | 2 | 0 (0) | 1 | 1 (100) | 3 | 1 (33) |
| 25-34 years | 9 | 3 (33) | 6 | 4 (67) | 16 | 8 (50) |
| 35-44 years | 29 | 7 (24) | 12 | 3 (25) | 42 | 10 (24) |
| 45+ years | 26 | 6 (23) | 7 | 1 (14) | 34 | 7 (21) |
| p value | | 0.803 | | 0.045 | | 0.212 |
| 2019* | | | | | | |
| <25 years | 1 | 0 (0) | 4 | 1 (25) | 5 | 1 (20) |
| 25-34 years | 8 | 1 (13) | 9 | 5 (56) | 17 | 6 (35) |
| 35-44 years | 32 | 7 (22) | 12 | 2 (17) | 44 | 9 (20) |
| 45+ years | 35 | 8 (23) | 13 | 2 (15) | 49 | 10 (20) |
| p value | | 0.858 | | 0.128 | | 0.650 |
| 2020 | | | | | | |
| <25 years | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| 25-34 years | 9 | 0 (0) | 5 | 2 (40) | 15 | 2 (13) |
| 35-44 years | 19 | 3 (16) | 12 | 3 (25) | 31 | 6 (19) |
| 45+ years | 32 | 10 (31) | 12 | 0 (0) | 44 | 10 (23) |
| p value | | 0.100 | | 0.160 | | 0.829 |
| 2021 | | | | | | |
| <25 years | 0 | 0 (0) | 2 | 1 (50) | 2 | 1 (50) |
| 25-34 years | 14 | 5 (36) | 5 | 1 (20) | 19 | 6 (32) |
| 35-44 years | 21 | 5 (24) | 11 | 2 (18) | 32 | 7 (22) |
| 45+ years | 27 | 4 (15) | 14 | 2 (14) | 41 | 6 (15) |
| p value | | 0.312 | | 0.689 | | 0.351 |
| 2022 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 8 | 0 (0) | 8 | 1 (13) | 17 | 1 (6) |
| 35-44 years | 22 | 3 (14) | 8 | 1 (13) | 30 | 4 (13) |
| 45+ years | 35 | 4 (11) | 15 | 1 (7) | 50 | 5 (10) |
| p value | | 0.861 | | 1.000 | | 0.820 |

Table 2.4.4 HCV RNA prevalence by years since first injection, gender and survey year

| Australian Capital Territory | Male | | Female | | Total | |
|------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Years of injection | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <3 years | 1 | 1 (100) | 0 | 0 (0) | 1 | 1 (100) |
| 3 to 10 years | 8 | 2 (25) | 4 | 3 (75) | 12 | 5 (42) |
| 11+ years | 56 | 13 (23) | 22 | 6 (27) | 79 | 20 (25) |
| p value | | 0.233 | | 0.091 | | 0.161 |
| 2019* | | | | | | |
| <3 years | 3 | 0 (0) | 5 | 0 (0) | 8 | 0 (0) |
| 3 to 10 years | 6 | 0 (0) | 6 | 4 (67) | 12 | 4 (33) |
| 11+ years | 65 | 15 (23) | 28 | 6 (21) | 93 | 20 (22) |
| p value | | 0.278 | | 0.027 | | 0.222 |
| 2020 | | | | | | |
| <3 years | 3 | 0 (0) | 1 | 0 (0) | 4 | 0 (0) |
| 3 to 10 years | 6 | 0 (0) | 2 | 1 (50) | 8 | 1 (13) |
| 11+ years | 50 | 13 (26) | 25 | 4 (16) | 76 | 17 (22) |
| p value | | 0.223 | | 0.431 | | 0.470 |
| 2021 | | | | | | |
| <3 years | 5 | 0 (0) | 5 | 1 (20) | 10 | 1 (10) |
| 3 to 10 years | 4 | 1 (25) | 4 | 1 (25) | 8 | 2 (25) |
| 11+ years | 52 | 12 (23) | 20 | 4 (20) | 72 | 16 (22) |
| p value | | 0.476 | | 0.974 | | 0.648 |
| 2022 | | | | | | |
| <3 years | 4 | 0 (0) | 4 | 0 (0) | 8 | 0 (0) |
| 3 to 10 years | 6 | 0 (0) | 6 | 1 (17) | 12 | 1 (8) |
| 11+ years | 55 | 7 (13) | 19 | 2 (11) | 74 | 9 (12) |
| p value | | 1.000 | | 1.000 | | 0.840 |

Table 2.4.5 HCV RNA prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| Australian Capital Territory | Male | | Female | | Total | |
|---------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Receptively shared syringe last month | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| No receptive sharing | 40 | 10 (25) | 20 | 8 (40) | 61 | 19 (31) |
| Receptive sharing | 19 | 5 (26) | 4 | 1 (25) | 23 | 6 (26) |
| p value | | 0.917 | | 0.554 | | 0.680 |
| 2019* | | | | | | |
| No receptive sharing | 55 | 12 (22) | 33 | 9 (27) | 88 | 20 (23) |
| Receptive sharing | 14 | 3 (21) | 1 | 0 (0) | 15 | 3 (20) |
| p value | | 0.951 | | 0.557 | | 0.732 |
| 2020 | | | | | | |
| No receptive sharing | 45 | 11 (24) | 22 | 3 (14) | 68 | 14 (21) |
| Receptive sharing | 10 | 1 (10) | 6 | 2 (33) | 16 | 3 (19) |
| p value | | 0.317 | | 0.264 | | 0.869 |
| 2021 | | | | | | |
| No receptive sharing | 44 | 9 (20) | 22 | 3 (14) | 66 | 12 (18) |
| Receptive sharing | 12 | 5 (42) | 6 | 3 (50) | 18 | 8 (44) |
| p value | | 0.133 | | 0.054 | | 0.020 |
| 2022 | | | | | | |
| No receptive sharing | 50 | 5 (10) | 26 | 2 (8) | 77 | 7 (9) |
| Receptive sharing | 11 | 1 (9) | 4 | 1 (25) | 15 | 2 (13) |
| p value | | 1.000 | | 0.360 | | 0.637 |

Table 2.4.6 HCV RNA prevalence by last drug injected, gender and survey year

| Australian Capital Territory | Male | | Female | | Total | |
|------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Last drug injected | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heroin | 29 | 7 (24) | 10 | 4 (40) | 41 | 12 (29) |
| Methamphetamine | 32 | 9 (28) | 11 | 2 (18) | 43 | 11 (26) |
| Other opioids | 4 | 0 (0) | 2 | 1 (50) | 6 | 1 (17) |
| Other drugs | 0 | 0 (0) | 3 | 2 (67) | 3 | 2 (67) |
| p value | | 0.441 | | 0.333 | | 0.419 |
| 2019* | | | | | | |
| Heroin | 30 | 6 (20) | 23 | 7 (30) | 53 | 12 (23) |
| Methamphetamine | 37 | 8 (22) | 12 | 2 (17) | 50 | 10 (20) |
| Other opioids | 5 | 0 (0) | 2 | 0 (0) | 7 | 0 (0) |
| Other drugs | 3 | 1 (33) | 1 | 1 (100) | 4 | 2 (50) |
| p value | | 0.655 | | 0.215 | | 0.266 |
| 2020 | | | | | | |
| Heroin | 31 | 6 (19) | 12 | 2 (17) | 43 | 8 (19) |
| Methamphetamine | 21 | 5 (24) | 12 | 2 (17) | 34 | 7 (21) |
| Other opioids | 2 | 0 (0) | 5 | 1 (20) | 7 | 1 (14) |
| Other drugs | 6 | 2 (33) | 1 | 0 (0) | 7 | 2 (29) |
| p value | | 0.756 | | 0.971 | | 0.913 |
| 2021 | | | | | | |
| Heroin | 28 | 5 (18) | 14 | 2 (14) | 42 | 7 (17) |
| Methamphetamine | 25 | 7 (28) | 11 | 2 (18) | 36 | 9 (25) |
| Other opioids | 4 | 0 (0) | 1 | 1 (100) | 5 | 1 (20) |
| Other drugs | 4 | 2 (50) | 4 | 1 (25) | 8 | 3 (38) |
| p value | | 0.306 | | 0.224 | | 0.569 |
| 2022 | | | | | | |
| Heroin | 24 | 3 (13) | 19 | 3 (16) | 43 | 6 (14) |
| Methamphetamine | 29 | 2 (7) | 10 | 0 (0) | 40 | 2 (5) |
| Other opioids | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| Other drugs | 8 | 1 (13) | 1 | 0 (0) | 9 | 1 (11) |
| p value | | 0.856 | | 0.381 | | 0.541 |

Table 2.4.7 HCV RNA prevalence by frequency of drug injection last month, gender and survey year

| Australian Capital Territory | Male | | Female | | Total | |
|-----------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| Frequency of injection last month | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| Less than daily | 25 | 5 (20) | 10 | 4 (40) | 35 | 9 (26) |
| Daily or more | 35 | 10 (29) | 13 | 5 (38) | 50 | 16 (32) |
| Not last month | 5 | 1 (20) | 3 | 0 (0) | 8 | 1 (13) |
| p value | | 0.701 | | 0.406 | | 0.504 |
| 2019* | | | | | | |
| Less than daily | 29 | 10 (34) | 17 | 3 (18) | 47 | 13 (28) |
| Daily or more | 40 | 5 (13) | 17 | 6 (35) | 57 | 11 (19) |
| Not last month | 5 | 0 (0) | 4 | 1 (25) | 9 | 1 (11) |
| p value | | 0.049 | | 0.524 | | 0.391 |
| 2020 | | | | | | |
| Less than daily | 21 | 2 (10) | 8 | 0 (0) | 30 | 2 (7) |
| Daily or more | 35 | 10 (29) | 20 | 5 (25) | 55 | 15 (27) |
| Not last month | 4 | 1 (25) | 2 | 0 (0) | 6 | 1 (17) |
| p value | | 0.242 | | 0.223 | | 0.073 |
| 2021 | | | | | | |
| Less than daily | 19 | 3 (16) | 12 | 0 (0) | 31 | 3 (10) |
| Daily or more | 37 | 11 (30) | 17 | 6 (35) | 54 | 17 (31) |
| Not last month | 5 | 0 (0) | 2 | 0 (0) | 7 | 0 (0) |
| p value | | 0.223 | | 0.047 | | 0.022 |
| 2022 | | | | | | |
| Less than daily | 26 | 3 (12) | 15 | 1 (7) | 41 | 4 (10) |
| Daily or more | 35 | 3 (9) | 15 | 2 (13) | 51 | 5 (10) |
| Not last month | 3 | 0 (0) | 0 | 0 (0) | 3 | 0 (0) |
| p value | | 1.000 | | 1.000 | | 1.000 |

Table 2.4.8 HCV RNA prevalence by imprisonment last year, gender and survey year

| Australian Capital Territory | Male | | Female | | Total | |
|------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| Imprisonment last year | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| No imprisonment | 50 | 12 (24) | 20 | 7 (35) | 71 | 20 (28) |
| Imprisonment | 6 | 1 (17) | 2 | 1 (50) | 8 | 2 (25) |
| p value | | 0.636 | | 0.706 | | 0.812 |
| 2019* | | | | | | |
| No imprisonment | 56 | 13 (23) | 35 | 7 (20) | 91 | 20 (22) |
| Imprisonment | 17 | 2 (12) | 2 | 2 (100) | 19 | 4 (21) |
| p value | | 0.305 | | 0.011 | | 0.928 |
| 2020 | | | | | | |
| No imprisonment | 49 | 9 (18) | 24 | 2 (8) | 74 | 11 (15) |
| Imprisonment | 8 | 3 (38) | 5 | 2 (40) | 13 | 5 (38) |
| p value | | 0.218 | | 0.062 | | 0.043 |
| 2021 | | | | | | |
| No imprisonment | 48 | 9 (19) | 28 | 6 (21) | 76 | 15 (20) |
| Imprisonment | 8 | 1 (13) | 3 | 0 (0) | 11 | 1 (9) |
| p value | | 0.669 | | 0.372 | | 0.394 |
| 2022 | | | | | | |
| No imprisonment | 56 | 7 (13) | 29 | 2 (7) | 86 | 9 (10) |
| Imprisonment | 7 | 0 (0) | 2 | 1 (50) | 9 | 1 (11) |
| p value | | 1.000 | | 0.187 | | 1.000 |

Table 2.4.9 HCV RNA prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Australian Capital Territory | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| Aboriginal and Torres Strait Islander origin | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| Non Indigenous | 55 | 11 (20) | 17 | 5 (24) | 73 | 16 (22) |
| Indigenous | 12 | 5 (42) | 8 | 5 (63) | 20 | 10 (50) |
| p value | | 0.099 | | 0.156 | | 0.017 |
| 2019* | | | | | | |
| Non Indigenous | 67 | 14 (21) | 29 | 5 (17) | 96 | 18 (19) |
| Indigenous | 9 | 2 (22) | 10 | 5 (50) | 19 | 7 (37) |
| p value | | 0.961 | | 0.044 | | 0.113 |
| 2020 | | | | | | |
| Non Indigenous | 49 | 10 (20) | 20 | 1 (5) | 70 | 11 (16) |
| Indigenous | 9 | 2 (22) | 9 | 4 (44) | 18 | 6 (33) |
| p value | | 0.902 | | 0.009 | | 0.091 |
| 2021 | | | | | | |
| Non Indigenous | 45 | 7 (16) | 19 | 4 (21) | 64 | 11 (17) |
| Indigenous | 15 | 7 (47) | 12 | 2 (17) | 27 | 9 (33) |
| p value | | 0.017 | | 0.763 | | 0.089 |
| 2022 | | | | | | |
| Non Indigenous | 51 | 6 (12) | 17 | 1 (6) | 68 | 7 (10) |
| Indigenous | 13 | 1 (8) | 13 | 1 (8) | 27 | 2 (7) |
| p value | | 1.000 | | 1.000 | | 1.000 |

Table 2.4.10 HCV RNA prevalence by main language spoken at home by parents, gender and survey year

| Australian Capital Territory | Male | | Female | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| Main language spoken at home by parents | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| English speaking | 61 | 16 (26) | 22 | 8 (36) | 85 | 24 (28) |
| Non-English speaking | 4 | 1 (25) | 4 | 1 (25) | 8 | 2 (25) |
| p value | | 0.964 | | 0.678 | | 0.856 |
| 2019* | | | | | | |
| English speaking | 70 | 16 (23) | 35 | 9 (26) | 106 | 24 (23) |
| Non-English speaking | 6 | 0 (0) | 3 | 1 (33) | 9 | 1 (11) |
| p value | | 0.186 | | 0.730 | | 0.368 |
| 2020 | | | | | | |
| English speaking | 55 | 11 (20) | 30 | 5 (17) | 86 | 16 (19) |
| Non-English speaking | 5 | 2 (40) | 0 | 0 (0) | 5 | 2 (40) |
| p value | | 0.299 | | -- | | 0.243 |
| 2021 | | | | | | |
| English speaking | 60 | 13 (22) | 29 | 5 (17) | 89 | 18 (20) |
| Non-English speaking | 2 | 1 (50) | 3 | 1 (33) | 5 | 2 (40) |
| p value | | 0.346 | | 0.497 | | 0.293 |
| 2022 | | | | | | |
| English speaking | 61 | 6 (10) | 30 | 3 (10) | 91 | 9 (10) |
| Non-English speaking | 3 | 1 (33) | 1 | 0 (0) | 4 | 1 (25) |
| p value | | 0.298 | | 1.000 | | 0.364 |

Table 2.4.11 HCV RNA prevalence by region/country of birth, gender and survey year

| Australian Capital Territory | Male | | Female | | Total | |
|------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Australia | 58 | 16 (28) | 24 | 9 (38) | 83 | 25 (30) |
| Other Oceania | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) |
| Other | 6 | 1 (17) | 1 | 0 (0) | 7 | 1 (14) |
| p value | | 0.799 | | 0.551 | | 0.588 |
| 2019* | | | | | | |
| Australia | 66 | 15 (23) | 35 | 9 (26) | 102 | 23 (23) |
| Other Oceania | 3 | 0 (0) | 3 | 1 (33) | 6 | 1 (17) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 2 | 1 (50) | 0 | 0 (0) | 2 | 1 (50) |
| Other | 5 | 0 (0) | 0 | 0 (0) | 5 | 0 (0) |
| p value | | 0.352 | | 0.733 | | 0.465 |
| 2020 | | | | | | |
| Australia | 51 | 11 (22) | 29 | 5 (17) | 81 | 16 (20) |
| Other Oceania | 3 | 0 (0) | 0 | 0 (0) | 3 | 0 (0) |
| Asia | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| UK & Ireland | 1 | 1 (100) | 1 | 0 (0) | 2 | 1 (50) |
| Other | 2 | 1 (50) | 0 | 0 (0) | 2 | 1 (50) |
| p value | | 0.285 | | 0.649 | | 0.581 |
| 2021 | | | | | | |
| Australia | 57 | 14 (25) | 28 | 5 (18) | 85 | 19 (22) |
| Other Oceania | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 2 | 0 (0) | 1 | 0 (0) | 3 | 0 (0) |
| Other | 1 | 0 (0) | 3 | 1 (33) | 4 | 1 (25) |
| p value | | 0.663 | | 0.717 | | 0.695 |
| 2022 | | | | | | |
| Australia | 54 | 5 (9) | 27 | 3 (11) | 82 | 8 (10) |
| Other Oceania | 5 | 1 (20) | 0 | 0 (0) | 5 | 1 (20) |
| Asia | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) |
| UK & Ireland | 2 | 0 (0) | 2 | 0 (0) | 4 | 0 (0) |
| Other | 1 | 1 (100) | 0 | 0 (0) | 1 | 1 (100) |
| p value | | 0.145 | | 1.000 | | 0.162 |

New South Wales

Table 3.1.1 Number (percentage) of respondents by demographic characteristics and survey year

| New South Wales | 2018 | 2019 | 2020* | 2021* | 2022 |
|---|----------|----------|----------|----------|----------|
| Number of sites | 19 | 18 | 12 | 3 | 18 |
| N° surveyed | N=574 | N=553 | N=338 | N=46 | N=420 |
| Response rate (%) | 37% | 41% | 25% | 40% | 23% |
| Gender (%) | | | | | |
| Male | 420 (73) | 397 (72) | 242 (72) | 29 (63) | 275 (65) |
| Female | 147 (26) | 146 (26) | 91 (27) | 16 (35) | 133 (32) |
| Other | 5 (1) | 9 (2) | 5 (1) | 1 (2) | 10 (2) |
| Not reported | 2 (<1) | 1 (<1) | 0 (0) | 0 (0) | 2 (<1) |
| Sexual identity (%) | | | | | |
| Heterosexual | 417 (73) | 397 (72) | 257 (76) | 32 (70) | 327 (78) |
| Bisexual | 65 (11) | 69 (12) | 36 (11) | 5 (11) | 45 (11) |
| Gay/Lesbian | 36 (6) | 38 (7) | 21 (6) | 7 (15) | 31 (7) |
| Other | -- -- | -- -- | 9 (3) | 2 (4) | 15 (4) |
| Not reported | 56 (10) | 49 (9) | 15 (4) | 0 (0) | 2 (<1) |
| Aboriginal and Torres Strait Islander origin (%) | | | | | |
| No | 446 (78) | 424 (77) | 252 (75) | 31 (67) | 315 (75) |
| Yes | 94 (16) | 118 (21) | 67 (20) | 11 (24) | 94 (22) |
| Not reported | 34 (6) | 11 (2) | 19 (6) | 4 (9) | 11 (3) |
| Region/country of birth (%) | | | | | |
| Australia | 470 (82) | 471 (85) | 276 (82) | 36 (78) | 358 (85) |
| Other Oceania | 22 (4) | 22 (4) | 9 (3) | 3 (7) | 14 (3) |
| Asia | 6 (1) | 9 (2) | 12 (4) | 1 (2) | 8 (2) |
| UK & Ireland | 16 (3) | 12 (2) | 9 (3) | 2 (4) | 9 (2) |
| Other | 38 (7) | 29 (5) | 25 (7) | 2 (4) | 26 (6) |
| Not reported | 22 (4) | 10 (2) | 7 (2) | 2 (4) | 5 (1) |
| Main language spoken at home by parents (%) | | | | | |
| English | 512 (89) | 510 (92) | 303 (90) | 43 (93) | 391 (93) |
| Non-English | 43 (7) | 37 (7) | 28 (8) | 2 (4) | 22 (5) |
| Not reported | 19 (3) | 6 (1) | 7 (2) | 1 (2) | 7 (2) |
| Age and time since first injection (years) | | | | | |
| <i>Median age</i> | 44 | 44 | 42.5 | 49 | 45 |
| <i>Age range</i> | 18-69 | 18-73 | 18-67 | 27-66 | 17-71 |
| Age group (%) | | | | | |
| <25 years | 24 (4) | 23 (4) | 10 (3) | 0 (0) | 21 (5) |
| 25+ years | 550 (96) | 530 (96) | 328 (97) | 46 (100) | 397 (95) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 2 (<1) |
| Median age first injection | | | | | |
| <i>Median age first injection</i> | 19 | 19 | 20 | 20 | 20 |
| <i>Age range</i> | 10-54 | 10-63 | 10-60 | 13-45 | 10-55 |
| <i>N° not reported</i> | 40 | 22 | 15 | 1 | 28 |
| Median yrs since first injection | | | | | |
| <i>Median yrs since first injection</i> | 23 | 23 | 22 | 26 | 24 |
| <i>Range</i> | <1-53 | <1-59 | <1-52 | 4-49 | <1-56 |
| Years since first injection | | | | | |
| <3 years | 28 (5) | 40 (7) | 27 (8) | 0 (0) | 37 (9) |
| 3+ years | 505 (88) | 491 (89) | 296 (88) | 45 (98) | 355 (85) |
| Not reported | 41 (7) | 22 (4) | 15 (4) | 1 (2) | 28 (7) |
| Imprisonment last year (%) | | | | | |
| No | 459 (80) | 454 (82) | 265 (78) | 41 (89) | 367 (87) |
| Yes | 68 (12) | 67 (12) | 41 (12) | 4 (9) | 37 (9) |
| Not reported | 47 (8) | 32 (6) | 32 (9) | 1 (2) | 16 (4) |
| N° in prison | | | | | |
| <i>N° in prison</i> | N=68 | N=67 | N=41 | N=4 | N=37 |
| <i>Injected in prison</i> | 26 (38) | 26 (39) | 15 (37) | 1 (25) | 10 (27) |

* Implementation of the 2020 and 2021 ANSPS was impacted by the COVID-19 pandemic, see page 1 for more information

Table 3.1.2 Number (percentage) of respondents by last drug injected, frequency of injecting and survey year

| New South Wales | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|-------------|-------------|-------------|-------------|-------------|
| N° surveyed | N=574 | N=553 | N=338 | N=46 | N=420 |
| Drug last injected (%) | | | | | |
| Cocaine | 11 (2) | 11 (2) | 2 (1) | 1 (2) | 9 (2) |
| Methamphetamine | 203 (35) | 213 (39) | 118 (35) | 25 (54) | 143 (34) |
| Heroin | 175 (30) | 172 (31) | 94 (28) | 16 (35) | 126 (30) |
| Pharm. opioids | 26 (5) | 15 (3) | 8 (2) | 1 (2) | 6 (1) |
| Methadone | 41 (7) | 30 (5) | 32 (9) | 2 (4) | 32 (8) |
| Buprenorphine | 5 (1) | 8 (1) | 2 (1) | 0 (0) | 4 (1) |
| Buprenorphine/naloxone | 3 (1) | 3 (1) | 4 (1) | 0 (0) | 1 (<1) |
| PIEDs | 64 (11) | 56 (10) | 47 (14) | 1 (2) | 69 (16) |
| More than one | 37 (6) | 24 (4) | 22 (7) | 0 (0) | 20 (5) |
| Other | 4 (1) | 13 (2) | 2 (1) | 0 (0) | 3 (1) |
| Not reported | 5 (1) | 8 (1) | 7 (2) | 0 (0) | 7 (2) |
| Ever injected opioids | | | | | |
| No | 135 (24) | 133 (24) | 109 (32) | 12 (26) | 113 (27) |
| Yes | 419 (73) | 418 (76) | 229 (68) | 34 (74) | 305 (73) |
| Not reported | 20 (3) | 2 (<1) | 0 (0) | 0 (0) | 2 (<1) |
| Frequency of injection last month (%) | | | | | |
| Not last month | 94 (16) | 80 (14) | 42 (12) | 11 (24) | 66 (16) |
| Less than weekly | 90 (16) | 85 (15) | 51 (15) | 9 (20) | 72 (17) |
| Weekly not daily | 136 (24) | 123 (22) | 84 (25) | 12 (26) | 97 (23) |
| Daily or more | 232 (40) | 254 (46) | 155 (46) | 14 (30) | 175 (42) |
| Not reported | 22 (4) | 11 (2) | 6 (2) | 0 (0) | 10 (2) |
| Present when someone injected for the first time in the last 12 months (%) | | | | | |
| No | 464 (81) | 432 (78) | 276 (82) | 40 (87) | 340 (81) |
| Yes | 77 (13) | 96 (17) | 47 (14) | 6 (13) | 71 (17) |
| Not reported | 33 (6) | 25 (5) | 15 (4) | 0 (0) | 9 (2) |
| Overdosed in last 12 months | | | | | |
| No | 436 (76) | 432 (78) | 287 (85) | 39 (85) | 363 (86) |
| Yes | 103 (18) | 92 (17) | 35 (10) | 7 (15) | 48 (11) |
| Not reported | 35 (6) | 29 (5) | 16 (5) | 0 (0) | 9 (2) |

Table 3.1.3 Number (percentage) of respondents by injecting behaviour in the last month and survey year

| New South Wales | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|--------------|--------------|--------------|-------------|--------------|
| N° injected last month | N=458 | N=462 | N=290 | N=35 | N=344 |
| Places injected last month (%) | | | | | |
| All private | 298 (65) | 261 (56) | 184 (63) | 25 (71) | 220 (64) |
| Any public | 158 (35) | 200 (43) | 104 (36) | 10 (29) | 122 (35) |
| Not reported | 2 (<1) | 1 (<1) | 2 (1) | 0 (0) | 2 (1) |
| Use of new and sterile needles and syringes last month (%) | | | | | |
| All injections | 363 (79) | 355 (77) | 230 (79) | 29 (83) | 273 (79) |
| Most of the time | 71 (16) | 83 (18) | 44 (15) | 4 (11) | 55 (16) |
| Half of the time | 7 (2) | 8 (2) | 3 (1) | 1 (3) | 5 (1) |
| Some of the time | 6 (1) | 9 (2) | 3 (1) | 1 (3) | 4 (1) |
| Not last month | 6 (1) | 3 (1) | 1 (<1) | 0 (0) | 1 (<1) |
| Not reported | 5 (1) | 4 (1) | 9 (3) | 0 (0) | 6 (2) |
| Re-used someone else's used needle & syringe last month (%) | | | | | |
| None | 384 (84) | 388 (84) | 257 (89) | 31 (89) | 288 (84) |
| Once | 25 (5) | 24 (5) | 11 (4) | 2 (6) | 17 (5) |
| Twice | 13 (3) | 17 (4) | 6 (2) | 1 (3) | 12 (3) |
| 3-5 times | 20 (4) | 16 (3) | 3 (1) | 1 (3) | 14 (4) |
| >5 times | 7 (2) | 10 (2) | 8 (3) | 0 (0) | 9 (3) |
| Not reported | 9 (2) | 7 (2) | 5 (2) | 0 (0) | 4 (1) |
| Number of people needle & syringe was re-used after last month (%) | | | | | |
| None | 384 (84) | 388 (84) | 257 (89) | 31 (89) | 288 (84) |
| One | 31 (7) | 26 (6) | 8 (3) | 2 (6) | 19 (6) |
| Two | 5 (1) | 7 (2) | 3 (1) | 0 (0) | 2 (1) |
| Three to five | 5 (1) | 5 (1) | 5 (2) | 0 (0) | 3 (1) |
| More than five | 3 (1) | 2 (<1) | 3 (1) | 0 (0) | 5 (1) |
| Don't know | 12 (3) | 15 (3) | 6 (2) | 1 (3) | 13 (4) |
| Not reported | 18 (4) | 19 (4) | 8 (3) | 1 (3) | 14 (4) |
| Relationship to people needle & syringe was used after last month (%) [more than one could be selected] | | | | | |
| Regular sex partner | 25 (5) | 19 (4) | 13 (4) | 0 (0) | 19 (6) |
| Casual sex partner | 0 (0) | 2 (<1) | 3 (1) | 0 (0) | 3 (1) |
| Close friend | 15 (3) | 16 (3) | 6 (2) | 0 (0) | 9 (3) |
| Acquaintance | 8 (2) | 12 (3) | 6 (2) | 2 (6) | 4 (1) |
| Other | 8 (2) | 8 (2) | 4 (1) | 1 (3) | 9 (3) |
| Equipment used after someone else last month (%) [more than one could be selected] | | | | | |
| Spoon | 87 (19) | 100 (22) | 55 (19) | 4 (11) | 60 (17) |
| Water | 72 (16) | 78 (17) | 39 (13) | 3 (9) | 50 (15) |
| Filter | 35 (8) | 37 (8) | 21 (7) | 1 (3) | 23 (7) |
| Drug mix | 33 (7) | 51 (11) | 23 (8) | 2 (6) | 36 (10) |
| None | 330 (72) | 317 (69) | 209 (72) | 32 (91) | 245 (71) |
| Injected by someone after that person injected themselves or others last month (%) | | | | | |
| No | 394 (86) | 396 (86) | 238 (82) | 30 (86) | 286 (83) |
| Yes | 56 (12) | 64 (14) | 45 (16) | 5 (14) | 53 (15) |
| Not reported | 8 (2) | 2 (<1) | 7 (2) | 0 (0) | 5 (1) |
| Source of needle acquisition | | | | | |
| Needle Syringe Program | 393 (86) | 398 (86) | 240 (83) | 32 (91) | 296 (86) |
| Chemist/Pharmacy | 83 (18) | 79 (17) | 55 (19) | 3 (9) | 49 (14) |
| Personal sources | 56 (12) | 76 (16) | 35 (12) | 3 (9) | 34 (10) |
| Dispensing/Vending Machine | 124 (27) | 159 (34) | 92 (32) | 12 (34) | 97 (28) |
| Other sources | 26 (6) | 14 (3) | 14 (5) | 2 (6) | 12 (3) |

Table 3.1.4 Number (percentage) of respondents by treatment for drug use and survey year

| New South Wales | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|-------------|-------------|-------------|-------------|-------------|
| N° surveyed | N=574 | N=553 | N=338 | N=46 | N=420 |
| Ever any treatment/therapy for drug use (%) | | | | | |
| No | 153 (27) | 168 (30) | 103 (30) | 11 (24) | 127 (30) |
| Yes | 402 (70) | 381 (69) | 234 (69) | 35 (76) | 293 (70) |
| Not reported | 19 (3) | 4 (1) | 1 (<1) | 0 (0) | 0 (0) |
| History of methadone maintenance treatment (%) | | | | | |
| Currently | 183 (32) | 187 (34) | 90 (27) | 13 (28) | 115 (27) |
| Previously | 132 (23) | 103 (19) | 67 (20) | 6 (13) | 97 (23) |
| Never | 225 (39) | 252 (46) | 158 (47) | 24 (52) | 195 (46) |
| Not reported | 34 (6) | 11 (2) | 23 (7) | 3 (7) | 13 (3) |
| History of buprenorphine (Subutex®) treatment (%) | | | | | |
| Currently | 47 (8) | 33 (6) | 24 (7) | 2 (4) | 29 (7) |
| Previously | 147 (26) | 148 (27) | 74 (22) | 10 (22) | 98 (23) |
| Never | 348 (61) | 363 (66) | 218 (65) | 30 (65) | 283 (67) |
| Not reported | 32 (6) | 9 (2) | 22 (7) | 4 (9) | 10 (2) |
| History of buprenorphine-naloxone (Suboxone®) treatment (%) | | | | | |
| Currently | 35 (6) | 26 (5) | 17 (5) | 0 (0) | 18 (4) |
| Previously | 130 (23) | 126 (23) | 67 (20) | 10 (22) | 94 (22) |
| Never | 374 (65) | 389 (70) | 251 (74) | 33 (72) | 304 (72) |
| Not reported | 35 (6) | 12 (2) | 3 (1) | 3 (7) | 4 (1) |

Table 3.1.5 Number (percentage) of respondents by testing for HIV and HCV, HCV treatment by survey year

| New South Wales | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|-------------|-------------|-------------|-------------|-------------|
| N° surveyed | N=574 | N=553 | N=338 | N=46 | N=420 |
| Previous HIV test (%) | | | | | |
| Yes, ever | 463 (81) | 463 (84) | 255 (75) | 38 (83) | 336 (80) |
| Yes, last year | 307 (53) | 274 (50) | 156 (46) | 22 (48) | 204 (49) |
| >1 year ago | 156 (27) | 189 (34) | 99 (29) | 16 (35) | 132 (31) |
| Never tested | 70 (12) | 63 (11) | 60 (18) | 4 (9) | 68 (16) |
| Not reported | 41 (7) | 27 (5) | 23 (7) | 4 (9) | 16 (4) |
| Previous HCV test (%) | | | | | |
| Yes, ever | 452 (79) | 455 (82) | 271 (80) | 38 (83) | 329 (78) |
| Yes, last year | 315 (55) | 309 (56) | 167 (49) | 29 (63) | 204 (49) |
| >1 year ago | 137 (24) | 146 (26) | 104 (31) | 9 (20) | 125 (30) |
| Never tested | 53 (9) | 54 (10) | 31 (9) | 2 (4) | 49 (12) |
| Not reported | 69 (12) | 44 (8) | 36 (11) | 6 (13) | 42 (10) |
| Lifetime treatment for HCV (%)# | | | | | |
| N° self-reported HCV diagnosis | N=163 | N=147 | N=64 | N=9 | N=78 |
| Antiviral treatment | 97 (60) | 93 (63) | 46 (72) | 5 (56) | 58 (74) |
| No antiviral treatment | 65 (40) | 53 (36) | 18 (28) | 4 (44) | 19 (24) |
| Not reported | 1 (1) | 1 (1) | 0 (0) | 0 (0) | 1 (1) |
| Treatment for HCV in past 12 months (%)** | | | | | |
| N° self-reported HCV diagnosis | N=131 | N=114 | N=47 | N=8 | N=43 |
| Antiviral treatment | 57 (44) | 53 (46) | 22 (47) | 4 (50) | 19 (44) |
| No antiviral treatment | 73 (56) | 60 (53) | 25 (53) | 4 (50) | 23 (53) |
| Not reported | 1 (1) | 1 (1) | 0 (0) | 0 (0) | 1 (2) |

among people who tested HCV antibody positive and did not report spontaneous clearance

* excludes people who reported treatment induced clearance more than 12 months previously

Table 3.1.6 Number (percentage) of respondents by sexual behaviour in the month prior to survey and survey year

| New South Wales | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|-------------|-------------|-------------|-------------|-------------|
| N° surveyed | N=574 | N=553 | N=338 | N=46 | N=420 |
| Sex with a regular partner last month (%) | | | | | |
| No | 279 (49) | 279 (50) | 143 (42) | 30 (65) | 222 (53) |
| Yes | 255 (44) | 245 (44) | 174 (51) | 11 (24) | 182 (43) |
| Not reported | 40 (7) | 29 (5) | 21 (6) | 5 (11) | 16 (4) |
| Condom used with regular partner last month (%) | | | | | |
| N° surveyed | N=255 | N=245 | N=174 | N=11 | N=182 |
| Never | 182 (71) | 171 (70) | 127 (73) | 9 (82) | 134 (74) |
| Sometimes | 26 (10) | 35 (14) | 21 (12) | 2 (18) | 18 (10) |
| Every time | 32 (13) | 26 (11) | 20 (11) | 0 (0) | 21 (12) |
| Not reported | 15 (6) | 13 (5) | 6 (3) | 0 (0) | 9 (5) |
| Sex with other partner(s) last month (%) | | | | | |
| No | 437 (76) | 421 (76) | 252 (75) | 37 (80) | 317 (75) |
| Yes | 82 (14) | 91 (16) | 61 (18) | 5 (11) | 85 (20) |
| Not reported | 55 (10) | 41 (7) | 25 (7) | 4 (9) | 18 (4) |
| Condom used with other partner(s) last month (%) | | | | | |
| N° surveyed | N=82 | N=91 | N=61 | N=5 | N=85 |
| Never | 33 (40) | 35 (38) | 25 (41) | 2 (40) | 43 (51) |
| Sometimes | 21 (26) | 20 (22) | 19 (31) | 2 (40) | 21 (25) |
| Every time | 25 (30) | 28 (31) | 15 (25) | 1 (20) | 20 (24) |
| Not reported | 3 (4) | 8 (9) | 2 (3) | 0 (0) | 1 (1) |
| Sex work last month (%) | | | | | |
| No | 487 (85) | 474 (86) | 293 (87) | 40 (87) | 364 (87) |
| Yes | 30 (5) | 28 (5) | 20 (6) | 1 (2) | 29 (7) |
| Not reported | 57 (10) | 51 (9) | 25 (7) | 5 (11) | 27 (6) |
| Condom used at last sex work (%) | | | | | |
| Yes | 19 (63) | 19 (68) | 10 (50) | 1 (100) | 20 (69) |

Table 3.1.7 Number (percentage) of respondents by COVID-19 related impacts, 2020-2021

| New South Wales | 2020 | 2021 | 2022 |
|---|--------------|-------------|--------------|
| N° surveyed | N=338 | N=46 | N=420 |
| COVID-19 test | | | |
| Yes | 138 (41) | 25 (54) | 384 (91) |
| No | 155 (46) | 8 (17) | 33 (8) |
| Don't know | 9 (3) | 0 (0) | -- -- |
| Not reported | 36 (11) | 13 (28) | 3 (1) |
| Self-reported positive COVID-19 test result | | | |
| Yes | 2 (1) | 0 (0) | 167 (43) |
| Yes, last 12 months | -- -- | -- -- | 132 (34) |
| Yes, more than a year ago | -- -- | -- -- | 35 (9) |
| COVID-19 vaccine uptake | | | |
| ≥1 dose | -- -- | 38 (83) | 351 (84) |
| COVID-19 vaccine doses | | | |
| One dose | -- -- | 8 (17) | 22 (5) |
| Two doses | -- -- | 30 (65) | 136 (32) |
| Three or more doses | -- -- | -- -- | 193 (46) |
| No doses | -- -- | 0 (0) | 59 (14) |
| Not reported | -- -- | 8 (17) | 10 (2) |
| Vaccine type * | | | N=351 |
| Astra Zeneca | -- -- | 12 (26) | 150 (43) |
| Pfizer | -- -- | 21 (46) | 231 (66) |
| Don't know | -- -- | 1 (2) | 25 (7) |
| Other | -- -- | -- -- | 24 (7) |
| Not reported | -- -- | 12 (26) | -- -- |
| * 2021 only one response permitted, 2022 multiple responses permitted | | | |
| Last vaccine location | | | N=420 |
| GP or medical centre | -- -- | -- -- | 130 (31) |
| Pharmacy/ chemist | -- -- | -- -- | 66 (16) |
| Vaccine hub | -- -- | -- -- | 68 (16) |
| Opioid treatment/ methadone clinic | -- -- | -- -- | 18 (4) |
| NSP | -- -- | -- -- | 18 (4) |
| Prison | -- -- | -- -- | 18 (4) |
| Other | -- -- | -- -- | 10 (2) |
| Multiple response | -- -- | -- -- | 29 (7) |
| Not reported | -- -- | -- -- | 63 (15) |
| Vaccine hesitancy - if not had three or more doses | | | |
| Haven't got around to it | -- -- | -- -- | 73 (17) |
| Difficulty booking appointment | -- -- | -- -- | 4 (1) |
| Concerns about side effects | -- -- | -- -- | 29 (7) |
| Concerns about vaccine safety | -- -- | -- -- | 20 (5) |
| Not at risk of serious illness | -- -- | -- -- | 34 (8) |
| Other | -- -- | -- -- | 45 (11) |
| Multiple response | -- -- | -- -- | 23 (5) |
| Not reported | -- -- | -- -- | 192 (46) |

HIV antibody prevalence

Table 3.2.1 HIV antibody prevalence by gender and survey year

| New South Wales Survey year | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | 395 | 15 (3.8) | 144 | 1 (0.7) | 546 | 17 (3.1) |
| 2019 | 387 | 22 (5.7) | 143 | 1 (0.7) | 540 | 24 (4.4) |
| 2020 | 232 | 11 (4.7) | 88 | 0 (0.0) | 325 | 13 (4.0) |
| 2021 | 27 | 2 (7.4) | 16 | 1 (6.3) | 44 | 3 (6.8) |
| 2022 | 270 | 15 (5.6) | 133 | 4 (3.0) | 415 | 22 (5.3) |
| X ² p trend | | 0.372 | | 0.047 | | 0.104 |

Table 3.2.2 HIV antibody prevalence by sexual identity, gender and survey year

| New South Wales Sexual identity | Male | | Female | | Total | |
|------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Heterosexual | 301 | 0 (0.0) | 90 | 1 (1.1) | 392 | 1 (0.3) |
| Bisexual | 26 | 2 (7.7) | 36 | 0 (0.0) | 64 | 2 (3.1) |
| Homosexual | 26 | 10 (38.5) | 7 | 0 (0.0) | 34 | 11 (32.4) |
| p value | | <0.001 | | 1.000 | | <0.001 |
| 2019 | | | | | | |
| Heterosexual | 299 | 1 (0.3) | 88 | 1 (1.1) | 388 | 2 (0.5) |
| Bisexual | 26 | 4 (15.3) | 36 | 0 (0.0) | 67 | 5 (7.5) |
| Homosexual | 31 | 17 (54.8) | 6 | 0 (0.0) | 38 | 17 (44.7) |
| p value | | <0.001 | | 1.000 | | <0.001 |
| 2020 | | | | | | |
| Heterosexual | 194 | 2 (1.0) | 53 | 0 (0.0) | 248 | 2 (0.8) |
| Bisexual | 8 | 0 (0.0) | 24 | 0 (0.0) | 34 | 1 (2.9) |
| Homosexual | 18 | 9 (50.0) | 1 | 0 (0.0) | 20 | 10 (50.0) |
| p value | | <0.001 | | -- | | <0.001 |
| 2021 | | | | | | |
| Heterosexual | 21 | 0 (0.0) | 10 | 1 (10.0) | 31 | 1 (3.2) |
| Bisexual | 1 | 0 (0.0) | 3 | 0 (0.0) | 5 | 0 (0.0) |
| Homosexual | 5 | 2 (40.0) | 1 | 0 (0.0) | 6 | 2 (33.3) |
| p value | | 0.043 | | 1.000 | | 0.204 |
| 2022 | | | | | | |
| Heterosexual | 226 | 1 (0.4) | 96 | 2 (2.1) | 323 | 3 (0.9) |
| Bisexual | 15 | 2 (13.3) | 25 | 1 (4.0) | 45 | 5 (11.1) |
| Homosexual | 23 | 12 (52.2) | 5 | 0 (0.0) | 30 | 12 (40.0) |
| p value | | <0.001 | | 0.174 | | <0.001 |

Table 3.2.3 HIV antibody prevalence by age group, gender and survey year

| New South Wales Age group | Male | | Female | | Total | |
|------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| <25 years | 14 | 1 (7.1) | 7 | 0 (0.0) | 22 | 1 (4.6) |
| 25-34 years | 54 | 5 (9.3) | 24 | 0 (0.0) | 79 | 5 (6.3) |
| 35-44 years | 127 | 3 (2.4) | 55 | 1 (1.8) | 184 | 4 (2.2) |
| 45+ years | 200 | 6 (3.0) | 58 | 0 (0.0) | 261 | 7 (2.7) |
| p value | | 0.094 | | 0.597 | | 0.229 |
| 2019 | | | | | | |
| <25 years | 15 | 0 (0.0) | 6 | 0 (0.0) | 21 | 0 (0.0) |
| 25-34 years | 66 | 3 (4.6) | 25 | 0 (0.0) | 93 | 3 (3.2) |
| 35-44 years | 118 | 3 (2.5) | 51 | 0 (0.0) | 171 | 3 (1.8) |
| 45+ years | 188 | 16 (8.5) | 61 | 1 (1.6) | 255 | 18 (7.1) |
| p value | | 0.150 | | 1.000 | | 0.051 |
| 2020 | | | | | | |
| <25 years | 8 | 0 (0.0) | 1 | 0 (0.0) | 10 | 0 (0.0) |
| 25-34 years | 49 | 1 (2.0) | 16 | 0 (0.0) | 65 | 1 (1.5) |
| 35-44 years | 70 | 1 (1.4) | 37 | 0 (0.0) | 107 | 1 (0.9) |
| 45+ years | 105 | 9 (8.6) | 34 | 0 (0.0) | 143 | 11 (7.7) |
| p value | | 0.154 | | -- | | 0.039 |
| 2021 | | | | | | |
| <25 years | 0 | 0 (0.0) | 0 | 0 (0.0) | 0 | 0 (0.0) |
| 25-34 years | 3 | 0 (0.0) | 3 | 0 (0.0) | 6 | 0 (0.0) |
| 35-44 years | 7 | 2 (28.6) | 3 | 1 (33.3) | 10 | 3 (30.0) |
| 45+ years | 17 | 0 (0.0) | 10 | 0 (0.0) | 28 | 0 (0.0) |
| p value | | 0.128 | | 0.375 | | 0.011 |
| 2022 | | | | | | |
| <25 years | 15 | 0 (0.0) | 5 | 0 (0.0) | 21 | 0 (0.0) |
| 25-34 years | 45 | 1 (2.2) | 17 | 0 (0.0) | 63 | 1 (1.6) |
| 35-44 years | 69 | 3 (4.4) | 44 | 2 (4.6) | 116 | 6 (5.2) |
| 45+ years | 139 | 10 (7.2) | 67 | 2 (3.0) | 213 | 14 (6.6) |
| p value | | 0.570 | | 0.831 | | 0.406 |

Table 3.2.4 HIV antibody prevalence by last drug injected, gender and survey year

| New South Wales Last drug injected | Male | | Female | | Total | |
|---------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Heroin | 107 | 3 (2.8) | 56 | 0 (0.0) | 165 | 3 (1.8) |
| Methamphetamine | 140 | 9 (6.4) | 47 | 1 (2.1) | 191 | 11 (5.8) |
| Other opioids | 47 | 1 (2.1) | 27 | 0 (0.0) | 74 | 1 (1.4) |
| PIEDs | 56 | 0 (0.0) | 3 | 0 (0.0) | 60 | 0 (0.0) |
| Other drugs | 42 | 2 (4.8) | 10 | 0 (0.0) | 52 | 2 (3.9) |
| p value | | 0.247 | | 0.608 | | 0.106 |
| 2019 | | | | | | |
| Heroin | 110 | 1 (0.9) | 51 | 0 (0.0) | 168 | 2 (1.2) |
| Methamphetamine | 153 | 19 (12.4) | 51 | 1 (2.0) | 207 | 20 (9.7) |
| Other opioids | 36 | 2 (5.6) | 19 | 0 (0.0) | 55 | 2 (3.6) |
| PIEDs | 54 | 0 (0.0) | 2 | 0 (0.0) | 56 | 0 (0.0) |
| Other drugs | 29 | 0 (0.0) | 17 | 0 (0.0) | 46 | 0 (0.0) |
| p value | | <0.001 | | 1.000 | | <0.001 |
| 2020 | | | | | | |
| Heroin | 62 | 1 (1.6) | 28 | 0 (0.0) | 91 | 1 (1.1) |
| Methamphetamine | 80 | 9 (11.3) | 30 | 0 (0.0) | 113 | 10 (8.9) |
| Other opioids | 27 | 0 (0.0) | 19 | 0 (0.0) | 46 | 0 (0.0) |
| PIEDs | 43 | 0 (0.0) | 2 | 0 (0.0) | 45 | 0 (0.0) |
| Other drugs | 16 | 0 (0.0) | 7 | 0 (0.0) | 24 | 1 (4.2) |
| p value | | 0.019 | | -- | | 0.012 |
| 2021 | | | | | | |
| Heroin | 9 | 0 (0.0) | 7 | 0 (0.0) | 16 | 0 (0.0) |
| Methamphetamine | 14 | 2 (14.3) | 8 | 1 (12.5) | 23 | 3 (13.0) |
| Other opioids | 2 | 0 (0.0) | 1 | 0 (0.0) | 3 | 0 (0.0) |
| PIEDs | 1 | 0 (0.0) | 0 | 0 (0.0) | 1 | 0 (0.0) |
| Other drugs | 1 | 0 (0.0) | 0 | 0 (0.0) | 1 | 0 (0.0) |
| p value | | 0.641 | | 1.000 | | 0.486 |
| 2022 | | | | | | |
| Heroin | 69 | 0 (0.0) | 53 | 0 (0.0) | 124 | 1 (0.8) |
| Methamphetamine | 84 | 13 (15.5) | 52 | 4 (7.7) | 41 | 17 (12.1) |
| Other opioids | 28 | 0 (0.0) | 13 | 0 (0.0) | 43 | 0 (0.0) |
| PIEDs | 64 | 1 (1.6) | 4 | 0 (0.0) | 68 | 1 (1.5) |
| Other drugs | 22 | 1 (4.6) | 8 | 0 (0.0) | 32 | 2 (6.3) |
| p value | | <0.001 | | 0.213 | | <0.001 |

Table 3.2.5 HIV antibody prevalence by years of injection, gender and survey year

| New South Wales Years of injection | Male | | Female | | Total | |
|---------------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | | | | | | |
| <3 years | 19 | 1 (5.3) | 5 | 0 (0.0) | 24 | 1 (4.2) |
| 3 to 10 years | 60 | 5 (8.3) | 13 | 0 (0.0) | 73 | 5 (6.9) |
| 11+ years | 316 | 9 (2.9) | 126 | 1 (0.8) | 449 | 11 (2.5) |
| p value | | 0.090 | | 1.000 | | 0.098 |
| 2019 | | | | | | |
| <3 years | 31 | 1 (3.2) | 8 | 0 (0.0) | 39 | 1 (2.6) |
| 3 to 10 years | 63 | 7 (11.1) | 11 | 0 (0.0) | 75 | 7 (9.3) |
| 11+ years | 275 | 14 (5.1) | 122 | 1 (0.8) | 405 | 16 (4.0) |
| p value | | 0.144 | | 1.000 | | 0.121 |
| 2020 | | | | | | |
| <3 years | 19 | 1 (5.3) | 8 | 0 (0.0) | 27 | 1 (3.7) |
| 3 to 10 years | 42 | 3 (7.1) | 7 | 0 (0.0) | 50 | 3 (6.0) |
| 11+ years | 161 | 7 (4.4) | 70 | 0 (0.0) | 235 | 9 (3.8) |
| p value | | 0.582 | | -- | | 0.698 |
| 2021 | | | | | | |
| <3 years | 0 | 0 (0.0) | 0 | 0 (0.0) | 0 | 0 (0.0) |
| 3 to 10 years | 3 | 0 (0.0) | 2 | 0 (0.0) | 5 | 0 (0.0) |
| 11+ years | 24 | 2 (8.3) | 13 | 1 (7.7) | 38 | 3 (7.9) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2022 | | | | | | |
| <3 years | 30 | 0 (0.0) | 5 | 0 (0.0) | 36 | 0 (0.0) |
| 3 to 10 years | 36 | 1 (2.8) | 13 | 0 (0.0) | 49 | 1 (2.0) |
| 11+ years | 183 | 12 (6.6) | 109 | 4 (3.7) | 302 | 19 (6.3) |
| p value | | 0.414 | | 1.000 | | 0.222 |

Table 3.2.6 HIV antibody prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| New South Wales Receptively shared syringe last month | Male | | Female | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | | | | | | |
| No receptive sharing | 284 | 12 (4.2) | 87 | 0 (0.0) | 374 | 12 (3.2) |
| Receptive sharing | 48 | 2 (4.2) | 17 | 0 (0.0) | 65 | 2 (3.1) |
| p value | | 1.000 | | -- | | 1.000 |
| 2019 | | | | | | |
| No receptive sharing | 277 | 18 (6.5) | 98 | 0 (0.0) | 381 | 19 (5.0) |
| Receptive sharing | 47 | 3 (6.4) | 17 | 1 (5.9) | 65 | 4 (6.2) |
| p value | | 1.000 | | 0.148 | | 0.760 |
| 2020 | | | | | | |
| No receptive sharing | 174 | 7 (4.0) | 69 | 0 (0.0) | 247 | 9 (3.6) |
| Receptive sharing | 16 | 0 (0.0) | 11 | 0 (0.0) | 27 | 0 (0.0) |
| p value | | 1.000 | | -- | | 0.606 |
| 2021 | | | | | | |
| No receptive sharing | 18 | 1 (5.6) | 11 | 1 (9.1) | 30 | 2 (6.7) |
| Receptive sharing | 2 | 1 (50.0) | 2 | 0 (0.0) | 4 | 1 (25.0) |
| p value | | 0.195 | | 1.000 | | 0.322 |
| 2022 | | | | | | |
| No receptive sharing | 188 | 12 (6.4) | 88 | 1 (1.1) | 285 | 15 (5.3) |
| Receptive sharing | 33 | 2 (6.1) | 17 | 1 (5.9) | 52 | 3 (5.8) |
| p value | | 1.000 | | 0.299 | | 0.747 |

Table 3.2.7 HIV antibody prevalence by frequency of drug injection last month, gender and survey year

| New South Wales | | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Frequency of injection | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | |
| 2018 | | | | | | | |
| Less than daily | 166 | 6 (3.6) | 52 | 0 (0.0) | 220 | 6 (2.7) | |
| Daily or more | 169 | 8 (4.7) | 57 | 0 (0.0) | 227 | 8 (3.5) | |
| Not last month | 56 | 1 (1.8) | 33 | 1 (3.0) | 92 | 3 (3.3) | |
| p value | | 0.725 | | 0.232 | | 0.898 | |
| 2019 | | | | | | | |
| Less than daily | 155 | 13 (8.4) | 49 | 0 (0.0) | 205 | 13 (6.3) | |
| Daily or more | 171 | 8 (4.7) | 69 | 1 (1.5) | 246 | 10 (4.1) | |
| Not last month | 51 | 1 (2.0) | 24 | 0 (0.0) | 78 | 1 (1.3) | |
| p value | | 0.209 | | 1.000 | | 0.195 | |
| 2020 | | | | | | | |
| Less than daily | 95 | 3 (3.2) | 33 | 0 (0.0) | 129 | 3 (2.3) | |
| Daily or more | 98 | 4 (4.1) | 48 | 0 (0.0) | 149 | 6 (4.0) | |
| Not last month | 34 | 4 (11.8) | 7 | 0 (0.0) | 42 | 4 (9.5) | |
| p value | | 0.137 | | -- | | 0.119 | |
| 2021 | | | | | | | |
| Less than daily | 13 | 1 (7.7) | 7 | 0 (0.0) | 20 | 1 (5.0) | |
| Daily or more | 7 | 1 (14.3) | 6 | 1 (16.7) | 0 | 0 (0.0) | |
| Not last month | 7 | 0 (0.0) | 3 | 0 (0.0) | 10 | 0 (0.0) | |
| p value | | 1.000 | | 0.562 | | 0.444 | |
| 2022 | | | | | | | |
| Less than daily | 117 | 8 (6.8) | 47 | 1 (2.1) | 168 | 9 (5.4) | |
| Daily or more | 107 | 6 (5.6) | 59 | 1 (1.7) | 173 | 9 (5.2) | |
| Not last month | 42 | 1 (2.4) | 22 | 2 (9.1) | 64 | 3 (4.7) | |
| p value | | 0.678 | | 0.229 | | 1.000 | |

Table 3.2.8 HIV antibody prevalence by imprisonment last year, gender and survey year

| New South Wales | | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Imprisonment last year | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | |
| 2018 | | | | | | | |
| No imprisonment | 320 | 14 (4.4) | 123 | 1 (0.8) | 448 | 16 (3.6) | |
| Imprisonment | 52 | 1 (1.9) | 15 | 0 (0.0) | 68 | 1 (1.5) | |
| p value | | 0.705 | | 1.000 | | 0.712 | |
| 2019 | | | | | | | |
| No imprisonment | 315 | 21 (6.7) | 123 | 1 (0.8) | 445 | 23 (5.2) | |
| Imprisonment | 48 | 0 (0.0) | 14 | 0 (0.0) | 64 | 0 (0.0) | |
| p value | | 0.091 | | 1.000 | | 0.098 | |
| 2020 | | | | | | | |
| No imprisonment | 180 | 11 (6.1) | 72 | 0 (0.0) | 257 | 13 (5.1) | |
| Imprisonment | 28 | 0 (0.0) | 12 | 0 (0.0) | 40 | 0 (0.0) | |
| p value | | 0.367 | | -- | | 0.228 | |
| 2021 | | | | | | | |
| No imprisonment | 24 | 1 (4.2) | 14 | 1 (7.1) | 39 | 2 (5.1) | |
| Imprisonment | 2 | 0 (0.0) | 2 | 0 (0.0) | 4 | 0 (0.0) | |
| p value | | 1.000 | | 1.000 | | 1.000 | |
| 2022 | | | | | | | |
| No imprisonment | 236 | 15 (6.4) | 116 | 2 (1.7) | 363 | 20 (5.5) | |
| Imprisonment | 24 | 0 (0.0) | 12 | 2 (16.7) | 37 | 2 (5.4) | |
| p value | | 0.374 | | 0.044 | | 1.000 | |

Table 3.2.9 HIV antibody prevalence by condom use at last sex with regular partner, gender and survey year

| New South Wales | | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Condom use at last sex | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | |
| 2018 | | | | | | | |
| No condom use | 128 | 3 (2.3) | 49 | 0 (0.0) | 178 | 3 (1.7) | |
| Condom use | 37 | 5 (13.5) | 16 | 0 (0.0) | 56 | 5 (8.9) | |
| p value | | 0.015 | | -- | | 0.021 | |
| 2019 | | | | | | | |
| No condom use | 120 | 8 (6.7) | 49 | 0 (0.0) | 169 | 8 (4.7) | |
| Condom use | 44 | 1 (2.3) | 13 | 1 (7.7) | 59 | 2 (3.4) | |
| p value | | 0.447 | | 0.210 | | 1.000 | |
| 2020 | | | | | | | |
| No condom use | 83 | 6 (7.2) | 37 | 0 (0.0) | 122 | 7 (5.7) | |
| Condom use | 28 | 1 (3.6) | 13 | 0 (0.0) | 41 | 1 (2.4) | |
| p value | | 0.677 | | -- | | 0.681 | |
| 2021 | | | | | | | |
| No condom use | 4 | 0 (0.0) | 5 | 0 (0.0) | 9 | 0 (0.0) | |
| Condom use | 0 | 0 (0.0) | 1 | 0 (0.0) | 2 | 0 (0.0) | |
| p value | | -- | | -- | | -- | |
| 2022 | | | | | | | |
| No condom use | 79 | 7 (8.9) | 50 | 2 (4.0) | 133 | 10 (7.5) | |
| Condom use | 25 | 0 (0.0) | 12 | 0 (0.0) | 39 | 1 (2.6) | |
| p value | | 0.192 | | 1.000 | | 0.460 | |

Table 3.2.10 HIV antibody prevalence by sex work last month, gender and survey year

| New South Wales | | Male | | Female | | Total | |
|---------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Sex work last month | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | |
| 2018 | | | | | | | |
| No sex work | 354 | 10 (2.8) | 117 | 1 (0.9) | 476 | 12 (2.5) | |
| Sex work | 13 | 2 (15.4) | 15 | 0 (0.0) | 30 | 2 (6.7) | |
| p value | | 0.063 | | 1.000 | | 0.199 | |
| 2019 | | | | | | | |
| No sex work | 338 | 18 (5.3) | 121 | 1 (0.8) | 465 | 20 (4.3) | |
| Sex work | 12 | 2 (16.7) | 12 | 0 (0.0) | 27 | 2 (7.4) | |
| p value | | 0.145 | | 1.000 | | 0.343 | |
| 2020 | | | | | | | |
| No sex work | 210 | 10 (4.8) | 72 | 0 (0.0) | 285 | 11 (3.9) | |
| Sex work | 7 | 1 (14.3) | 10 | 0 (0.0) | 19 | 2 (10.5) | |
| p value | | 0.309 | | -- | | 0.191 | |
| 2021 | | | | | | | |
| No sex work | 23 | 1 (4.4) | 14 | 1 (7.1) | 38 | 2 (5.3) | |
| Sex work | 0 | 0 (0.0) | 1 | 0 (0.0) | 1 | 0 (0.0) | |
| p value | | -- | | 1.000 | | 1.000 | |
| 2022 | | | | | | | |
| No sex work | 238 | 9 (3.8) | 112 | 4 (3.6) | 360 | 16 (4.4) | |
| Sex work | 11 | 5 (45.5) | 16 | 0 (0.0) | 29 | 5 (17.2) | |
| p value | | <0.001 | | 1.000 | | 0.003 | |

Table 3.2.11 HIV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| New South Wales Aboriginal and Torres Strait Islander origin | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Non Indigenous | 321 | 9 (2.8) | 110 | 0 (0.0) | 437 | 9 (2.1) |
| Indigenous | 62 | 5 (8.1) | 30 | 0 (0.0) | 93 | 6 (6.5) |
| p value | | 0.043 | | -- | | 0.020 |
| 2019 | | | | | | |
| Non Indigenous | 315 | 17 (5.4) | 94 | 0 (0.0) | 417 | 17 (4.1) |
| Indigenous | 65 | 5 (7.7) | 47 | 1 (2.1) | 113 | 7 (6.2) |
| p value | | 0.557 | | 0.333 | | 0.337 |
| 2020 | | | | | | |
| Non Indigenous | 184 | 10 (5.4) | 56 | 0 (0.0) | 243 | 11 (4.5) |
| Indigenous | 35 | 1 (2.9) | 29 | 0 (0.0) | 66 | 2 (3.0) |
| p value | | 1.000 | | -- | | 0.742 |
| 2021 | | | | | | |
| Non Indigenous | 19 | 2 (10.5) | 9 | 0 (0.0) | 29 | 2 (6.9) |
| Indigenous | 7 | 0 (0.0) | 4 | 1 (25.0) | 11 | 1 (9.1) |
| p value | | 1.000 | | 0.308 | | 1.000 |
| 2022 | | | | | | |
| Non Indigenous | 213 | 12 (5.6) | 90 | 2 (2.2) | 311 | 15 (4.8) |
| Indigenous | 49 | 3 (6.1) | 40 | 2 (5.0) | 93 | 7 (7.5) |
| p value | | 1.000 | | 0.586 | | 0.313 |

Table 3.2.12 HIV antibody prevalence by main language spoken at home by parents, gender and survey year

| New South Wales Main language spoken at home by parents | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| English speaking | 361 | 15 (4.2) | 135 | 0 (0.0) | 501 | 16 (3.2) |
| Non-English speaking | 33 | 0 (0.0) | 8 | 0 (0.0) | 43 | 0 (0.0) |
| p value | | 0.625 | | -- | | 0.628 |
| 2019 | | | | | | |
| English speaking | 354 | 21 (5.9) | 136 | 0 (0.0) | 498 | 22 (4.4) |
| Non-English speaking | 29 | 1 (3.5) | 7 | 1 (14.3) | 37 | 2 (5.4) |
| p value | | 1.000 | | 0.049 | | 0.678 |
| 2020 | | | | | | |
| English speaking | 206 | 10 (4.9) | 81 | 0 (0.0) | 292 | 12 (4.1) |
| Non-English speaking | 24 | 1 (4.2) | 4 | 0 (0.0) | 28 | 1 (3.6) |
| p value | | 1.000 | | -- | | 1.000 |
| 2021 | | | | | | |
| English speaking | 25 | 2 (8.0) | 15 | 1 (6.7) | 41 | 3 (7.3) |
| Non-English speaking | 2 | 0 (0.0) | 0 | 0 (0.0) | 2 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2022 | | | | | | |
| English speaking | 248 | 14 (5.7) | 127 | 4 (3.2) | 387 | 21 (5.4) |
| Non-English speaking | 16 | 1 (6.3) | 5 | 0 (0.0) | 21 | 1 (4.8) |
| p value | | 1.000 | | 1.000 | | 1.000 |

Table 3.2.13 HIV antibody prevalence by region/country of birth, gender and survey year

| New South Wales | | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Region/country of birth | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | |
| 2018 | | | | | | | |
| Australia | 330 | 12 (3.6) | 125 | 0 (0.0) | 460 | 12 (2.6) | |
| Other Oceania | 14 | 1 (7.1) | 7 | 0 (0.0) | 22 | 2 (9.1) | |
| Asia | 5 | 0 (0.0) | 0 | 0 (0.0) | 5 | 0 (0.0) | |
| UK & Ireland | 13 | 1 (7.7) | 3 | 0 (0.0) | 16 | 1 (6.3) | |
| Other | 31 | 1 (3.2) | 6 | 0 (0.0) | 38 | 1 (2.6) | |
| p value | | 0.478 | | -- | | 0.240 | |
| 2019 | | | | | | | |
| Australia | 329 | 18 (5.5) | 127 | 1 (0.8) | 461 | 20 (4.3) | |
| Other Oceania | 14 | 1 (7.1) | 5 | 0 (0.0) | 22 | 1 (4.6) | |
| Asia | 9 | 3 (33.3) | 0 | 0 (0.0) | 9 | 3 (33.3) | |
| UK & Ireland | 7 | 0 (0.0) | 4 | 0 (0.0) | 11 | 0 (0.0) | |
| Other | 20 | 0 (0.0) | 7 | 0 (0.0) | 28 | 0 (0.0) | |
| p value | | 0.044 | | 1.000 | | 0.024 | |
| 2020 | | | | | | | |
| Australia | 189 | 7 (3.7) | 72 | 0 (0.0) | 265 | 8 (3.0) | |
| Other Oceania | 8 | 1 (12.5) | 1 | 0 (0.0) | 9 | 1 (11.1) | |
| Asia | 10 | 1 (10.0) | 2 | 0 (0.0) | 12 | 1 (8.3) | |
| UK & Ireland | 4 | 0 (0.0) | 4 | 0 (0.0) | 8 | 0 (0.0) | |
| Other | 18 | 1 (5.6) | 7 | 0 (0.0) | 25 | 1 (4.0) | |
| p value | | 0.277 | | -- | | 0.296 | |
| 2021 | | | | | | | |
| Australia | 21 | 1 (4.8) | 14 | 1 (7.1) | 35 | 2 (5.7) | |
| Other Oceania | 1 | 0 (0.0) | 1 | 0 (0.0) | 3 | 0 (0.0) | |
| Asia | 1 | 0 (0.0) | 0 | 0 (0.0) | 1 | 0 (0.0) | |
| UK & Ireland | 1 | 0 (0.0) | 1 | 0 (0.0) | 2 | 0 (0.0) | |
| Other | 1 | 0 (0.0) | 0 | 0 (0.0) | 1 | 0 (0.0) | |
| p value | | 1.000 | | 1.000 | | 1.000 | |
| 2022 | | | | | | | |
| Australia | 224 | 10 (4.5) | 121 | 4 (3.3) | 355 | 16 (4.5) | |
| Other Oceania | 12 | 2 (16.7) | 1 | 0 (0.0) | 14 | 2 (14.3) | |
| Asia | 5 | 0 (0.0) | 2 | 0 (0.0) | 7 | 0 (0.0) | |
| UK & Ireland | 7 | 1 (14.3) | 2 | 0 (0.0) | 9 | 1 (11.1) | |
| Other | 19 | 2 (10.5) | 6 | 0 (0.0) | 25 | 2 (8.0) | |
| p value | | 0.115 | | 1.000 | | 0.196 | |

HCV antibody prevalence

Table 3.3.1 HCV antibody prevalence by gender and survey year

| New South Wales Survey year | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | 395 | 183 (46) | 144 | 60 (42) | 546 | 247 (45) |
| 2019 | 384 | 154 (40) | 141 | 72 (51) | 535 | 231 (43) |
| 2020 | 233 | 79 (34) | 88 | 34 (39) | 326 | 115 (35) |
| 2021 | 28 | 7 (25) | 16 | 6 (38) | 45 | 13 (29) |
| 2022 | 257 | 68 (26) | 125 | 50 (40) | 393 | 121 (31) |
| X ² p trend | | <0.001 | | 0.360 | | <0.001 |

Table 3.3.2 HCV antibody prevalence by sexual identity, gender and survey year

| New South Wales Sexual identity | Male | | Female | | Total | |
|------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heterosexual | 301 | 143 (48) | 90 | 34 (38) | 392 | 178 (45) |
| Bisexual | 26 | 12 (46) | 36 | 17 (47) | 64 | 31 (48) |
| Homosexual | 26 | 8 (31) | 7 | 3 (43) | 34 | 11 (32) |
| p value | | 0.259 | | 0.616 | | 0.282 |
| 2019 | | | | | | |
| Heterosexual | 297 | 126 (42) | 85 | 46 (54) | 383 | 173 (45) |
| Bisexual | 25 | 8 (32) | 37 | 18 (49) | 67 | 29 (43) |
| Homosexual | 31 | 4 (13) | 6 | 2 (33) | 38 | 6 (16) |
| p value | | 0.003 | | 0.575 | | 0.002 |
| 2020 | | | | | | |
| Heterosexual | 194 | 69 (36) | 53 | 18 (34) | 248 | 88 (35) |
| Bisexual | 8 | 3 (38) | 24 | 15 (63) | 34 | 18 (53) |
| Homosexual | 19 | 2 (11) | 1 | 0 (0) | 21 | 2 (10) |
| p value | | 0.107 | | 0.030 | | 0.007 |
| 2021 | | | | | | |
| Heterosexual | 21 | 6 (29) | 10 | 5 (50) | 31 | 11 (35) |
| Bisexual | 1 | 0 (0) | 3 | 1 (33) | 5 | 1 (20) |
| Homosexual | 6 | 1 (17) | 1 | 0 (0) | 7 | 1 (14) |
| p value | | 1.000 | | 1.000 | | 0.787 |
| 2022 | | | | | | |
| Heterosexual | 212 | 61 (29) | 89 | 41 (46) | 302 | 103 (34) |
| Bisexual | 15 | 3 (20) | 24 | 6 (25) | 44 | 9 (20) |
| Homosexual | 24 | 1 (4) | 5 | 0 (0) | 31 | 2 (6) |
| p value | | 0.009 | | 0.041 | | 0.001 |

Table 3.3.3 HCV antibody prevalence by age group, gender and survey year

| New South Wales Age group | Male | | Female | | Total | |
|------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <25 years | 14 | 2 (14) | 7 | 0 (0) | 22 | 2 (9) |
| 25-34 years | 54 | 12 (22) | 24 | 11 (46) | 79 | 24 (30) |
| 35-44 years | 126 | 66 (52) | 55 | 24 (44) | 183 | 91 (50) |
| 45+ years | 201 | 103 (51) | 58 | 25 (43) | 262 | 130 (50) |
| p value | | <0.001 | | 0.151 | | <0.001 |
| 2019 | | | | | | |
| <25 years | 15 | 3 (20) | 7 | 1 (14) | 22 | 4 (18) |
| 25-34 years | 66 | 13 (20) | 25 | 16 (64) | 93 | 29 (31) |
| 35-44 years | 116 | 52 (45) | 51 | 26 (51) | 169 | 80 (47) |
| 45+ years | 187 | 86 (46) | 58 | 29 (50) | 251 | 118 (47) |
| p value | | <0.001 | | 0.144 | | 0.003 |
| 2020 | | | | | | |
| <25 years | 8 | 0 (0) | 1 | 0 (0) | 10 | 0 (0) |
| 25-34 years | 49 | 10 (20) | 16 | 3 (19) | 65 | 13 (20) |
| 35-44 years | 70 | 22 (31) | 37 | 15 (41) | 107 | 37 (35) |
| 45+ years | 106 | 47 (44) | 34 | 16 (47) | 144 | 65 (45) |
| p value | | 0.003 | | 0.202 | | <0.001 |
| 2021 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 3 | 0 (0) | 3 | 0 (0) | 6 | 0 (0) |
| 35-44 years | 8 | 2 (25) | 3 | 1 (33) | 11 | 3 (27) |
| 45+ years | 17 | 5 (29) | 10 | 5 (50) | 28 | 10 (36) |
| p value | | 0.831 | | 0.494 | | 0.253 |
| 2022 | | | | | | |
| <25 years | 15 | 0 (0) | 4 | 1 (25) | 19 | 1 (5) |
| 25-34 years | 45 | 3 (7) | 16 | 6 (38) | 62 | 9 (15) |
| 35-44 years | 66 | 15 (23) | 42 | 12 (29) | 111 | 27 (24) |
| 45+ years | 129 | 48 (37) | 63 | 31 (49) | 199 | 82 (41) |
| p value | | <0.001 | | 0.169 | | <0.001 |

Table 3.3.4 HCV antibody prevalence by years since first injection, gender and survey year

| New South Wales Years of injection | Male | | Female | | Total | |
|---------------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| <3 years | 19 | 1 (5) | 5 | 0 (0) | 24 | 1 (4) |
| 3 to 10 years | 60 | 6 (10) | 13 | 3 (23) | 73 | 9 (12) |
| 11+ years | 288 | 164 (57) | 119 | 55 (46) | 413 | 223 (54) |
| p value | | <0.001 | | 0.040 | | <0.001 |
| 2019 | | | | | | |
| <3 years | 30 | 2 (7) | 9 | 0 (0) | 39 | 2 (5) |
| 3 to 10 years | 63 | 11 (17) | 11 | 3 (27) | 75 | 15 (20) |
| 11+ years | 273 | 134 (49) | 119 | 69 (58) | 400 | 206 (52) |
| p value | | <0.001 | | <0.001 | | <0.001 |
| 2020 | | | | | | |
| <3 years | 19 | 2 (11) | 8 | 0 (0) | 27 | 2 (7) |
| 3 to 10 years | 42 | 7 (17) | 7 | 3 (43) | 50 | 10 (20) |
| 11+ years | 161 | 68 (42) | 70 | 31 (44) | 235 | 101 (43) |
| p value | | <0.001 | | 0.042 | | <0.001 |
| 2021 | | | | | | |
| <3 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 3 to 10 years | 3 | 1 (33) | 2 | 0 (0) | 5 | 1 (20) |
| 11+ years | 25 | 6 (24) | 13 | 6 (46) | 39 | 12 (31) |
| p value | | 1.000 | | 0.486 | | 1.000 |
| 2022 | | | | | | |
| <3 years | 30 | 0 (0) | 4 | 0 (0) | 34 | 0 (0) |
| 3 to 10 years | 36 | 2 (6) | 12 | 2 (17) | 48 | 4 (8) |
| 11+ years | 171 | 62 (36) | 103 | 44 (43) | 284 | 109 (38) |
| p value | | <0.001 | | 0.064 | | <0.001 |

Table 3.3.5 HCV antibody prevalence by re-use of someone else's used needle and syringe last month, years since first injection and survey year

| New South Wales Receptively shared syringe last month | <3 years | | 3+ years | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| No receptive sharing | 16 | 0 (0) | 338 | 160 (47) | 374 | 168 (45) |
| Receptive sharing | 0 | 0 (0) | 59 | 36 (61) | 65 | 40 (62) |
| p value | | -- | | 0.052 | | 0.013 |
| 2019 | | | | | | |
| No receptive sharing | 28 | 1 (4) | 340 | 156 (46) | 378 | 163 (43) |
| Receptive sharing | 4 | 1 (25) | 56 | 34 (61) | 63 | 37 (59) |
| p value | | 0.238 | | 0.040 | | 0.021 |
| 2020 | | | | | | |
| No receptive sharing | 22 | 2 (9) | 220 | 83 (38) | 248 | 87 (35) |
| Receptive sharing | 1 | 0 (0) | 25 | 13 (52) | 27 | 13 (48) |
| p value | | 1.000 | | 0.166 | | 0.180 |
| 2021 | | | | | | |
| No receptive sharing | -- | -- -- | 29 | 9 (31) | 30 | 9 (30) |
| Receptive sharing | -- | -- -- | 4 | 2 (50) | 4 | 2 (50) |
| p value | | -- | | 0.586 | | 0.580 |
| 2022 | | | | | | |
| No receptive sharing | 26 | 0 (0) | 233 | 70 (30) | 268 | 74 (28) |
| Receptive sharing | 2 | 0 (0) | 44 | 29 (66) | 50 | 31 (62) |
| p value | | -- | | <0.001 | | <0.001 |

Table 3.3.6 HCV antibody prevalence by last drug injected, years since first injection and survey year

| New South Wales Last drug injected | <3 years | | 3+ years | | Total | |
|---------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heroin | 2 | 0 (0) | 156 | 94 (60) | 162 | 96 (59) |
| Methamphetamine | 10 | 1 (10) | 167 | 71 (43) | 190 | 77 (41) |
| Other opioids | 1 | 0 (0) | 65 | 40 (62) | 73 | 44 (60) |
| PIEDs | 11 | 0 (0) | 43 | 1 (2) | 60 | 1 (2) |
| Other drugs | 0 | 0 (0) | 46 | 21 (46) | 50 | 23 (46) |
| p value | | 0.691 | | <0.001 | | <0.001 |
| 2019 | | | | | | |
| Heroin | 6 | 0 (0) | 150 | 91 (61) | 163 | 95 (58) |
| Methamphetamine | 15 | 2 (13) | 185 | 75 (41) | 204 | 79 (39) |
| Other opioids | 1 | 0 (0) | 52 | 25 (48) | 53 | 25 (47) |
| PIEDs | 13 | 0 (0) | 37 | 0 (0) | 54 | 0 (0) |
| Other drugs | 1 | 0 (0) | 41 | 27 (66) | 44 | 29 (66) |
| p value | | 0.564 | | <0.001 | | <0.001 |
| 2020 | | | | | | |
| Heroin | 4 | 1 (25) | 85 | 42 (49) | 90 | 43 (48) |
| Methamphetamine | 7 | 1 (14) | 105 | 35 (33) | 114 | 37 (32) |
| Other opioids | 2 | 0 (0) | 42 | 21 (50) | 45 | 22 (49) |
| PIEDs | 12 | 0 (0) | 26 | 0 (0) | 44 | 0 (0) |
| Other drugs | 1 | 0 (0) | 21 | 12 (57) | 23 | 12 (52) |
| p value | | 0.499 | | <0.001 | | <0.001 |
| 2021 | | | | | | |
| Heroin | -- | -- -- | 16 | 5 (31) | 16 | 5 (31) |
| Methamphetamine | -- | -- -- | 23 | 6 (26) | 24 | 6 (26) |
| Other opioids | -- | -- -- | 3 | 2 (67) | 3 | 2 (67) |
| PIEDs | -- | -- -- | 1 | 0 (0) | 1 | 0 (0) |
| Other drugs | -- | -- -- | 1 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | 0.561 | | 0.538 |
| 2022 | | | | | | |
| Heroin | 1 | 0 (0) | 111 | 49 (44) | 115 | 50 (43) |
| Methamphetamine | 4 | 0 (0) | 120 | 33 (28) | 132 | 37 (28) |
| Other opioids | 0 | 0 (0) | 37 | 21 (57) | 37 | 21 (57) |
| PIEDs | 28 | 0 (0) | 32 | 1 (3) | 66 | 2 (3) |
| Other drugs | 0 | 0 (0) | 26 | 8 (31) | 29 | 9 (31) |
| p value | | -- | | <0.001 | | <0.001 |

Table 3.3.7 HCV antibody prevalence by frequency of drug injection last month, years since first injection and survey year

| New South Wales Frequency of injection last month | <3 years | | 3+ years | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| Less than daily | 10 | 0 (0) | 202 | 93 (46) | 221 | 95 (43) |
| Daily or more | 6 | 0 (0) | 203 | 105 (52) | 226 | 115 (51) |
| Not last month | 8 | 1 (13) | 75 | 30 (40) | 92 | 33 (36) |
| p value | | 0.583 | | 0.190 | | 0.037 |
| 2019 | | | | | | |
| Less than daily | 15 | 1 (7) | 181 | 70 (39) | 201 | 75 (37) |
| Daily or more | 17 | 1 (6) | 220 | 123 (56) | 245 | 128 (52) |
| Not last month | 7 | 0 (0) | 66 | 26 (39) | 78 | 26 (33) |
| p value | | 1.000 | | 0.001 | | 0.001 |
| 2020 | | | | | | |
| Less than daily | 11 | 2 (18) | 116 | 38 (33) | 130 | 40 (31) |
| Daily or more | 12 | 0 (0) | 133 | 60 (45) | 149 | 62 (42) |
| Not last month | 4 | 0 (0) | 32 | 12 (38) | 42 | 12 (29) |
| p value | | 0.436 | | 0.135 | | 0.101 |
| 2021 | | | | | | |
| Less than daily | -- | -- -- | 19 | 6 (32) | 20 | 6 (30) |
| Daily or more | -- | -- -- | 14 | 5 (36) | 14 | 5 (36) |
| Not last month | -- | -- -- | 11 | 2 (18) | 11 | 2 (18) |
| p value | | -- | | 0.645 | | 0.644 |
| 2022 | | | | | | |
| Less than daily | 21 | 0 (0) | 130 | 39 (30) | 159 | 41 (26) |
| Daily or more | 7 | 0 (0) | 149 | 60 (40) | 163 | 64 (39) |
| Not last month | 5 | 0 (0) | 49 | 14 (29) | 61 | 15 (25) |
| p value | | -- | | 0.127 | | 0.016 |

Table 3.3.8 HCV antibody prevalence by imprisonment last year, years since first injection and survey year

| New South Wales Imprisonment last year | <3 years | | 3+ years | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| No imprisonment | 18 | 0 (0) | 401 | 185 (46) | 448 | 194 (43) |
| Imprisonment | 6 | 1 (17) | 57 | 32 (56) | 68 | 36 (53) |
| p value | | 0.250 | | 0.157 | | 0.136 |
| 2019 | | | | | | |
| No imprisonment | 36 | 2 (6) | 389 | 175 (45) | 440 | 181 (41) |
| Imprisonment | 2 | 0 (0) | 58 | 29 (50) | 64 | 31 (48) |
| p value | | 1.000 | | 0.475 | | 0.269 |
| 2020 | | | | | | |
| No imprisonment | 24 | 2 (8) | 223 | 80 (36) | 258 | 83 (32) |
| Imprisonment | 3 | 0 (0) | 34 | 17 (50) | 40 | 18 (45) |
| p value | | 1.000 | | 0.113 | | 0.111 |
| 2021 | | | | | | |
| No imprisonment | 0 | 0 (0) | 39 | 12 (31) | 40 | 12 (30) |
| Imprisonment | 0 | 0 (0) | 4 | 0 (0) | 4 | 0 (0) |
| p value | | -- | | 0.563 | | 0.562 |
| 2022 | | | | | | |
| No imprisonment | 32 | 0 (0) | 288 | 93 (32) | 342 | 99 (29) |
| Imprisonment | 1 | 0 (0) | 31 | 14 (45) | 36 | 15 (42) |
| p value | | -- | | 0.149 | | 0.114 |

Table 3.3.9 HCV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| New South Wales Aboriginal and Torres Strait Islander origin | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Non Indigenous | 321 | 144 (45) | 110 | 41 (37) | 437 | 189 (43) |
| Indigenous | 62 | 34 (55) | 30 | 16 (53) | 93 | 50 (54) |
| p value | | 0.149 | | 0.112 | | 0.064 |
| 2019 | | | | | | |
| Non Indigenous | 311 | 116 (37) | 92 | 44 (48) | 411 | 165 (40) |
| Indigenous | 66 | 34 (52) | 47 | 26 (55) | 114 | 60 (53) |
| p value | | 0.032 | | 0.403 | | 0.017 |
| 2020 | | | | | | |
| Non Indigenous | 185 | 60 (32) | 56 | 20 (36) | 244 | 82 (34) |
| Indigenous | 35 | 15 (43) | 29 | 12 (41) | 66 | 27 (41) |
| p value | | 0.233 | | 0.609 | | 0.270 |
| 2021 | | | | | | |
| Non Indigenous | 20 | 5 (25) | 9 | 3 (33) | 30 | 8 (27) |
| Indigenous | 7 | 2 (29) | 4 | 2 (50) | 11 | 4 (36) |
| p value | | 1.000 | | 1.000 | | 0.701 |
| 2022 | | | | | | |
| Non Indigenous | 203 | 51 (25) | 82 | 31 (38) | 292 | 84 (29) |
| Indigenous | 46 | 15 (33) | 40 | 17 (43) | 90 | 33 (37) |
| p value | | 0.355 | | 0.618 | | 0.155 |

Table 3.3.10 HCV antibody prevalence by main language spoken at home by parents, gender and survey year

| New South Wales Main language spoken at home by parents | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| English speaking | 361 | 169 (47) | 135 | 58 (43) | 501 | 230 (46) |
| Non-English speaking | 33 | 14 (42) | 8 | 2 (25) | 43 | 17 (40) |
| p value | | 0.628 | | 0.468 | | 0.421 |
| 2019 | | | | | | |
| English speaking | 352 | 140 (40) | 134 | 69 (51) | 494 | 213 (43) |
| Non-English speaking | 28 | 13 (46) | 7 | 3 (43) | 36 | 17 (47) |
| p value | | 0.489 | | 0.715 | | 0.631 |
| 2020 | | | | | | |
| English speaking | 207 | 69 (33) | 81 | 32 (40) | 293 | 103 (35) |
| Non-English speaking | 24 | 10 (42) | 4 | 2 (50) | 28 | 12 (43) |
| p value | | 0.415 | | 1.000 | | 0.417 |
| 2021 | | | | | | |
| English speaking | 26 | 6 (23) | 15 | 5 (33) | 42 | 11 (26) |
| Non-English speaking | 2 | 1 (50) | 0 | 0 (0) | 2 | 1 (50) |
| p value | | 0.444 | | -- | | 0.476 |
| 2022 | | | | | | |
| English speaking | 236 | 61 (26) | 119 | 49 (41) | 366 | 113 (31) |
| Non-English speaking | 16 | 5 (31) | 5 | 1 (20) | 21 | 6 (29) |
| p value | | 0.634 | | 0.647 | | 0.824 |

Table 3.3.11 HCV antibody prevalence by region/country of birth, gender and survey year

| New South Wales | | Male | | Female | | Total | |
|--------------------------------|------------------|------------------------|------------------|------------------------|------------------|------------------------|--|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | |
| 2018 | | | | | | | |
| Australia | 330 | 154 (47) | 125 | 52 (42) | 460 | 209 (45) | |
| Other Oceania | 14 | 7 (50) | 7 | 3 (43) | 22 | 10 (45) | |
| Asia | 5 | 4 (80) | 0 | 0 (0) | 5 | 4 (80) | |
| UK & Ireland | 13 | 6 (46) | 3 | 2 (67) | 16 | 8 (50) | |
| Other | 31 | 11 (35) | 6 | 2 (33) | 38 | 14 (37) | |
| p value | | 0.445 | | 0.860 | | 0.474 | |
| 2019 | | | | | | | |
| Australia | 327 | 132 (40) | 125 | 64 (51) | 457 | 199 (44) | |
| Other Oceania | 14 | 7 (50) | 5 | 3 (60) | 22 | 11 (50) | |
| Asia | 9 | 3 (33) | 0 | 0 (0) | 9 | 3 (33) | |
| UK & Ireland | 7 | 3 (43) | 4 | 3 (75) | 11 | 6 (55) | |
| Other | 19 | 6 (32) | 7 | 2 (29) | 27 | 8 (30) | |
| p value | | 0.857 | | 0.517 | | 0.493 | |
| 2020 | | | | | | | |
| Australia | 190 | 65 (34) | 72 | 26 (36) | 266 | 93 (35) | |
| Other Oceania | 8 | 1 (13) | 1 | 0 (0) | 9 | 1 (11) | |
| Asia | 10 | 3 (30) | 2 | 1 (50) | 12 | 4 (33) | |
| UK & Ireland | 4 | 0 (0) | 4 | 2 (50) | 8 | 2 (25) | |
| Other | 18 | 10 (56) | 7 | 4 (57) | 25 | 14 (56) | |
| p value | | 0.136 | | 0.709 | | 0.130 | |
| 2021 | | | | | | | |
| Australia | 21 | 5 (24) | 14 | 6 (43) | 35 | 11 (31) | |
| Other Oceania | 1 | 0 (0) | 1 | 0 (0) | 3 | 0 (0) | |
| Asia | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) | |
| UK & Ireland | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) | |
| Other | 2 | 1 (50) | 0 | 0 (0) | 2 | 1 (50) | |
| p value | | 0.764 | | 1.000 | | 0.720 | |
| 2022 | | | | | | | |
| Australia | 210 | 60 (29) | 114 | 46 (40) | 333 | 108 (32) | |
| Other Oceania | 12 | 3 (25) | 1 | 0 (0) | 14 | 4 (29) | |
| Asia | 5 | 1 (20) | 2 | 1 (50) | 7 | 2 (29) | |
| UK & Ireland | 7 | 0 (0) | 2 | 1 (50) | 9 | 1 (11) | |
| Other | 20 | 4 (20) | 5 | 1 (20) | 25 | 5 (20) | |
| p value | | 0.557 | | 0.950 | | 0.544 | |

HCV RNA prevalence

Table 3.4.1 HCV RNA prevalence by gender and survey year

| New South Wales Survey year | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | 270 | 50 (19) | 98 | 20 (20) | 373 | 71 (19) |
| 2019* | 320 | 46 (14) | 118 | 26 (22) | 446 | 72 (16) |
| 2020 | 221 | 20 (9) | 76 | 9 (12) | 300 | 29 (10) |
| 2021 | 26 | 2 (8) | 14 | 2 (14) | 41 | 4 (10) |
| 2022 | 262 | 19 (7) | 131 | 14 (11) | 405 | 35 (9) |
| X ² p trend | | <0.001 | | 0.012 | | <0.001 |

* 2018, 2019: Weighted for gender and HCV antibody status

Totals include respondents where gender was reported as other or not reported

Table 3.4.2 HCV RNA prevalence by sexual identity, gender and survey year

| New South Wales Sexual identity | Male | | Female | | Total | |
|------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heterosexual | 202 | 37 (18) | 59 | 12 (20) | 262 | 49 (19) |
| Bisexual | 17 | 5 (29) | 22 | 4 (18) | 41 | 9 (22) |
| Homosexual | 20 | 3 (15) | 6 | 3 (50) | 27 | 6 (22) |
| p value | | 0.501 | | 0.266 | | 0.842 |
| 2019* | | | | | | |
| Heterosexual | 254 | 36 (14) | 69 | 14 (20) | 323 | 50 (15) |
| Bisexual | 20 | 2 (10) | 32 | 8 (25) | 57 | 10 (18) |
| Homosexual | 26 | 0 (0) | 6 | 0 (0) | 33 | 0 (0) |
| p value | | 0.112 | | 0.416 | | 0.036 |
| 2020 | | | | | | |
| Heterosexual | 184 | 19 (10) | 46 | 6 (13) | 231 | 25 (11) |
| Bisexual | 8 | 0 (0) | 20 | 3 (15) | 29 | 3 (10) |
| Homosexual | 17 | 0 (0) | 1 | 0 (0) | 18 | 0 (0) |
| p value | | 0.242 | | 0.903 | | 0.339 |
| 2021 | | | | | | |
| Heterosexual | 21 | 2 (10) | 9 | 2 (22) | 30 | 4 (13) |
| Bisexual | 1 | 0 (0) | 2 | 0 (0) | 4 | 0 (0) |
| Homosexual | 4 | 0 (0) | 1 | 0 (0) | 5 | 0 (0) |
| p value | | 0.773 | | 0.670 | | 0.512 |
| 2022 | | | | | | |
| Heterosexual | 221 | 19 (9) | 95 | 14 (15) | 317 | 33 (10) |
| Bisexual | 14 | 0 (0) | 25 | 0 (0) | 44 | 0 (0) |
| Homosexual | 21 | 0 (0) | 4 | 0 (0) | 27 | 1 (3) |
| p value | | 0.591 | | 0.347 | | 0.165 |

Table 3.4.3 HCV RNA prevalence by age group, gender and survey year

| New South Wales Age group | Male | | Female | | Total | |
|------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <25 years | 12 | 1 (8) | 5 | 0 (0) | 18 | 1 (6) |
| 25-34 years | 31 | 5 (16) | 13 | 2 (15) | 44 | 7 (16) |
| 35-44 years | 81 | 21 (26) | 35 | 7 (20) | 117 | 30 (26) |
| 45+ years | 146 | 23 (16) | 45 | 10 (22) | 193 | 33 (17) |
| p value | | 0.132 | | 0.588 | | 0.109 |
| 2019* | | | | | | |
| <25 years | 12 | 2 (17) | 6 | 1 (17) | 18 | 3 (17) |
| 25-34 years | 56 | 5 (9) | 22 | 5 (23) | 80 | 10 (13) |
| 35-44 years | 96 | 22 (23) | 41 | 10 (24) | 138 | 32 (23) |
| 45+ years | 156 | 17 (11) | 49 | 10 (20) | 210 | 27 (13) |
| p value | | 0.035 | | 0.899 | | 0.049 |
| 2020 | | | | | | |
| <25 years | 7 | 0 (0) | 1 | 0 (0) | 9 | 0 (0) |
| 25-34 years | 46 | 4 (9) | 16 | 1 (6) | 62 | 5 (8) |
| 35-44 years | 69 | 8 (12) | 30 | 5 (17) | 99 | 13 (13) |
| 45+ years | 99 | 8 (8) | 29 | 3 (10) | 130 | 11 (8) |
| p value | | 0.715 | | 0.719 | | 0.436 |
| 2021 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 3 | 0 (0) | 2 | 0 (0) | 5 | 0 (0) |
| 35-44 years | 6 | 1 (17) | 3 | 0 (0) | 9 | 1 (11) |
| 45+ years | 17 | 1 (6) | 9 | 2 (22) | 27 | 3 (11) |
| p value | | 0.604 | | 0.523 | | 0.735 |
| 2022 | | | | | | |
| <25 years | 14 | 0 (0) | 5 | 0 (0) | 20 | 0 (0) |
| 25-34 years | 45 | 2 (4) | 17 | 2 (12) | 63 | 4 (6) |
| 35-44 years | 67 | 3 (4) | 42 | 3 (7) | 112 | 6 (5) |
| 45+ years | 134 | 14 (10) | 67 | 9 (13) | 208 | 25 (12) |
| p value | | 0.323 | | 0.786 | | 0.103 |

Table 3.4.4 HCV RNA prevalence by years since first injection, gender and survey year

| New South Wales Years of injection | Male | | Female | | Total | |
|---------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <3 years | 15 | 1 (7) | 5 | 0 (0) | 20 | 1 (5) |
| 3 to 10 years | 37 | 2 (5) | 5 | 2 (40) | 42 | 4 (10) |
| 11+ years | 197 | 44 (22) | 83 | 16 (19) | 284 | 60 (21) |
| p value | | 0.030 | | 0.304 | | 0.045 |
| 2019* | | | | | | |
| <3 years | 25 | 2 (8) | 7 | 0 (0) | 32 | 2 (6) |
| 3 to 10 years | 49 | 3 (6) | 11 | 2 (18) | 60 | 6 (10) |
| 11+ years | 232 | 37 (16) | 98 | 23 (23) | 337 | 59 (18) |
| p value | | 0.139 | | 0.331 | | 0.085 |
| 2020 | | | | | | |
| <3 years | 18 | 2 (11) | 7 | 0 (0) | 25 | 2 (8) |
| 3 to 10 years | 40 | 3 (8) | 6 | 1 (17) | 47 | 4 (9) |
| 11+ years | 152 | 14 (9) | 61 | 8 (13) | 215 | 22 (10) |
| p value | | 0.898 | | 0.567 | | 0.893 |
| 2021 | | | | | | |
| <3 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 3 to 10 years | 3 | 1 (33) | 1 | 0 (0) | 4 | 1 (25) |
| 11+ years | 23 | 1 (4) | 12 | 2 (17) | 36 | 3 (8) |
| p value | | 0.076 | | 0.657 | | 0.292 |
| 2022 | | | | | | |
| <3 years | 29 | 0 (0) | 5 | 0 (0) | 35 | 0 (0) |
| 3 to 10 years | 36 | 1 (3) | 13 | 1 (8) | 49 | 2 (4) |
| 11+ years | 177 | 18 (10) | 107 | 13 (12) | 294 | 33 (11) |
| p value | | 0.083 | | 1.000 | | 0.034 |

Table 3.4.5 HCV RNA prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| New South Wales Receptively shared syringe last month | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| No receptive sharing | 193 | 32 (17) | 57 | 10 (18) | 252 | 42 (17) |
| Receptive sharing | 34 | 14 (41) | 13 | 3 (23) | 47 | 17 (36) |
| p value | | 0.002 | | 0.767 | | 0.004 |
| 2019* | | | | | | |
| No receptive sharing | 232 | 32 (14) | 81 | 20 (25) | 318 | 53 (17) |
| Receptive sharing | 36 | 10 (28) | 12 | 4 (33) | 50 | 14 (28) |
| p value | | 0.037 | | 0.643 | | 0.056 |
| 2020 | | | | | | |
| No receptive sharing | 166 | 16 (10) | 60 | 9 (15) | 228 | 25 (11) |
| Receptive sharing | 15 | 4 (27) | 10 | 0 (0) | 25 | 4 (16) |
| p value | | 0.044 | | 0.190 | | 0.453 |
| 2021 | | | | | | |
| No receptive sharing | 17 | 0 (0) | 10 | 2 (20) | 28 | 2 (7) |
| Receptive sharing | 1 | 1 (100) | 2 | 0 (0) | 3 | 1 (33) |
| p value | | <0.001 | | 0.488 | | 0.145 |
| 2022 | | | | | | |
| No receptive sharing | 182 | 12 (7) | 87 | 10 (11) | 278 | 23 (8) |
| Receptive sharing | 33 | 5 (15) | 16 | 4 (25) | 51 | 10 (20) |
| p value | | 0.094 | | 0.225 | | 0.013 |

Table 3.4.6 HCV RNA prevalence by last drug injected, gender and survey year

| New South Wales Last drug injected | Male | | Female | | Total | |
|---------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heroin | 79 | 16 (20) | 38 | 8 (21) | 119 | 24 (20) |
| Methamphetamine | 91 | 22 (24) | 31 | 5 (16) | 125 | 27 (22) |
| Other opioids | 27 | 6 (22) | 18 | 4 (22) | 45 | 10 (22) |
| PIEDs | 34 | 1 (3) | 0 | 0 (0) | 36 | 1 (3) |
| Other drugs | 31 | 4 (13) | 2 | 2 (100) | 40 | 6 (15) |
| p value | | 0.080 | | 0.972 | | 0.095 |
| 2019* | | | | | | |
| Heroin | 95 | 20 (21) | 37 | 9 (24) | 138 | 28 (20) |
| Methamphetamine | 123 | 16 (13) | 44 | 9 (20) | 169 | 25 (15) |
| Other opioids | 30 | 5 (17) | 17 | 3 (18) | 47 | 8 (17) |
| PIEDs | 43 | 0 (0) | 2 | 0 (0) | 45 | 0 (0) |
| Other drugs | 22 | 4 (18) | 13 | 6 (46) | 36 | 10 (28) |
| p value | | 0.024 | | 0.350 | | 0.007 |
| 2020 | | | | | | |
| Heroin | 58 | 8 (14) | 27 | 1 (3) | 86 | 9 (11) |
| Methamphetamine | 76 | 7 (9) | 22 | 4 (18) | 100 | 11 (11) |
| Other opioids | 25 | 3 (12) | 19 | 2 (11) | 44 | 5 (11) |
| PIEDs | 39 | 0 (0) | 2 | 0 (0) | 41 | 0 (0) |
| Other drugs | 15 | 2 (13) | 6 | 2 (33) | 21 | 4 (19) |
| p value | | 0.211 | | 0.238 | | 0.151 |
| 2021 | | | | | | |
| Heroin | 9 | 1 (11) | 5 | 1 (20) | 14 | 2 (14) |
| Methamphetamine | 13 | 1 (8) | 8 | 1 (13) | 22 | 2 (9) |
| Other opioids | 2 | 0 (0) | 1 | 0 (0) | 3 | 0 (0) |
| PIEDs | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| Other drugs | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | 0.975 | | 0.852 | | 0.928 |
| 2022 | | | | | | |
| Heroin | 66 | 9 (14) | 51 | 7 (14) | 119 | 16 (13) |
| Methamphetamine | 80 | 5 (6) | 50 | 3 (6) | 135 | 8 (6) |
| Other opioids | 27 | 3 (11) | 13 | 4 (31) | 42 | 9 (21) |
| PIEDs | 62 | 0 (0) | 3 | 0 (0) | 65 | 0 (0) |
| Other drugs | 22 | 2 (9) | 7 | 0 (0) | 30 | 2 (7) |
| p value | | 0.050 | | 0.095 | | 0.001 |

Table 3.4.7 HCV RNA prevalence by frequency of drug injection last month, gender and survey year

| New South Wales | | Male | | Female | | Total | |
|-----------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|--|
| Frequency of injection last month | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | |
| 2018* | | | | | | | |
| Less than daily | 119 | 20 (17) | 33 | 7 (21) | 154 | 27 (18) | |
| Daily or more | 111 | 27 (24) | 39 | 5 (13) | 151 | 32 (21) | |
| Not last month | 36 | 3 (8) | 25 | 6 (24) | 62 | 9 (15) | |
| p value | | 0.086 | | 0.419 | | 0.508 | |
| 2019* | | | | | | | |
| Less than daily | 133 | 15 (11) | 40 | 10 (25) | 174 | 24 (14) | |
| Daily or more | 136 | 27 (20) | 55 | 14 (25) | 197 | 42 (21) | |
| Not last month | 46 | 3 (7) | 21 | 2 (10) | 69 | 5 (7) | |
| p value | | 0.036 | | 0.263 | | 0.013 | |
| 2020 | | | | | | | |
| Less than daily | 92 | 9 (10) | 29 | 5 (17) | 122 | 14 (11) | |
| Daily or more | 92 | 11 (12) | 42 | 4 (10) | 135 | 15 (11) | |
| Not last month | 32 | 0 (0) | 5 | 0 (0) | 38 | 0 (0) | |
| p value | | 0.129 | | 0.428 | | 0.092 | |
| 2021 | | | | | | | |
| Less than daily | 12 | 0 (0) | 7 | 1 (14) | 19 | 1 (5) | |
| Daily or more | 6 | 1 (17) | 5 | 1 (20) | 12 | 2 (17) | |
| Not last month | 8 | 1 (13) | 2 | 0 (0) | 10 | 1 (10) | |
| p value | | 0.379 | | 0.792 | | 0.581 | |
| 2022 | | | | | | | |
| Less than daily | 114 | 5 (4) | 46 | 5 (11) | 164 | 10 (6) | |
| Daily or more | 104 | 12 (12) | 58 | 9 (16) | 169 | 23 (14) | |
| Not last month | 40 | 2 (5) | 22 | 0 (0) | 62 | 2 (3) | |
| p value | | 0.139 | | 0.137 | | 0.016 | |

Table 3.4.8 HCV RNA prevalence by imprisonment last year, gender and survey year

| New South Wales | | Male | | Female | | Total | |
|------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|--|
| Imprisonment last year | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | |
| 2018* | | | | | | | |
| No imprisonment | 213 | 37 (17) | 84 | 19 (23) | 301 | 56 (19) | |
| Imprisonment | 40 | 9 (23) | 10 | 1 (10) | 51 | 10 (20) | |
| p value | | 0.465 | | 0.300 | | 0.943 | |
| 2019* | | | | | | | |
| No imprisonment | 264 | 33 (13) | 104 | 21 (20) | 374 | 55 (15) | |
| Imprisonment | 37 | 9 (24) | 10 | 4 (40) | 49 | 13 (27) | |
| p value | | 0.061 | | 0.151 | | 0.039 | |
| 2020 | | | | | | | |
| No imprisonment | 172 | 15 (9) | 62 | 9 (15) | 237 | 24 (10) | |
| Imprisonment | 26 | 3 (12) | 11 | 0 (0) | 37 | 3 (8) | |
| p value | | 0.641 | | 0.177 | | 0.702 | |
| 2021 | | | | | | | |
| No imprisonment | 24 | 2 (8) | 12 | 2 (17) | 37 | 4 (11) | |
| Imprisonment | 2 | 0 (0) | 2 | 0 (0) | 4 | 0 (0) | |
| p value | | 0.671 | | 0.533 | | 0.489 | |
| 2022 | | | | | | | |
| No imprisonment | 228 | 15 (7) | 114 | 13 (11) | 353 | 30 (9) | |
| Imprisonment | 24 | 2 (8) | 12 | 1 (8) | 37 | 3 (8) | |
| p value | | 0.669 | | 1.000 | | 1.000 | |

Table 3.4.9 HCV RNA prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| New South Wales | | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|--|
| Aboriginal and Torres Strait Islander origin | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | |
| 2018* | | | | | | | |
| Non Indigenous | 222 | 38 (17) | 72 | 15 (21) | 299 | 53 (18) | |
| Indigenous | 39 | 12 (31) | 23 | 3 (13) | 63 | 15 (24) | |
| p value | | 0.048 | | 0.317 | | 0.318 | |
| 2019* | | | | | | | |
| Non Indigenous | 263 | 31 (12) | 78 | 13 (17) | 348 | 45 (13) | |
| Indigenous | 53 | 13 (25) | 37 | 11 (30) | 91 | 24 (26) | |
| p value | | 0.014 | | 0.090 | | 0.001 | |
| 2020 | | | | | | | |
| Non Indigenous | 174 | 14 (8) | 47 | 6 (13) | 223 | 20 (9) | |
| Indigenous | 34 | 3 (9) | 27 | 3 (11) | 62 | 6 (10) | |
| p value | | 0.880 | | 0.843 | | 0.864 | |
| 2021 | | | | | | | |
| Non Indigenous | 18 | 0 (0) | 7 | 2 (29) | 26 | 2 (8) | |
| Indigenous | 7 | 2 (29) | 4 | 0 (0) | 11 | 2 (18) | |
| p value | | 0.018 | | 0.237 | | 0.348 | |
| 2022 | | | | | | | |
| Non Indigenous | 205 | 15 (7) | 88 | 12 (14) | 301 | 28 (9) | |
| Indigenous | 49 | 3 (6) | 40 | 1 (3) | 93 | 5 (5) | |
| p value | | 1.000 | | 0.062 | | 0.232 | |

Table 3.4.10 HCV RNA prevalence by main language spoken at home by parents, gender and survey year

| New South Wales | | Male | | Female | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|--|
| Main language spoken at home by parents | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | |
| 2018* | | | | | | | |
| English speaking | 252 | 48 (19) | 92 | 19 (21) | 347 | 68 (20) | |
| Non-English speaking | 18 | 2 (11) | 6 | 1 (17) | 26 | 3 (12) | |
| p value | | 0.364 | | 0.695 | | 0.273 | |
| 2019* | | | | | | | |
| English speaking | 298 | 42 (14) | 112 | 25 (22) | 417 | 67 (16) | |
| Non-English speaking | 21 | 4 (19) | 6 | 1 (17) | 28 | 5 (18) | |
| p value | | 0.558 | | 0.750 | | 0.835 | |
| 2020 | | | | | | | |
| English speaking | 196 | 17 (9) | 71 | 8 (11) | 270 | 25 (9) | |
| Non-English speaking | 23 | 3 (13) | 4 | 1 (25) | 27 | 4 (15) | |
| p value | | 0.491 | | 0.411 | | 0.354 | |
| 2021 | | | | | | | |
| English speaking | 24 | 2 (8) | 13 | 1 (8) | 38 | 3 (8) | |
| Non-English speaking | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) | |
| p value | | 0.671 | | -- | | 0.679 | |
| 2022 | | | | | | | |
| English speaking | 240 | 18 (8) | 125 | 14 (11) | 377 | 34 (9) | |
| Non-English speaking | 16 | 1 (6) | 5 | 0 (0) | 21 | 1 (5) | |
| p value | | 1.000 | | 1.000 | | 1.000 | |

Table 3.4.11 HCV RNA prevalence by region/country of birth, gender and survey year

| New South Wales | | | | | | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Region/Country of birth | Male | | Female | | Total | |
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Australia | 225 | 43 (19) | 84 | 18 (21) | 312 | 62 (20) |
| Other Oceania | 9 | 1 (11) | 5 | 1 (20) | 15 | 2 (13) |
| Asia | 4 | 2 (50) | 0 | 0 (0) | 4 | 2 (50) |
| UK & Ireland | 11 | 2 (18) | 2 | 1 (50) | 13 | 3 (23) |
| Other | 20 | 2 (10) | 6 | 0 (0) | 26 | 2 (8) |
| p value | | 0.440 | | 0.430 | | 0.268 |
| 2019* | | | | | | |
| Australia | 277 | 41 (15) | 104 | 25 (24) | 386 | 65 (17) |
| Other Oceania | 11 | 0 (0) | 4 | 1 (25) | 17 | 2 (12) |
| Asia | 6 | 2 (33) | 0 | 0 (0) | 6 | 2 (33) |
| UK & Ireland | 3 | 0 (0) | 4 | 0 (0) | 7 | 0 (0) |
| Other | 17 | 2 (12) | 6 | 0 (0) | 24 | 2 (8) |
| p value | | 0.385 | | 0.380 | | 0.367 |
| 2020 | | | | | | |
| Australia | 181 | 16 (9) | 62 | 7 (11) | 246 | 23 (9) |
| Other Oceania | 8 | 0 (0) | 0 | 0 (0) | 8 | 0 (0) |
| Asia | 10 | 0 (0) | 2 | 0 (0) | 12 | 0 (0) |
| UK & Ireland | 3 | 0 (0) | 4 | 1 (25) | 7 | 1 (14) |
| Other | 16 | 4 (25) | 7 | 1 (14) | 23 | 5 (22) |
| p value | | 0.196 | | 0.890 | | 0.252 |
| 2021 | | | | | | |
| Australia | 20 | 2 (10) | 12 | 2 (17) | 32 | 4 (13) |
| Other Oceania | 1 | 0 (0) | 1 | 0 (0) | 3 | 0 (0) |
| Asia | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| UK & Ireland | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) |
| Other | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| p value | | 0.969 | | 0.823 | | 0.893 |
| 2022 | | | | | | |
| Australia | 216 | 18 (8) | 119 | 14 (12) | 345 | 33 (10) |
| Other Oceania | 11 | 0 (0) | 1 | 0 (0) | 13 | 1 (8) |
| Asia | 5 | 1 (20) | 2 | 0 (0) | 7 | 1 (14) |
| UK & Ireland | 7 | 0 (0) | 2 | 0 (0) | 9 | 0 (0) |
| Other | 20 | 0 (0) | 6 | 0 (0) | 26 | 0 (0) |
| p value | | 0.416 | | 1.000 | | 0.383 |

Northern Territory

Table 4.1.1 Number (percentage) of respondents by demographic characteristics and survey year

| Northern Territory | 2018 | 2019 | 2020* | 2021* | 2022 |
|---|---------|---------|----------|----------|----------|
| Number of sites | 3 | 3 | 3 | 3 | 3 |
| N° surveyed | N=67 | N=85 | N=37 | N=40 | N=20 |
| Response rate (%) | 37% | 42% | 23% | 18% | 15% |
| Gender (%) | | | | | |
| Male | 41 (61) | 53 (62) | 25 (68) | 25 (63) | 14 (70) |
| Female | 26 (39) | 31 (36) | 12 (32) | 15 (38) | 4 (20) |
| Other | 0 (0) | 1 (1) | 0 (0) | 0 (0) | 2 (10) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Sexual identity (%) | | | | | |
| Heterosexual | 60 (90) | 68 (80) | 30 (81) | 29 (73) | 15 (75) |
| Bisexual | 3 (4) | 8 (9) | 4 (11) | 7 (18) | 3 (15) |
| Gay/Lesbian | 4 (6) | 2 (2) | 0 (0) | 2 (5) | 0 (0) |
| Other | -- -- | -- -- | 1 (3) | 2 (5) | 2 (10) |
| Not reported | 0 (0) | 7 (8) | 2 (5) | 0 (0) | 0 (0) |
| Aboriginal and Torres Strait Islander origin (%) | | | | | |
| No | 41 (61) | 55 (65) | 25 (68) | 20 (50) | 15 (75) |
| Yes | 25 (37) | 30 (35) | 10 (27) | 20 (50) | 4 (20) |
| Not reported | 1 (1) | 0 (0) | 2 (5) | 0 (0) | 1 (5) |
| Region/country of birth (%) | | | | | |
| Australia | 62 (93) | 79 (93) | 35 (95) | 39 (98) | 19 (95) |
| Other Oceania | 2 (3) | 1 (1) | 1 (3) | 0 (0) | 0 (0) |
| Asia | 1 (1) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| UK & Ireland | 0 (0) | 1 (1) | 0 (0) | 0 (0) | 1 (5) |
| Other | 1 (1) | 3 (4) | 0 (0) | 0 (0) | 0 (0) |
| Not reported | 1 (1) | 1 (1) | 1 (3) | 1 (3) | 0 (0) |
| Main language spoken at home by parents (%) | | | | | |
| English | 64 (96) | 80 (94) | 33 (89) | 39 (98) | 20 (100) |
| Non-English | 1 (1) | 4 (5) | 2 (5) | 1 (3) | 0 (0) |
| Not reported | 2 (3) | 1 (1) | 2 (5) | 0 (0) | 0 (0) |
| Age and time since first injection (years) | | | | | |
| <i>Median age</i> | 43 | 43.5 | 45 | 50 | 47.5 |
| <i>Age range</i> | 24-61 | 30-70 | 21-64 | 31-65 | 34-68 |
| Age group (%) | | | | | |
| <25 years | 1 (1) | 0 (0) | 1 (3) | 0 (0) | 0 (0) |
| 25+ years | 65 (97) | 84 (99) | 36 (97) | 40 (100) | 20 (100) |
| Not reported | 1 (1) | 1 (1) | 0 (0) | 0 (0) | 0 (0) |
| Median age first injection | | | | | |
| <i>Median age first injection</i> | 17.5 | 18 | 19 | 17 | 18.5 |
| <i>Age range</i> | 10-43 | 11-65 | 10-42 | 10-42 | 10-38 |
| <i>N° not reported</i> | 5 | 3 | 0 | 1 | 2 |
| Median yrs since first injection | | | | | |
| <i>Median yrs since first injection</i> | 23.5 | 22.5 | 24 | 29 | 27 |
| <i>Range</i> | 1-47 | 2-50 | 4-51 | 1-50 | 13-51 |
| Years since first injection | | | | | |
| <3 years | 2 (3) | 1 (1) | 0 (0) | 1 (3) | 0 (0) |
| 3+ years | 60 (90) | 81 (95) | 37 (100) | 38 (95) | 18 (90) |
| Not reported | 5 (7) | 3 (4) | 0 (0) | 1 (3) | 2 (10) |
| Imprisonment last year (%) | | | | | |
| No | 59 (88) | 68 (80) | 32 (86) | 36 (90) | 19 (95) |
| Yes | 4 (6) | 10 (12) | 3 (8) | 4 (10) | 0 (0) |
| Not reported | 4 (6) | 7 (8) | 2 (5) | 0 (0) | 1 (5) |
| N° in prison | | | | | |
| <i>N° in prison</i> | N=4 | N=10 | N=3 | N=4 | N=0 |
| Injected in prison | 1 (25) | 1 (10) | 0 (0) | 0 (0) | 0 (0) |

* Implementation of the 2020 and 2021 ANSPS was impacted by the COVID-19 pandemic, see page 1 for more information

Table 4.1.2 Number (percentage) of respondents by last drug injected, frequency of injecting and survey year

| Northern Territory | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|-------------|-------------|-------------|-------------|-------------|
| N° surveyed | N=67 | N=85 | N=37 | N=40 | N=20 |
| Drug last injected (%) | | | | | |
| Cocaine | 0 (0) | 2 (2) | 0 (0) | 1 (3) | 0 (0) |
| Methamphetamine | 27 (40) | 46 (54) | 25 (68) | 27 (68) | 11 (55) |
| Heroin | 1 (1) | 0 (0) | 0 (0) | 1 (3) | 0 (0) |
| Pharm. opioids | 24 (36) | 23 (27) | 7 (19) | 4 (10) | 6 (30) |
| Methadone | 1 (1) | 6 (7) | 0 (0) | 3 (8) | 0 (0) |
| Buprenorphine | 5 (7) | 4 (5) | 3 (8) | 1 (3) | 1 (5) |
| Buprenorphine/naloxone | 2 (3) | 0 (0) | 1 (3) | 0 (0) | 0 (0) |
| PIEDs | 2 (3) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| More than one | 5 (7) | 2 (2) | 1 (3) | 1 (3) | 2 (10) |
| Other | 0 (0) | 1 (1) | 0 (0) | 1 (3) | 0 (0) |
| Not reported | 0 (0) | 1 (1) | 0 (0) | 1 (3) | 0 (0) |
| Ever injected opioids | | | | | |
| No | 12 (18) | 19 (22) | 10 (27) | 11 (28) | 3 (15) |
| Yes | 53 (79) | 65 (76) | 27 (73) | 29 (73) | 17 (85) |
| Not reported | 2 (3) | 1 (1) | 0 (0) | 0 (0) | 0 (0) |
| Frequency of injection last month (%) | | | | | |
| Not last month | 1 (1) | 6 (7) | 5 (14) | 5 (13) | 1 (5) |
| Less than weekly | 6 (9) | 8 (9) | 8 (22) | 10 (25) | 6 (30) |
| Weekly not daily | 14 (21) | 12 (14) | 7 (19) | 7 (18) | 5 (25) |
| Daily or more | 44 (66) | 59 (69) | 17 (46) | 16 (40) | 8 (40) |
| Not reported | 2 (3) | 0 (0) | 0 (0) | 2 (5) | 0 (0) |
| Present when someone injected for the first time in the last 12 months (%) | | | | | |
| No | 54 (81) | 63 (74) | 33 (89) | 31 (78) | 17 (85) |
| Yes | 8 (12) | 17 (20) | 2 (5) | 9 (23) | 3 (15) |
| Not reported | 5 (7) | 5 (6) | 2 (5) | 0 (0) | 0 (0) |
| Overdosed in last 12 months | | | | | |
| No | 56 (84) | 72 (85) | 33 (89) | 35 (88) | 18 (90) |
| Yes | 6 (9) | 8 (9) | 1 (3) | 4 (10) | 1 (5) |
| Not reported | 5 (7) | 5 (6) | 3 (8) | 1 (3) | 1 (5) |

Table 4.1.3 Number (percentage) of respondents by injecting behaviour in the last month and survey year

| Northern Territory | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|-------------|-------------|-------------|-------------|-------------|
| N° injected last month | N=64 | N=79 | N=32 | N=33 | N=19 |
| Places injected last month (%) | | | | | |
| All private | 40 (63) | 51 (65) | 19 (59) | 15 (45) | 13 (68) |
| Any public | 24 (38) | 28 (35) | 12 (38) | 18 (55) | 6 (32) |
| Not reported | 0 (0) | 0 (0) | 1 (3) | 0 (0) | 0 (0) |
| Use of new and sterile needles and syringes last month (%) | | | | | |
| All injections | 53 (83) | 67 (85) | 26 (81) | 27 (82) | 18 (95) |
| Most of the time | 11 (17) | 8 (10) | 2 (6) | 4 (12) | 1 (5) |
| Half of the time | 0 (0) | 2 (3) | 2 (6) | 1 (3) | 0 (0) |
| Some of the time | 0 (0) | 1 (1) | 1 (3) | 1 (3) | 0 (0) |
| Not last month | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Not reported | 0 (0) | 1 (1) | 1 (3) | 0 (0) | 0 (0) |
| Re-used someone else's used needle & syringe last month (%) | | | | | |
| None | 55 (86) | 70 (89) | 27 (84) | 28 (85) | 19 (100) |
| Once | 6 (9) | 3 (4) | 1 (3) | 0 (0) | 0 (0) |
| Twice | 1 (2) | 0 (0) | 1 (3) | 2 (6) | 0 (0) |
| 3-5 times | 1 (2) | 4 (5) | 1 (3) | 1 (3) | 0 (0) |
| >5 times | 1 (2) | 1 (1) | 1 (3) | 2 (6) | 0 (0) |
| Not reported | 0 (0) | 1 (1) | 1 (3) | 0 (0) | 0 (0) |
| Number of people needle & syringe was re-used after last month (%) | | | | | |
| None | 55 (86) | 70 (89) | 27 (84) | 28 (85) | 19 (100) |
| One | 5 (8) | 2 (3) | 0 (0) | 1 (3) | 0 (0) |
| Two | 0 (0) | 1 (1) | 1 (3) | 2 (6) | 0 (0) |
| Three to five | 1 (2) | 1 (1) | 0 (0) | 0 (0) | 0 (0) |
| More than five | 1 (2) | 0 (0) | 1 (3) | 0 (0) | 0 (0) |
| Don't know | 1 (2) | 2 (3) | 2 (6) | 2 (6) | 0 (0) |
| Not reported | 1 (2) | 3 (4) | 1 (3) | 0 (0) | 0 (0) |
| Relationship to people needle & syringe was used after last month (%) [more than one could be selected] | | | | | |
| Regular sex partner | 6 (9) | 3 (4) | 2 (6) | 0 (0) | 0 (0) |
| Casual sex partner | 1 (2) | 1 (1) | 0 (0) | 0 (0) | 0 (0) |
| Close friend | 2 (3) | 0 (0) | 2 (6) | 4 (12) | 0 (0) |
| Acquaintance | 0 (0) | 3 (4) | 0 (0) | 1 (3) | 0 (0) |
| Other | 1 (2) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Equipment used after someone else last month (%) [more than one could be selected] | | | | | |
| Spoon | 8 (13) | 15 (19) | 3 (9) | 2 (6) | 0 (0) |
| Water | 10 (16) | 10 (13) | 5 (16) | 6 (18) | 1 (5) |
| Filter | 8 (13) | 7 (9) | 3 (9) | 3 (9) | 0 (0) |
| Drug mix | 8 (13) | 4 (5) | 0 (0) | 1 (3) | 0 (0) |
| None | 50 (78) | 55 (70) | 25 (78) | 22 (67) | 18 (95) |
| Injected by someone after that person injected themselves or others last month (%) | | | | | |
| No | 54 (84) | 63 (80) | 26 (81) | 28 (85) | 18 (95) |
| Yes | 10 (16) | 16 (20) | 5 (16) | 5 (15) | 1 (5) |
| Not reported | 0 (0) | 0 (0) | 1 (3) | 0 (0) | 0 (0) |
| Source of needle acquisition | | | | | |
| Needle Syringe Program | 58 (91) | 67 (85) | 28 (88) | 30 (91) | 18 (95) |
| Chemist/Pharmacy | 11 (17) | 9 (11) | 0 (0) | 2 (6) | 2 (11) |
| Personal sources | 5 (8) | 7 (9) | 1 (3) | 4 (12) | 1 (5) |
| Dispensing/Vending Machine | 15 (23) | 17 (22) | 5 (16) | 3 (9) | 3 (16) |
| Other sources | 0 (0) | 1 (1) | 1 (3) | 1 (3) | 0 (0) |

Table 4.1.4 Number (percentage) of respondents by treatment for drug use and survey year

| Northern Territory | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|---------|---------|---------|---------|---------|
| N° surveyed | N=67 | N=85 | N=37 | N=40 | N=20 |
| Ever any treatment/therapy for drug use (%) | | | | | |
| No | 31 (46) | 39 (46) | 16 (43) | 20 (50) | 6 (30) |
| Yes | 35 (52) | 46 (54) | 21 (57) | 20 (50) | 14 (70) |
| Not reported | 1 (1) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| History of methadone maintenance treatment (%) | | | | | |
| Currently | 1 (1) | 10 (12) | 1 (3) | 6 (15) | 1 (5) |
| Previously | 11 (16) | 17 (20) | 7 (19) | 4 (10) | 4 (20) |
| Never | 50 (75) | 57 (67) | 28 (76) | 29 (73) | 14 (70) |
| Not reported | 5 (7) | 1 (1) | 1 (3) | 1 (3) | 1 (5) |
| History of buprenorphine (Subutex®) treatment (%) | | | | | |
| Currently | 6 (9) | 5 (6) | 6 (16) | 3 (8) | 2 (10) |
| Previously | 15 (22) | 17 (20) | 4 (11) | 2 (5) | 6 (30) |
| Never | 44 (66) | 62 (73) | 26 (70) | 35 (88) | 10 (50) |
| Not reported | 2 (3) | 1 (1) | 1 (3) | 0 (0) | 2 (10) |
| History of buprenorphine-naloxone (Suboxone®) treatment (%) | | | | | |
| Currently | 2 (3) | 3 (4) | 4 (11) | 1 (3) | 1 (5) |
| Previously | 16 (24) | 18 (21) | 6 (16) | 4 (10) | 7 (35) |
| Never | 46 (69) | 64 (75) | 27 (73) | 35 (88) | 11 (55) |
| Not reported | 3 (4) | 0 (0) | 0 (0) | 0 (0) | 1 (5) |

Table 4.1.5 Number (percentage) of respondents by testing for HIV and HCV, HCV treatment by survey year

| Northern Territory | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|---------|---------|---------|---------|---------|
| N° surveyed | N=67 | N=85 | N=37 | N=40 | N=20 |
| Previous HIV test (%) | | | | | |
| Yes, ever | 54 (81) | 69 (81) | 26 (70) | 34 (85) | 16 (80) |
| Yes, last year | 34 (51) | 43 (51) | 14 (38) | 18 (45) | 6 (30) |
| >1 year ago | 20 (30) | 26 (31) | 12 (32) | 16 (40) | 10 (50) |
| Never tested | 10 (15) | 11 (13) | 9 (24) | 6 (15) | 2 (10) |
| Not reported | 3 (4) | 5 (6) | 2 (5) | 0 (0) | 2 (10) |
| Previous HCV test (%) | | | | | |
| Yes, ever | 51 (76) | 66 (78) | 27 (73) | 34 (85) | 17 (85) |
| Yes, last year | 34 (51) | 44 (52) | 10 (27) | 20 (50) | 9 (45) |
| >1 year ago | 17 (25) | 22 (26) | 17 (46) | 14 (35) | 8 (40) |
| Never tested | 6 (9) | 8 (9) | 3 (8) | 4 (10) | 1 (5) |
| Not reported | 10 (15) | 11 (13) | 7 (19) | 2 (5) | 2 (10) |
| Lifetime treatment for HCV (%)# | | | | | |
| N° self-reported HCV diagnosis | N=19 | N=23 | N=2 | N=6 | N=4 |
| Antiviral treatment | 3 (16) | 16 (70) | 2 (100) | 3 (50) | 1 (25) |
| No antiviral treatment | 16 (84) | 7 (30) | 0 (0) | 3 (50) | 3 (75) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Treatment for HCV in past 12 months (%)** | | | | | |
| N° self-reported HCV diagnosis | N=19 | N=16 | N=1 | N=4 | N=4 |
| Antiviral treatment | 3 (16) | 8 (50) | 1 (100) | 0 (0) | 1 (25) |
| No antiviral treatment | 16 (84) | 8 (50) | 0 (0) | 4 (100) | 3 (75) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |

among people who tested HCV antibody positive and did not report spontaneous clearance

* excludes people who reported treatment induced clearance more than 12 months previously

Table 4.1.6 Number (percentage) of respondents by sexual behaviour in the month prior to survey and survey year

| Northern Territory | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|-------------|-------------|-------------|-------------|-------------|
| N° surveyed | N=67 | N=85 | N=37 | N=40 | N=20 |
| Sex with a regular partner last month (%) | | | | | |
| No | 29 (43) | 43 (51) | 22 (59) | 25 (63) | 12 (60) |
| Yes | 33 (49) | 36 (42) | 13 (35) | 14 (35) | 7 (35) |
| Not reported | 5 (7) | 6 (7) | 2 (5) | 1 (3) | 1 (5) |
| Condom used with regular partner last month (%) | | | | | |
| N° surveyed | N=33 | N=36 | N=13 | N=14 | N=7 |
| Never | 22 (67) | 28 (78) | 11 (85) | 8 (57) | 6 (86) |
| Sometimes | 4 (12) | 5 (14) | 1 (8) | 1 (7) | 0 (0) |
| Every time | 4 (12) | 3 (8) | 1 (8) | 5 (36) | 1 (14) |
| Not reported | 3 (9) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Sex with other partner(s) last month (%) | | | | | |
| No | 44 (66) | 58 (68) | 31 (84) | 29 (73) | 15 (75) |
| Yes | 17 (25) | 20 (24) | 4 (11) | 10 (25) | 5 (25) |
| Not reported | 6 (9) | 7 (8) | 2 (5) | 1 (3) | 0 (0) |
| Condom used with other partner(s) last month (%) | | | | | |
| N° surveyed | N=17 | N=20 | N=4 | N=10 | N=5 |
| Never | 7 (41) | 7 (35) | 0 (0) | 3 (30) | 2 (40) |
| Sometimes | 1 (6) | 6 (30) | 0 (0) | 0 (0) | 1 (20) |
| Every time | 7 (41) | 6 (30) | 4 (100) | 6 (60) | 1 (20) |
| Not reported | 2 (12) | 1 (5) | 0 (0) | 1 (10) | 1 (20) |
| Sex work last month (%) | | | | | |
| No | 53 (79) | 67 (79) | 31 (84) | 32 (80) | 16 (80) |
| Yes | 6 (9) | 11 (13) | 3 (8) | 8 (20) | 1 (5) |
| Not reported | 8 (12) | 7 (8) | 3 (8) | 0 (0) | 3 (15) |
| Condom used at last sex work (%) | | | | | |
| Yes | 3 (50) | 7 (64) | 3 (100) | 7 (88) | 1 (100) |

Table 4.1.7 Number (percentage) of respondents by COVID-19 related impacts, 2020-2022

| Northern Territory | 2020 | 2021 | 2022 |
|---|---------|---------|---------|
| N° surveyed | N=37 | N=40 | N=20 |
| COVID-19 test | | | |
| Yes | 8 (22) | 9 (23) | 17 (85) |
| No | 25 (68) | 21 (53) | 2 (10) |
| Don't know | 1 (3) | 1 (3) | -- -- |
| Not reported | 3 (8) | 9 (23) | 1 (5) |
| Self-reported positive COVID-19 test result | | | |
| Yes | 0 (0) | 0 (0) | 5 (29) |
| Yes, last 12 months | -- -- | -- -- | 5 (29) |
| Yes, more than a year ago | -- -- | -- -- | 0 (0) |
| COVID-19 vaccine uptake | | | |
| ≥1 dose | -- -- | 23 (58) | 15 (75) |
| COVID-19 vaccine doses | | | |
| One dose | -- -- | 5 (13) | 0 (0) |
| Two doses | -- -- | 18 (45) | 7 (35) |
| Three or more doses | -- -- | -- -- | 8 (40) |
| No doses | -- -- | 13 (33) | 5 (25) |
| Not reported | -- -- | 4 (10) | 0 (0) |
| Vaccine type * | | | N=15 |
| Astra Zeneca | -- -- | 7 (18) | 3 (20) |
| Pfizer | -- -- | 11 (28) | 10 (67) |
| Don't know | -- -- | 1 (3) | 2 (13) |
| Other | -- -- | -- -- | 0 (0) |
| Not reported | -- -- | 21 (53) | -- -- |
| * 2021 only one response permitted, 2022 multiple responses permitted | | | |
| Last vaccine location | | | N=20 |
| GP or medical centre | -- -- | -- -- | 7 (35) |
| Pharmacy/ chemist | -- -- | -- -- | 0 (0) |
| Vaccine hub | -- -- | -- -- | 5 (25) |
| Opioid treatment/ methadone clinic | -- -- | -- -- | 0 (0) |
| NSP | -- -- | -- -- | 0 (0) |
| Prison | -- -- | -- -- | 0 (0) |
| Other | -- -- | -- -- | 1 (5) |
| Multiple response | -- -- | -- -- | 1 (5) |
| Not reported | -- -- | -- -- | 6 (30) |
| Vaccine hesitancy - if not had three or more doses | | | |
| Haven't got around to it | -- -- | -- -- | 3 (15) |
| Difficulty booking appointment | -- -- | -- -- | 0 (0) |
| Concerns about side effects | -- -- | -- -- | 2 (10) |
| Concerns about vaccine safety | -- -- | -- -- | 2 (10) |
| Not at risk of serious illness | -- -- | -- -- | 0 (0) |
| Other | -- -- | -- -- | 4 (20) |
| Multiple response | -- -- | -- -- | 1 (5) |
| Not reported | -- -- | -- -- | 8 (40) |

HIV antibody prevalence

Table 4.2.1 HIV antibody prevalence by gender and survey year

| Northern Territory Survey year | Male | | Female | | Total | |
|-----------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | 41 | 0 (0.0) | 25 | 0 (0.0) | 66 | 0 (0.0) |
| 2019 | 52 | 1 (1.9) | 29 | 1 (3.5) | 82 | 2 (2.4) |
| 2020 | 20 | 0 (0.0) | 10 | 0 (0.0) | 30 | 0 (0.0) |
| 2021 | 25 | 1 (4.0) | 14 | 1 (7.1) | 39 | 2 (5.1) |
| 2022 | 14 | 0 (0.0) | 4 | 0 (0.0) | 20 | 0 (0.0) |
| X ² p trend | | 0.557 | | 0.409 | | 0.369 |

HCV antibody prevalence

Table 4.3.1 HCV antibody prevalence by gender and survey year

| Northern Territory Survey year | Male | | Female | | Total | |
|-----------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | 41 | 15 (37) | 25 | 11 (44) | 66 | 26 (39) |
| 2019 | 51 | 23 (45) | 29 | 7 (24) | 81 | 30 (37) |
| 2020 | 20 | 5 (25) | 10 | 0 (0) | 30 | 5 (17) |
| 2021 | 24 | 7 (29) | 14 | 2 (14) | 38 | 9 (24) |
| 2022 | 14 | 5 (36) | 4 | 0 (0) | 20 | 5 (25) |
| X ² p trend | | 0.410 | | 0.005 | | 0.030 |

Table 4.3.2 HCV antibody prevalence by sexual identity, gender and survey year

| Northern Territory Sexual identity | Male | | Female | | Total | |
|---------------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| Heterosexual | 40 | 15 (38) | 20 | 9 (45) | 60 | 24 (40) |
| Bisexual | 0 | 0 (0) | 2 | 0 (0) | 2 | 0 (0) |
| Homosexual | 1 | 0 (0) | 3 | 2 (67) | 4 | 2 (50) |
| p value | | 1.000 | | 0.480 | | 0.656 |
| 2019 | | | | | | |
| Heterosexual | 43 | 20 (47) | 21 | 4 (19) | 65 | 24 (37) |
| Bisexual | 4 | 2 (50) | 3 | 1 (33) | 7 | 3 (43) |
| Homosexual | 1 | 0 (0) | 1 | 1 (100) | 2 | 1 (50) |
| p value | | 1.000 | | 0.234 | | 1.000 |
| 2020 | | | | | | |
| Heterosexual | 19 | 5 (26) | 5 | 0 (0) | 24 | 5 (21) |
| Bisexual | 1 | 0 (0) | 2 | 0 (0) | 3 | 0 (0) |
| Homosexual | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2021 | | | | | | |
| Heterosexual | 18 | 6 (33) | 9 | 1 (11) | 27 | 7 (26) |
| Bisexual | 4 | 1 (25) | 3 | 0 (0) | 7 | 1 (14) |
| Homosexual | 1 | 0 (0) | 1 | 1 (100) | 2 | 1 (50) |
| p value | | 1.000 | | 0.308 | | 0.790 |
| 2022 | | | | | | |
| Heterosexual | 12 | 4 (33) | 3 | 0 (0) | 15 | 4 (27) |
| Bisexual | 2 | 1 (50) | 1 | 0 (0) | 3 | 1 (33) |
| Homosexual | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | 1.000 | | -- | | 1.000 |

Table 4.3.3 HCV antibody prevalence by age group, gender and survey year

| Northern Territory Age group | Male | | Female | | Total | |
|---------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <25 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 25-34 years | 5 | 1 (20) | 4 | 1 (25) | 9 | 2 (22) |
| 35-44 years | 18 | 5 (28) | 7 | 4 (57) | 25 | 9 (36) |
| 45+ years | 16 | 8 (50) | 14 | 6 (43) | 30 | 14 (47) |
| p value | | 0.417 | | 0.654 | | 0.501 |
| 2019 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 9 | 3 (33) | 5 | 0 (0) | 14 | 3 (21) |
| 35-44 years | 14 | 7 (50) | 15 | 5 (33) | 29 | 12 (41) |
| 45+ years | 27 | 13 (48) | 9 | 2 (22) | 37 | 15 (41) |
| p value | | 0.799 | | 0.376 | | 0.459 |
| 2020 | | | | | | |
| <25 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 25-34 years | 3 | 0 (0) | 0 | 0 (0) | 3 | 0 (0) |
| 35-44 years | 4 | 1 (25) | 3 | 0 (0) | 7 | 1 (14) |
| 45+ years | 12 | 4 (33) | 7 | 0 (0) | 19 | 4 (21) |
| p value | | 0.830 | | -- | | 1.000 |
| 2021 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| 35-44 years | 7 | 2 (29) | 4 | 1 (4) | 11 | 3 (27) |
| 45+ years | 15 | 5 (33) | 10 | 1 (10) | 25 | 6 (24) |
| p value | | 1.000 | | 0.505 | | 1.000 |
| 2022 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 35-44 years | 4 | 1 (25) | 1 | 0 (0) | 5 | 1 (20) |
| 45+ years | 9 | 4 (44) | 3 | 0 (0) | 14 | 4 (29) |
| p value | | 1.000 | | -- | | 1.000 |

Table 4.3.4 HCV antibody prevalence by years since first injection, gender and survey year

| Northern Territory Years of injection | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <3 years | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| 3 to 10 years | 6 | 3 (50) | 2 | 1 (50) | 8 | 4 (50) |
| 11+ years | 31 | 10 (32) | 21 | 8 (38) | 52 | 18 (35) |
| p value | | 0.555 | | 1.000 | | 0.419 |
| 2019 | | | | | | |
| <3 years | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| 3 to 10 years | 5 | 2 (40) | 5 | 0 (0) | 10 | 2 (20) |
| 11+ years | 43 | 20 (47) | 23 | 7 (30) | 67 | 27 (40) |
| p value | | 1.000 | | 0.461 | | 0.370 |
| 2020 | | | | | | |
| <3 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 3 to 10 years | 3 | 0 (0) | 0 | 0 (0) | 3 | 0 (0) |
| 11+ years | 17 | 5 (29) | 10 | 0 (0) | 27 | 5 (19) |
| p value | | 0.539 | | -- | | 1.000 |
| 2021 | | | | | | |
| <3 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 3 to 10 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 11+ years | 22 | 7 (32) | 14 | 2 (14) | 36 | 9 (25) |
| p value | | 1.000 | | -- | | 1.000 |
| 2022 | | | | | | |
| <3 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 3 to 10 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 11+ years | 12 | 4 (33) | 4 | 0 (0) | 18 | 4 (22) |
| p value | | -- | | -- | | -- |

Table 4.3.5 HCV antibody prevalence by re-use of someone else's used needle and syringe last month, years since first injection and survey year

| Northern Territory Receptively shared syringe last month | <3 years | | 3+ years | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| No receptive sharing | 2 | 2 (100) | 49 | 20 (41) | 55 | 24 (44) |
| Receptive sharing | 0 | 0 (0) | 9 | 2 (22) | 9 | 2 (22) |
| p value | | -- | | 0.459 | | 0.291 |
| 2019 | | | | | | |
| No receptive sharing | 1 | 0 (0) | 62 | 22 (35) | 66 | 23 (35) |
| Receptive sharing | 0 | 0 (0) | 8 | 4 (50) | 8 | 4 (50) |
| p value | | -- | | 0.458 | | 0.452 |
| 2020 | | | | | | |
| No receptive sharing | 0 | 0 (0) | 22 | 3 (14) | 22 | 3 (14) |
| Receptive sharing | 0 | 0 (0) | 3 | 0 (0) | 3 | 0 (0) |
| p value | | -- | | 1.000 | | 1.000 |
| 2021 | | | | | | |
| No receptive sharing | 0 | 0 (0) | 26 | 5 (19) | 26 | 5 (19) |
| Receptive sharing | 0 | 0 (0) | 5 | 2 (40) | 5 | 2 (40) |
| p value | | -- | | 0.562 | | 0.562 |
| 2022 | | | | | | |
| No receptive sharing | 0 | 0 (0) | 18 | 4 (22) | 19 | 5 (26) |
| Receptive sharing | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |

Table 4.3.6 HCV antibody prevalence by last drug injected, years since first injection and survey year

| Northern Territory Last drug injected | <3 years | | 3+ years | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Methamphetamine | 1 | 1 (100) | 24 | 11 (46) | 26 | 12 (46) |
| Pharm. Opioids | 0 | 0 (0) | 21 | 6 (29) | 23 | 8 (35) |
| OAT* | 0 | 0 (0) | 7 | 4 (57) | 8 | 5 (63) |
| Other drugs | 1 | 1 (100) | 7 | 1 (14) | 8 | 1 (13) |
| p value | | -- | | 0.241 | | 0.181 |
| 2019 | | | | | | |
| Methamphetamine | 1 | 0 (0) | 39 | 12 (31) | 42 | 13 (31) |
| Pharm. Opioids | 0 | 0 (0) | 23 | 11 (48) | 23 | 11 (48) |
| OAT* | 0 | 0 (0) | 10 | 3 (30) | 10 | 3 (30) |
| Other drugs | 0 | 0 (0) | 4 | 2 (50) | 5 | 2 (40) |
| p value | | -- | | 0.511 | | 0.563 |
| 2020 | | | | | | |
| Methamphetamine | 0 | 0 (0) | 19 | 2 (11) | 19 | 2 (11) |
| Pharm. Opioids | 0 | 0 (0) | 6 | 2 (33) | 6 | 2 (33) |
| OAT* | 0 | 0 (0) | 4 | 0 (0) | 4 | 0 (0) |
| Other drugs | 0 | 0 (0) | 1 | 1 (100) | 1 | 1 (100) |
| p value | | -- | | 0.057 | | 0.057 |
| 2021 | | | | | | |
| Methamphetamine | 1 | 0 (0) | 23 | 3 (13) | 24 | 3 (13) |
| Pharm. Opioids | 0 | 0 (0) | 4 | 1 (25) | 4 | 1 (25) |
| OAT* | 0 | 0 (0) | 4 | 2 (50) | 4 | 2 (50) |
| Other drugs | 0 | 0 (0) | 4 | 2 (50) | 4 | 2 (50) |
| p value | | -- | | 0.203 | | 0.179 |
| 2022 | | | | | | |
| Methamphetamine | 0 | 0 (0) | 9 | 2 (22) | 11 | 3 (27) |
| Pharm. Opioids | 0 | 0 (0) | 6 | 1 (17) | 6 | 1 (17) |
| OAT* | 0 | 0 (0) | 1 | 1 (100) | 1 | 1 (100) |
| Other drugs | 0 | 0 (0) | 2 | 0 (0) | 2 | 0 (0) |
| p value | | -- | | 0.243 | | 0.270 |

*OAT = methadone, buprenorphine or buprenorphine/naloxone

Table 4.3.7 HCV antibody prevalence by frequency of drug injection last month, years since first injection and survey year

| Northern Territory Frequency of injection last month | <3 years | | 3+ years | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| Less than daily | 1 | 1 (100) | 18 | 9 (50) | 20 | 10 (50) |
| Daily or more | 1 | 1 (100) | 40 | 13 (33) | 44 | 16 (36) |
| Not last month | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | 0.363 | | 0.481 |
| 2019 | | | | | | |
| Less than daily | 0 | 0 (0) | 19 | 9 (47) | 20 | 9 (45) |
| Daily or more | 1 | 0 (0) | 52 | 17 (33) | 55 | 18 (33) |
| Not last month | 0 | 0 (0) | 6 | 3 (50) | 6 | 3 (50) |
| p value | | -- | | 0.378 | | 0.483 |
| 2020 | | | | | | |
| Less than daily | 0 | 0 (0) | 12 | 2 (17) | 12 | 2 (17) |
| Daily or more | 0 | 0 (0) | 14 | 1 (7) | 14 | 1 (7) |
| Not last month | 0 | 0 (0) | 4 | 2 (50) | 4 | 2 (50) |
| p value | | -- | | 0.134 | | 0.134 |
| 2021 | | | | | | |
| Less than daily | 0 | 0 (0) | 17 | 2 (12) | 17 | 2 (12) |
| Daily or more | 0 | 0 (0) | 14 | 5 (36) | 14 | 5 (36) |
| Not last month | 1 | 0 (0) | 4 | 1 (25) | 5 | 1 (20) |
| p value | | -- | | 0.220 | | 0.345 |
| 2022 | | | | | | |
| Less than daily | 0 | 0 (0) | 10 | 2 (20) | 11 | 3 (27) |
| Daily or more | 0 | 0 (0) | 8 | 2 (25) | 8 | 2 (25) |
| Not last month | 0 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | 1.000 | | 1.000 |

Table 4.3.8 HCV antibody prevalence by imprisonment last year, years since first injection and survey year

| Northern Territory Imprisonment last year | <3 years | | 3+ years | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| No imprisonment | 2 | 2 (100) | 54 | 20 (37) | 59 | 23 (39) |
| Imprisonment | 0 | 0 (0) | 3 | 2 (67) | 4 | 3 (75) |
| p value | | -- | | 0.553 | | 0.297 |
| 2019 | | | | | | |
| No imprisonment | 1 | 0 (0) | 65 | 27 (42) | 66 | 27 (41) |
| Imprisonment | 0 | 0 (0) | 7 | 2 (29) | 8 | 2 (25) |
| p value | | -- | | 0.694 | | 0.469 |
| 2020 | | | | | | |
| No imprisonment | 0 | 0 (0) | 25 | 5 (20) | 25 | 5 (20) |
| Imprisonment | 0 | 0 (0) | 3 | 0 (0) | 3 | 0 (0) |
| p value | | -- | | 1.000 | | 1.000 |
| 2021 | | | | | | |
| No imprisonment | 1 | 0 (0) | 32 | 9 (28) | 34 | 9 (26) |
| Imprisonment | 0 | 0 (0) | 4 | 0 (0) | 4 | 0 (0) |
| p value | | -- | | 0.553 | | 0.554 |
| 2022 | | | | | | |
| No imprisonment | 0 | 0 (0) | 17 | 4 (24) | 19 | 5 (26) |
| Imprisonment | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |

Table 4.3.9 HCV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Northern Territory Aboriginal and Torres Strait Islander origin | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Non Indigenous | 27 | 9 (33) | 14 | 7 (50) | 41 | 16 (39) |
| Indigenous | 14 | 6 (43) | 11 | 4 (36) | 25 | 10 (40) |
| p value | | 0.548 | | 0.689 | | 0.937 |
| 2019 | | | | | | |
| Non Indigenous | 39 | 16 (41) | 13 | 1 (8) | 53 | 17 (32) |
| Indigenous | 12 | 7 (58) | 16 | 6 (38) | 28 | 13 (46) |
| p value | | 0.292 | | 0.093 | | 0.203 |
| 2020 | | | | | | |
| Non Indigenous | 12 | 3 (25) | 9 | 0 (0) | 21 | 3 (14) |
| Indigenous | 6 | 1 (17) | 1 | 0 (0) | 7 | 1 (14) |
| p value | | 1.000 | | -- | | 1.000 |
| 2021 | | | | | | |
| Non Indigenous | 13 | 4 (31) | 6 | 0 (0) | 19 | 4 (21) |
| Indigenous | 11 | 3 (27) | 8 | 2 (25) | 19 | 5 (26) |
| p value | | 0.605 | | 0.473 | | 1.000 |
| 2022 | | | | | | |
| Non Indigenous | 10 | 4 (40) | 4 | 0 (0) | 15 | 4 (27) |
| Indigenous | 3 | 1 (33) | 0 | 0 (0) | 4 | 1 (25) |
| p value | | 1.000 | | -- | | 1.000 |

Table 4.3.10 HCV antibody prevalence by main language spoken at home by parents, gender and survey year

| Northern Territory Main language spoken at home by parents | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| English speaking | 39 | 14 (36) | 25 | 11 (44) | 64 | 25 (39) |
| Non-English speaking | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2019 | | | | | | |
| English speaking | 48 | 22 (46) | 28 | 7 (25) | 77 | 29 (38) |
| Non-English speaking | 3 | 1 (33) | 1 | 0 (0) | 4 | 1 (25) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2020 | | | | | | |
| English speaking | 18 | 4 (22) | 9 | 0 (0) | 27 | 4 (15) |
| Non-English speaking | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | -- | | 1.000 |
| 2021 | | | | | | |
| English speaking | 23 | 6 (26) | 14 | 2 (14) | 37 | 8 (22) |
| Non-English speaking | 1 | 1 (100) | 0 | 0 (0) | 1 | 1 (100) |
| p value | | 0.292 | | -- | | 0.237 |
| 2022 | | | | | | |
| English speaking | 14 | 5 (36) | 4 | 0 (0) | 20 | 5 (25) |
| Non-English speaking | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |

Table 4.3.11 HCV antibody prevalence by region/country of birth, gender and survey year

| Northern Territory Region/Country of birth | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Australia | 38 | 14 (37) | 24 | 11 (46) | 62 | 25 (40) |
| Other Oceania | 2 | 1 (50) | 0 | 0 (0) | 2 | 1 (50) |
| Asia | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| UK & Ireland | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| Other | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2019 | | | | | | |
| Australia | 47 | 21 (45) | 28 | 7 (25) | 76 | 28 (37) |
| Other Oceania | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 1 | 1 (100) | 0 | 0 (0) | 1 | 1 (100) |
| Other | 2 | 1 (50) | 1 | 0 (0) | 3 | 1 (33) |
| p value | | 0.849 | | 1.000 | | 0.824 |
| 2020 | | | | | | |
| Australia | 18 | 5 (28) | 10 | 0 (0) | 28 | 5 (18) |
| Other Oceania | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| Other | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2021 | | | | | | |
| Australia | 23 | 7 (30) | 14 | 2 (14) | 37 | 9 (24) |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| Other | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2022 | | | | | | |
| Australia | 13 | 5 (38) | 4 | 0 (0) | 19 | 5 (26) |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| Other | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | 1.000 | | -- | | 1.000 |

HCV RNA prevalence

Table 4.4.1 HCV RNA prevalence by gender and survey year

| Northern Territory Survey year | Male | | Female | | Total | |
|-----------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | 27 | 6 (22) | 16 | 4 (25) | 43 | 10 (23) |
| 2019* | 36 | 4 (11) | 18 | 2 (11) | 56 | 6 (11) |
| 2019 | 13 | 0 (0) | 5 | 0 (0) | 18 | 0 (0) |
| 2021 | 22 | 3 (14) | 11 | 0 (0) | 33 | 3 (9) |
| 2022 | 14 | 3 (21) | 4 | 0 (0) | 20 | 3 (15) |
| X ² p trend | | 0.799 | | 0.029 | | 0.174 |

* 2018, 2019: Weighted for gender and HCV antibody status

Totals include respondents where gender was reported as other or not reported

Table 4.4.2 HCV RNA prevalence by sexual identity, gender and survey year

| Northern Territory Sexual identity | Male | | Female | | Total | |
|---------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heterosexual | 27 | 6 (22) | 12 | 3 (25) | 39 | 9 (23) |
| Bisexual | 0 | 0 (0) | 2 | 0 (0) | 2 | 0 (0) |
| Homosexual | 0 | 0 (0) | 2 | 1 (50) | 2 | 1 (50) |
| p value | | -- | | 0.566 | | 0.549 |
| 2019* | | | | | | |
| Heterosexual | 28 | 4 (14) | 15 | 1 (7) | 43 | 5 (12) |
| Bisexual | 3 | 0 (0) | 1 | 0 (0) | 4 | 0 (0) |
| Homosexual | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) |
| p value | | 0.731 | | 0.935 | | 0.698 |
| 2020 | | | | | | |
| Heterosexual | 13 | 0 (0) | 3 | 0 (0) | 16 | 0 (0) |
| Bisexual | 0 | 0 (0) | 2 | 0 (0) | 2 | 0 (0) |
| Homosexual | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Heterosexual | 17 | 2 (12) | 8 | 0 (0) | 25 | 2 (8) |
| Bisexual | 3 | 1 (33) | 2 | 0 (0) | 5 | 1 (20) |
| Homosexual | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) |
| p value | | 0.564 | | -- | | 0.629 |
| 2022 | | | | | | |
| Heterosexual | 12 | 3 (25) | 3 | 0 (0) | 15 | 3 (20) |
| Bisexual | 2 | 0 (0) | 1 | 0 (0) | 3 | 0 (0) |
| Homosexual | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | 1.000 | | -- | | 1.000 |

Table 4.4.3 HCV RNA prevalence by age group, gender and survey year

| Northern Territory Age group | Male | | Female | | Total | |
|---------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <25 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 25-34 years | 2 | 0 (0) | 3 | 1 (33) | 5 | 1 (20) |
| 35-44 years | 11 | 2 (18) | 4 | 2 (50) | 16 | 4 (25) |
| 45+ years | 11 | 4 (36) | 9 | 1 (11) | 20 | 5 (25) |
| p value | | 0.575 | | 0.200 | | 0.889 |
| 2019* | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 4 | 0 (0) | 3 | 0 (0) | 7 | 0 (0) |
| 35-44 years | 7 | 0 (0) | 12 | 2 (17) | 19 | 2 (11) |
| 45+ years | 24 | 4 (17) | 6 | 0 (0) | 30 | 4 (13) |
| p value | | 0.405 | | 0.464 | | 0.617 |
| 2020 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| 35-44 years | 1 | 0 (0) | 2 | 0 (0) | 3 | 0 (0) |
| 45+ years | 10 | 0 (0) | 3 | 0 (0) | 13 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| 35-44 years | 6 | 1 (17) | 4 | 0 (0) | 10 | 1 (10) |
| 45+ years | 14 | 2 (14) | 7 | 0 (0) | 21 | 2 (10) |
| p value | | 0.832 | | -- | | 0.898 |
| 2022 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 35-44 years | 4 | 0 (0) | 1 | 0 (0) | 5 | 0 (0) |
| 45+ years | 9 | 3 (33) | 3 | 0 (0) | 14 | 3 (21) |
| p value | | 0.604 | | -- | | 0.601 |

Table 4.4.4 HCV RNA prevalence by years since first injection, gender and survey year

| Northern Territory Years of injection | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <3 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 3 to 10 years | 2 | 1 (50) | 1 | 0 (0) | 3 | 1 (33) |
| 11+ years | 22 | 4 (18) | 13 | 4 (31) | 35 | 8 (23) |
| p value | | 0.539 | | 0.562 | | 0.790 |
| 2019* | | | | | | |
| <3 years | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| 3 to 10 years | 2 | 0 (0) | 4 | 0 (0) | 6 | 0 (0) |
| 11+ years | 32 | 4 (13) | 16 | 2 (13) | 47 | 6 (13) |
| p value | | 0.566 | | 0.720 | | 0.606 |
| 2020 | | | | | | |
| <3 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 3 to 10 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 11+ years | 12 | 0 (0) | 5 | 0 (0) | 17 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| <3 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 3 to 10 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 11+ years | 20 | 3 (15) | 11 | 0 (0) | 31 | 3 (10) |
| p value | | 0.676 | | -- | | 0.744 |
| 2022 | | | | | | |
| <3 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 3 to 10 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 11+ years | 12 | 3 (25) | 4 | 0 (0) | 18 | 3 (17) |
| p value | | -- | | -- | | -- |

Table 4.4.5 HCV RNA prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| Northern Territory Receptively shared syringe last month | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| No receptive sharing | 20 | 5 (25) | 12 | 3 (25) | 33 | 8 (24) |
| Receptive sharing | 5 | 1 (20) | 3 | 1 (33) | 8 | 2 (25) |
| p value | | 0.808 | | 0.956 | | 0.869 |
| 2019* | | | | | | |
| No receptive sharing | 29 | 4 (14) | 17 | 1 (6) | 46 | 5 (11) |
| Receptive sharing | 4 | 0 (0) | 3 | 1 (33) | 7 | 1 (14) |
| p value | | 0.496 | | 0.153 | | 0.805 |
| 2020 | | | | | | |
| No receptive sharing | 9 | 0 (0) | 5 | 0 (0) | 14 | 0 (0) |
| Receptive sharing | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No receptive sharing | 14 | 1 (7) | 10 | 0 (0) | 24 | 1 (4) |
| Receptive sharing | 4 | 1 (25) | 1 | 0 (0) | 5 | 1 (20) |
| p value | | 0.316 | | -- | | 0.204 |
| 2022 | | | | | | |
| No receptive sharing | 13 | 3 (23) | 4 | 0 (0) | 19 | 3 (16) |
| Receptive sharing | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |

Table 4.4.6 HCV RNA prevalence by last drug injected, gender and survey year

| Northern Territory Last drug injected | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Methamphetamine | 13 | 4 (31) | 5 | 1 (20) | 18 | 5 (28) |
| Pharm. Opioids | 8 | 2 (25) | 7 | 2 (29) | 16 | 4 (25) |
| OAT | 4 | 0 (0) | 2 | 1 (50) | 6 | 1 (17) |
| Other drugs | 0 | 0 (0) | 2 | 0 (0) | 2 | 0 (0) |
| p value | | 0.467 | | 0.651 | | 0.755 |
| 2019* | | | | | | |
| Methamphetamine | 15 | 2 (13) | 11 | 1 (9) | 26 | 3 (12) |
| Pharm. Opioids | 12 | 2 (17) | 6 | 0 (0) | 18 | 2 (11) |
| OAT | 6 | 0 (0) | 1 | 0 (0) | 7 | 0 (0) |
| Other drugs | 3 | 0 (0) | 2 | 0 (0) | 5 | 0 (0) |
| p value | | 0.746 | | 0.849 | | 0.707 |
| 2020 | | | | | | |
| Methamphetamine | 5 | 0 (0) | 4 | 0 (0) | 9 | 0 (0) |
| Pharm. Opioids | 4 | 0 (0) | 1 | 0 (0) | 5 | 0 (0) |
| OAT | 3 | 0 (0) | 0 | 0 (0) | 3 | 0 (0) |
| Other drugs | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Methamphetamine | 11 | 1 (9) | 10 | 0 (0) | 21 | 1 (5) |
| Pharm. Opioids | 3 | 0 (0) | 0 | 0 (0) | 3 | 0 (0) |
| OAT | 4 | 1 (25) | 0 | 0 (0) | 4 | 1 (25) |
| Other drugs | 2 | 0 (0) | 1 | 0 (0) | 3 | 0 (0) |
| p value | | 0.667 | | -- | | 0.425 |
| 2022 | | | | | | |
| Methamphetamine | 7 | 2 (29) | 3 | 0 (0) | 11 | 2 (18) |
| Pharm. Opioids | 5 | 1 (20) | 1 | 0 (0) | 6 | 1 (17) |
| OAT | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| Other drugs | 1 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| p value | | 0.858 | | -- | | 0.890 |

OAT: methadone, buprenorphine or buprenorphine/naloxone

Table 4.4.7 HCV RNA prevalence by frequency of drug injection last month, gender and survey year

| Northern Territory Frequency of injection last month | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| Less than daily | 8 | 3 (38) | 6 | 1 (17) | 14 | 4 (29) |
| Daily or more | 18 | 3 (17) | 9 | 3 (33) | 27 | 6 (22) |
| Not last month | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| p value | | 0.289 | | 0.533 | | 0.816 |
| 2019* | | | | | | |
| Less than daily | 11 | 2 (18) | 5 | 1 (20) | 15 | 3 (20) |
| Daily or more | 23 | 2 (9) | 15 | 1 (7) | 37 | 3 (8) |
| Not last month | 2 | 0 (0) | 1 | 0 (0) | 3 | 0 (0) |
| p value | | 0.820 | | 0.630 | | 0.607 |
| 2020 | | | | | | |
| Less than daily | 5 | 0 (0) | 2 | 0 (0) | 7 | 0 (0) |
| Daily or more | 6 | 0 (0) | 3 | 0 (0) | 9 | 0 (0) |
| Not last month | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Less than daily | 9 | 1 (11) | 7 | 0 (0) | 16 | 1 (6) |
| Daily or more | 9 | 1 (11) | 4 | 0 (0) | 13 | 1 (8) |
| Not last month | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| p value | | 0.884 | | -- | | 0.918 |
| 2022 | | | | | | |
| Less than daily | 6 | 2 (33) | 3 | 0 (0) | 11 | 2 (18) |
| Daily or more | 7 | 1 (14) | 1 | 0 (0) | 8 | 1 (13) |
| Not last month | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | 0.654 | | -- | | 1.000 |

Table 4.4.8 HCV RNA prevalence by imprisonment last year, gender and survey year

| Northern Territory Imprisonment last year | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| No imprisonment | 24 | 4 (17) | 14 | 3 (21) | 38 | 8 (21) |
| Imprisonment | 2 | 1 (50) | 1 | 1 (100) | 3 | 2 (67) |
| p value | | 0.395 | | 0.134 | | 0.129 |
| 2019* | | | | | | |
| No imprisonment | 31 | 4 (13) | 18 | 1 (6) | 48 | 5 (10) |
| Imprisonment | 2 | 0 (0) | 2 | 1 (50) | 4 | 1 (25) |
| p value | | 0.561 | | 0.059 | | 0.452 |
| 2020 | | | | | | |
| No imprisonment | 11 | 0 (0) | 4 | 0 (0) | 15 | 0 (0) |
| Imprisonment | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No imprisonment | 19 | 3 (16) | 10 | 0 (0) | 29 | 3 (10) |
| Imprisonment | 3 | 0 (0) | 1 | 0 (0) | 4 | 0 (0) |
| p value | | 0.459 | | -- | | 0.500 |
| 2022 | | | | | | |
| No imprisonment | 14 | 3 (21) | 3 | 0 (0) | 19 | 3 (16) |
| Imprisonment | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |

Table 4.4.9 HCV RNA prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Northern Territory Aboriginal and Torres Strait Islander origin | Male | | Female | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| Non Indigenous | 20 | 4 (20) | 7 | 2 (29) | 27 | 6 (22) |
| Indigenous | 7 | 2 (29) | 9 | 2 (22) | 16 | 4 (25) |
| p value | | 0.681 | | 0.733 | | 0.525 |
| 2019* | | | | | | |
| Non Indigenous | 32 | 2 (6) | 9 | 0 (0) | 41 | 2 (5) |
| Indigenous | 4 | 2 (50) | 11 | 2 (18) | 15 | 4 (27) |
| p value | | 0.108 | | 0.183 | | 0.102 |
| 2020 | | | | | | |
| Non Indigenous | 9 | 0 (0) | 5 | 0 (0) | 14 | 0 (0) |
| Indigenous | 3 | 0 (0) | 0 | 0 (0) | 3 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Non Indigenous | 11 | 2 (18) | 4 | 0 (0) | 15 | 2 (13) |
| Indigenous | 11 | 1 (9) | 7 | 0 (0) | 18 | 1 (6) |
| p value | | 0.534 | | -- | | 0.439 |
| 2022 | | | | | | |
| Non Indigenous | 10 | 3 (30) | 4 | 0 (0) | 15 | 3 (20) |
| Indigenous | 3 | 0 (0) | 0 | 0 (0) | 4 | 0 (0) |
| p value | | 0.528 | | -- | | 1.000 |

Table 4.4.10 HCV RNA prevalence by main language spoken at home by parents, gender and survey year

| Northern Territory Main language spoken at home by parents | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| English speaking | 26 | 6 (23) | 16 | 4 (25) | 42 | 10 (24) |
| Non-English speaking | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2019* | | | | | | |
| English speaking | 34 | 4 (12) | 19 | 2 (11) | 53 | 6 (11) |
| Non-English speaking | 2 | 0 (0) | 1 | 0 (0) | 3 | 0 (0) |
| p value | | 0.644 | | 0.746 | | 0.569 |
| 2020 | | | | | | |
| English speaking | 12 | 0 (0) | 5 | 0 (0) | 17 | 0 (0) |
| Non-English speaking | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| English speaking | 21 | 3 (14) | 11 | 0 (0) | 32 | 3 (9) |
| Non-English speaking | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | 0.684 | | -- | | 0.748 |
| 2022 | | | | | | |
| English speaking | 14 | 3 (21) | 4 | 0 (0) | 20 | 3 (15) |
| Non-English speaking | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |

Table 4.4.11 HCV RNA prevalence by region/country of birth, gender and survey year

| Northern Territory | | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | |
| 2018* | | | | | | | |
| Australia | 26 | 6 (23) | 15 | 4 (27) | 41 | 10 (24) | |
| Other Oceania | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) | |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| UK & Ireland | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| Other | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) | |
| p value | | 0.597 | | 0.533 | | 0.719 | |
| 2019* | | | | | | | |
| Australia | 32 | 4 (13) | 19 | 2 (11) | 51 | 6 (12) | |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| UK & Ireland | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) | |
| Other | 2 | 0 (0) | 1 | 0 (0) | 3 | 0 (0) | |
| p value | | 0.859 | | 0.746 | | 0.823 | |
| 2020 | | | | | | | |
| Australia | 12 | 0 (0) | 5 | 0 (0) | 17 | 0 (0) | |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| UK & Ireland | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| Other | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| p value | | -- | | -- | | -- | |
| 2021 | | | | | | | |
| Australia | 21 | 3 (14) | 11 | 0 (0) | 32 | 3 (9) | |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| UK & Ireland | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| Other | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| p value | | -- | | -- | | -- | |
| 2022 | | | | | | | |
| Australia | 13 | 3 (23) | 4 | 0 (0) | 19 | 3 (16) | |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| UK & Ireland | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) | |
| Other | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| p value | | 1.000 | | -- | | 1.000 | |

Queensland

Table 5.1.1 Number (percentage) of respondents by demographic characteristics and survey year

| Queensland | 2018 | 2019 | 2020* | 2021* | 2022 |
|---|----------|----------|----------|----------|----------|
| Number of sites | 9 | 10 | 7 | 11 | 10 |
| N° surveyed | N=633 | N=574 | N=391 | N=458 | N=328 |
| Response rate (%) | 41% | 34% | 33% | 31% | 26% |
| Gender (%) | | | | | |
| Male | 441 (70) | 407 (71) | 285 (73) | 307 (67) | 229 (70) |
| Female | 191 (30) | 160 (28) | 104 (27) | 145 (32) | 92 (28) |
| Other | 1 (<1) | 3 (1) | 1 (<1) | 6 (1) | 6 (2) |
| Not reported | 0 (0) | 4 (1) | 1 (<1) | 0 (0) | 1 (<1) |
| Sexual identity (%) | | | | | |
| Heterosexual | 506 (80) | 427 (74) | 315 (81) | 382 (83) | 254 (77) |
| Bisexual | 62 (10) | 63 (11) | 38 (10) | 42 (9) | 34 (10) |
| Gay/Lesbian | 25 (4) | 27 (5) | 12 (3) | 24 (5) | 20 (6) |
| Other | -- -- | -- -- | 8 (2) | 9 (2) | 16 (5) |
| Not reported | 40 (6) | 57 (10) | 18 (5) | 1 (<1) | 4 (1) |
| Aboriginal and Torres Strait Islander origin (%) | | | | | |
| No | 511 (81) | 452 (79) | 317 (81) | 365 (80) | 268 (82) |
| Yes | 115 (18) | 116 (20) | 64 (16) | 79 (17) | 55 (17) |
| Not reported | 7 (1) | 6 (1) | 10 (3) | 14 (3) | 5 (2) |
| Region/country of birth (%) | | | | | |
| Australia | 570 (90) | 525 (91) | 352 (90) | 407 (89) | 293 (89) |
| Other Oceania | 20 (3) | 12 (2) | 15 (4) | 11 (2) | 11 (3) |
| Asia | 0 (0) | 4 (1) | 1 (<1) | 3 (1) | 1 (<1) |
| UK & Ireland | 24 (4) | 11 (2) | 11 (3) | 11 (2) | 8 (2) |
| Other | 15 (2) | 17 (3) | 8 (2) | 14 (3) | 8 (2) |
| Not reported | 4 (1) | 5 (1) | 4 (1) | 12 (3) | 7 (2) |
| Main language spoken at home by parents (%) | | | | | |
| English | 612 (97) | 550 (96) | 382 (98) | 441 (96) | 316 (96) |
| Non-English | 18 (3) | 22 (4) | 9 (2) | 11 (2) | 12 (4) |
| Not reported | 3 (<1) | 2 (<1) | 0 (0) | 6 (1) | 0 (0) |
| Age and time since first injection (years) | | | | | |
| <i>Median age</i> | 41 | 42 | 42 | 42 | 43 |
| <i>Age range</i> | 14-78 | 16-70 | 19-69 | 16-72 | 16-71 |
| Age group (%) | | | | | |
| <25 years | 37 (6) | 30 (5) | 19 (5) | 20 (4) | 15 (5) |
| 25+ years | 595 (94) | 543 (95) | 372 (95) | 438 (96) | 310 (95) |
| Not reported | 1 (<1) | 1 (<1) | 0 (0) | 0 (0) | 3 (<1) |
| Median age first injection | | | | | |
| <i>Median age first injection</i> | 19 | 19 | 19 | 19 | 19 |
| <i>Age range</i> | 10-64 | 10-60 | 11-59 | 12-52 | 10-55 |
| <i>N° not reported</i> | 22 | 26 | 4 | 14 | 17 |
| Median yrs since first injection | | | | | |
| <i>Median yrs since first injection</i> | 21 | 21 | 23 | 22 | 22 |
| <i>Range</i> | <1-52 | <1-51 | <1-53 | <1-58 | <1-51 |
| Years since first injection | | | | | |
| <3 years | 42 (7) | 51 (9) | 32 (8) | 27 (6) | 21 (6) |
| 3+ years | 569 (90) | 497 (87) | 354 (91) | 415 (91) | 290 (88) |
| Not reported | 22 (3) | 26 (5) | 5 (1) | 16 (3) | 17 (5) |
| Imprisonment last year (%) | | | | | |
| No | 535 (85) | 480 (84) | 336 (86) | 385 (84) | 288 (88) |
| Yes | 72 (11) | 66 (12) | 39 (10) | 46 (10) | 29 (9) |
| Not reported | 26 (4) | 28 (5) | 16 (4) | 27 (6) | 11 (3) |
| N° in prison | | | | | |
| <i>N° in prison</i> | N=72 | N=66 | N=39 | N=46 | N=29 |
| Injected in prison | 29 (40) | 30 (45) | 18 (46) | 27 (59) | 12 (41) |

* Implementation of the 2020 and 2021 ANSPS was impacted by the COVID-19 pandemic, see page 1 for more information

Table 5.1.2 Number (percentage) of respondents by last drug injected, frequency of injecting and survey year

| Queensland N° surveyed | 2018 N=633 | 2019 N=574 | 2020 N=391 | 2021 N=458 | 2022 N=328 |
|---|---------------|---------------|---------------|---------------|---------------|
| Drug last injected (%) | | | | | |
| Cocaine | 10 (2) | 3 (1) | 2 (1) | 2 (<1) | 3 (1) |
| Methamphetamine | 304 (48) | 281 (49) | 156 (40) | 194 (42) | 142 (43) |
| Heroin | 84 (13) | 93 (16) | 63 (16) | 72 (16) | 61 (19) |
| Pharm. opioids | 104 (16) | 75 (13) | 69 (18) | 73 (16) | 40 (12) |
| Methadone | 28 (4) | 29 (5) | 22 (6) | 20 (4) | 13 (4) |
| Buprenorphine | 33 (5) | 22 (4) | 23 (6) | 25 (5) | 12 (4) |
| Buprenorphine/naloxone | 17 (3) | 9 (2) | 4 (1) | 12 (3) | 8 (2) |
| PIEDs | 28 (4) | 27 (5) | 31 (8) | 34 (7) | 30 (9) |
| More than one | 21 (3) | 26 (5) | 16 (4) | 15 (3) | 11 (3) |
| Other | 3 (<1) | 8 (1) | 2 (1) | 4 (1) | 3 (1) |
| Not reported | 1 (<1) | 1 (<1) | 3 (1) | 7 (2) | 5 (2) |
| Ever injected opioids | | | | | |
| No | 151 (24) | 143 (25) | 101 (26) | 120 (26) | 99 (30) |
| Yes | 480 (76) | 428 (75) | 290 (74) | 333 (73) | 227 (69) |
| Not reported | 2 (<1) | 3 (1) | 0 (0) | 5 (1) | 2 (1) |
| Frequency of injection last month (%) | | | | | |
| Not last month | 41 (6) | 47 (8) | 46 (12) | 73 (16) | 35 (11) |
| Less than weekly | 90 (14) | 86 (15) | 59 (15) | 75 (16) | 63 (19) |
| Weekly not daily | 160 (25) | 133 (23) | 78 (20) | 89 (19) | 68 (21) |
| Daily or more | 332 (52) | 305 (53) | 204 (52) | 215 (47) | 157 (48) |
| Not reported | 10 (2) | 3 (1) | 4 (1) | 6 (1) | 5 (2) |
| Present when someone injected for the first time in the last 12 months (%) | | | | | |
| No | 499 (79) | 457 (80) | 320 (82) | 377 (82) | 284 (87) |
| Yes | 117 (18) | 96 (17) | 67 (17) | 73 (16) | 39 (12) |
| Not reported | 17 (3) | 21 (4) | 4 (1) | 8 (2) | 5 (2) |
| Overdosed in last 12 months | | | | | |
| No | 521 (82) | 478 (83) | 329 (84) | 400 (87) | 290 (88) |
| Yes | 96 (15) | 78 (14) | 58 (15) | 49 (11) | 34 (10) |
| Not reported | 16 (3) | 18 (3) | 4 (1) | 9 (2) | 4 (1) |

Table 5.1.3 Number (percentage) of respondents by injecting behaviour in the last month and survey year

| Queensland | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|-------------|-------------|-------------|-------------|-------------|
| N° injected last month | N=582 | N=524 | N=341 | N=379 | N=288 |
| Places injected last month (%) | | | | | |
| All private | 378 (65) | 327 (62) | 225 (66) | 249 (66) | 187 (65) |
| Any public | 198 (34) | 194 (37) | 116 (34) | 128 (34) | 101 (35) |
| Not reported | 6 (1) | 3 (1) | 0 (0) | 2 (1) | 0 (0) |
| Use of new and sterile needles and syringes last month (%) | | | | | |
| All injections | 445 (76) | 416 (79) | 274 (80) | 306 (81) | 221 (77) |
| Most of the time | 107 (18) | 87 (17) | 55 (16) | 57 (15) | 52 (18) |
| Half of the time | 8 (1) | 8 (2) | 6 (2) | 9 (2) | 9 (3) |
| Some of the time | 14 (2) | 8 (2) | 5 (1) | 5 (1) | 2 (1) |
| Not last month | 2 (<1) | 2 (<1) | 0 (0) | 2 (1) | 2 (1) |
| Not reported | 6 (1) | 3 (1) | 1 (<1) | 0 (0) | 2 (1) |
| Re-used someone else's used needle & syringe last month (%) | | | | | |
| None | 490 (84) | 464 (89) | 292 (86) | 327 (86) | 247 (86) |
| Once | 29 (5) | 13 (2) | 16 (5) | 12 (3) | 18 (6) |
| Twice | 17 (3) | 17 (3) | 10 (3) | 13 (3) | 12 (4) |
| 3-5 times | 23 (4) | 15 (3) | 13 (4) | 14 (4) | 5 (2) |
| >5 times | 18 (3) | 12 (2) | 9 (3) | 12 (3) | 6 (2) |
| Not reported | 5 (1) | 3 (1) | 1 (<1) | 1 (<1) | 0 (0) |
| Number of people needle & syringe was re-used after last month (%) | | | | | |
| None | 490 (84) | 464 (89) | 292 (86) | 327 (86) | 247 (86) |
| One | 37 (6) | 19 (4) | 18 (5) | 22 (6) | 13 (5) |
| Two | 7 (1) | 4 (1) | 7 (2) | 4 (1) | 7 (2) |
| Three to five | 5 (1) | 6 (1) | 3 (1) | 2 (1) | 1 (<1) |
| More than five | 10 (2) | 4 (1) | 2 (1) | 4 (1) | 3 (1) |
| Don't know | 20 (3) | 14 (3) | 9 (3) | 13 (3) | 7 (2) |
| Not reported | 13 (2) | 13 (2) | 10 (3) | 7 (2) | 10 (3) |
| Relationship to people needle & syringe was used after last month (%) [more than one could be selected] | | | | | |
| Regular sex partner | 29 (5) | 19 (4) | 16 (5) | 15 (4) | 13 (5) |
| Casual sex partner | 5 (1) | 3 (1) | 4 (1) | 3 (1) | 3 (1) |
| Close friend | 33 (6) | 19 (4) | 15 (4) | 14 (4) | 11 (4) |
| Acquaintance | 17 (3) | 9 (2) | 5 (1) | 10 (3) | 6 (2) |
| Other | 7 (1) | 6 (1) | 4 (1) | 7 (2) | 1 (<1) |
| Equipment used after someone else last month (%) [more than one could be selected] | | | | | |
| Spoon | 116 (20) | 102 (19) | 43 (13) | 74 (20) | 50 (17) |
| Water | 105 (18) | 102 (19) | 47 (14) | 65 (17) | 45 (16) |
| Filter | 63 (11) | 59 (11) | 17 (5) | 36 (10) | 19 (7) |
| Drug mix | 51 (9) | 63 (12) | 21 (6) | 38 (10) | 20 (7) |
| None | 413 (71) | 368 (70) | 263 (77) | 274 (72) | 210 (73) |
| Injected by someone after that person injected themselves or others last month (%) | | | | | |
| No | 475 (82) | 442 (84) | 305 (89) | 325 (86) | 237 (82) |
| Yes | 97 (17) | 78 (15) | 35 (10) | 53 (14) | 47 (16) |
| Not reported | 10 (2) | 4 (1) | 1 (<1) | 1 (<1) | 4 (1) |
| Source of needle acquisition | | | | | |
| Needle Syringe Program | 533 (92) | 473 (90) | 309 (91) | 339 (89) | 242 (84) |
| Chemist/Pharmacy | 143 (25) | 161 (31) | 89 (26) | 96 (25) | 69 (24) |
| Personal sources | 44 (8) | 55 (11) | 24 (7) | 37 (10) | 20 (7) |
| Dispensing/Vending Machine | 66 (11) | 87 (17) | 45 (13) | 47 (12) | 26 (9) |
| Other sources | 21 (4) | 12 (2) | 9 (3) | 9 (2) | 11 (4) |

Table 5.1.4 Number (percentage) of respondents by treatment for drug use and survey year

| Queensland | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|----------|----------|----------|----------|----------|
| N° surveyed | N=633 | N=574 | N=391 | N=458 | N=328 |
| Ever any treatment/therapy for drug use (%) | | | | | |
| No | 187 (30) | 179 (31) | 126 (32) | 152 (33) | 114 (35) |
| Yes | 443 (70) | 394 (69) | 264 (68) | 305 (67) | 213 (65) |
| Not reported | 3 (<1) | 1 (<1) | 1 (<1) | 1 (<1) | 1 (<1) |
| History of methadone maintenance treatment (%) | | | | | |
| Currently | 87 (14) | 81 (14) | 58 (15) | 42 (9) | 44 (13) |
| Previously | 141 (22) | 113 (20) | 77 (20) | 101 (22) | 65 (20) |
| Never | 391 (62) | 369 (64) | 247 (63) | 297 (65) | 210 (64) |
| Not reported | 14 (2) | 11 (2) | 9 (2) | 18 (4) | 9 (3) |
| History of buprenorphine (Subutex®) treatment (%) | | | | | |
| Currently | 56 (9) | 52 (9) | 45 (12) | 52 (11) | 31 (9) |
| Previously | 147 (23) | 131 (23) | 99 (25) | 93 (20) | 67 (20) |
| Never | 415 (66) | 381 (66) | 240 (61) | 299 (65) | 221 (67) |
| Not reported | 15 (2) | 10 (2) | 7 (2) | 14 (3) | 9 (3) |
| History of buprenorphine-naloxone (Suboxone®) treatment (%) | | | | | |
| Currently | 54 (9) | 44 (8) | 26 (7) | 36 (8) | 16 (5) |
| Previously | 139 (22) | 125 (22) | 90 (23) | 100 (22) | 73 (22) |
| Never | 428 (68) | 395 (69) | 273 (70) | 317 (69) | 234 (71) |
| Not reported | 12 (2) | 10 (2) | 2 (1) | 5 (1) | 5 (2) |

Table 5.1.5 Number (percentage) of respondents by testing for HIV and HCV, HCV treatment by survey year

| Queensland | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|----------|----------|----------|----------|----------|
| N° surveyed | N=619 | N=574 | N=391 | N=458 | N=328 |
| Previous HIV test (%) | | | | | |
| Yes, ever | 534 (84) | 489 (85) | 328 (84) | 350 (76) | 251 (77) |
| Yes, last year | 312 (49) | 287 (50) | 180 (46) | 146 (32) | 118 (36) |
| >1 year ago | 222 (35) | 202 (35) | 148 (38) | 204 (45) | 133 (41) |
| Never tested | 78 (12) | 69 (12) | 56 (14) | 95 (21) | 68 (21) |
| Not reported | 21 (3) | 16 (3) | 7 (2) | 13 (3) | 9 (3) |
| Previous HCV test (%) | | | | | |
| Yes, ever | 540 (85) | 482 (84) | 341 (87) | 367 (80) | 267 (81) |
| Yes, last year | 342 (54) | 322 (56) | 192 (49) | 198 (43) | 144 (44) |
| >1 year ago | 198 (31) | 160 (28) | 149 (38) | 169 (37) | 123 (38) |
| Never tested | 37 (6) | 46 (8) | 29 (7) | 59 (13) | 41 (13) |
| Not reported | 56 (9) | 46 (8) | 21 (5) | 32 (7) | 20 (6) |
| Lifetime treatment for HCV (%)# | | | | | |
| N° self-reported HCV diagnosis | N=172 | N=146 | N=100 | N=107 | N=54 |
| Antiviral treatment | 101 (59) | 92 (63) | 65 (65) | 80 (75) | 33 (61) |
| No antiviral treatment | 69 (40) | 54 (37) | 35 (35) | 25 (23) | 21 (39) |
| Not reported | 2 (1) | 0 (0) | 0 (0) | 2 (2) | 0 (0) |
| Treatment for HCV in past 12 months (%)** | | | | | |
| N° self-reported HCV diagnosis | N=129 | N=113 | N=62 | N=54 | N=33 |
| Antiviral treatment | 50 (39) | 48 (42) | 23 (37) | 20 (37) | 11 (33) |
| No antiviral treatment | 77 (60) | 65 (58) | 39 (63) | 32 (59) | 22 (67) |
| Not reported | 2 (2) | 0 (0) | 0 (0) | 2 (4) | 0 (0) |

among people who tested HCV antibody positive and did not report spontaneous clearance

* excludes people who reported treatment induced clearance more than 12 months previously

Table 5.1.6 Number (percentage) of respondents by sexual behaviour in the month prior to survey and survey year

| Queensland | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|----------|----------|----------|----------|----------|
| N° surveyed | N=633 | N=574 | N=391 | N=458 | N=328 |
| Sex with a regular partner last month (%) | | | | | |
| No | 319 (50) | 301 (52) | 217 (56) | 234 (51) | 172 (52) |
| Yes | 297 (47) | 256 (45) | 168 (43) | 210 (46) | 149 (45) |
| Not reported | 17 (3) | 17 (3) | 6 (2) | 14 (3) | 7 (2) |
| Condom used with regular partner last month (%) | | | | | |
| N° surveyed | N=297 | N=256 | N=168 | N=210 | N=149 |
| Never | 230 (77) | 197 (77) | 133 (79) | 161 (77) | 116 (78) |
| Sometimes | 32 (11) | 26 (10) | 12 (7) | 28 (13) | 17 (11) |
| Every time | 16 (5) | 22 (9) | 15 (9) | 14 (7) | 8 (5) |
| Not reported | 19 (6) | 11 (4) | 8 (5) | 7 (3) | 8 (5) |
| Sex with other partner(s) last month (%) | | | | | |
| No | 500 (79) | 457 (80) | 317 (81) | 366 (80) | 265 (81) |
| Yes | 108 (17) | 96 (17) | 58 (15) | 70 (15) | 48 (15) |
| Not reported | 25 (4) | 21 (4) | 16 (4) | 22 (5) | 15 (5) |
| Condom used with other partner(s) last month (%) | | | | | |
| N° surveyed | N=108 | N=96 | N=58 | N=70 | N=48 |
| Never | 49 (45) | 43 (45) | 28 (48) | 31 (44) | 26 (54) |
| Sometimes | 36 (33) | 25 (26) | 15 (26) | 22 (31) | 14 (29) |
| Every time | 17 (16) | 21 (22) | 14 (24) | 12 (17) | 6 (13) |
| Not reported | 6 (6) | 7 (7) | 1 (2) | 5 (7) | 2 (4) |
| Sex work last month (%) | | | | | |
| No | 581 (92) | 528 (92) | 363 (93) | 422 (92) | 304 (93) |
| Yes | 30 (5) | 21 (4) | 14 (4) | 14 (3) | 9 (3) |
| Not reported | 22 (3) | 25 (4) | 14 (4) | 22 (5) | 15 (5) |
| Condom used at last sex work (%) | | | | | |
| Yes | 16 (53) | 13 (62) | 8 (57) | 8 (57) | 7 (78) |

Table 5.1.7 Number (percentage) of respondents by COVID-19 related impacts, 2020-2022

| Queensland | 2020 | 2021 | 2022 |
|---|----------|----------|----------|
| N° surveyed | N=391 | N=458 | N=328 |
| COVID-19 test | | | |
| Yes | 78 (20) | 122 (27) | 244 (74) |
| No | 246 (63) | 205 (45) | 54 (16) |
| Don't know | 16 (4) | 12 (3) | -- -- |
| Not reported | 51 (13) | 119 (26) | 30 (9) |
| Self-reported positive COVID-19 test result | | | |
| Yes | 0 (0) | 0 (0) | 86 (35) |
| Yes, last 12 months | -- -- | -- -- | 80 (33) |
| Yes, more than a year ago | -- -- | -- -- | 6 (2) |
| COVID-19 vaccine uptake | | | |
| ≥1 dose | -- -- | 103 (22) | 228 (70) |
| COVID-19 vaccine doses | | | |
| One dose | -- -- | 53 (12) | 18 (5) |
| Two doses | -- -- | 50 (11) | 93 (28) |
| Three or more doses | -- -- | -- -- | 117 (36) |
| No doses | -- -- | 239 (52) | 68 (21) |
| Not reported | -- -- | 116 (25) | 32 (10) |
| Vaccine type * | | | N=228 |
| Astra Zeneca | -- -- | 48 (10) | 60 (26) |
| Pfizer | -- -- | 37 (8) | 149 (65) |
| Don't know | -- -- | 26 (6) | 32 (14) |
| Other | -- -- | -- -- | 27 (12) |
| Not reported | -- -- | 347 (76) | -- -- |
| * 2021 only one response permitted, 2022 multiple responses permitted | | | |
| Last vaccine location | | | N=328 |
| GP or medical centre | -- -- | -- -- | 89 (27) |
| Pharmacy/ chemist | -- -- | -- -- | 55 (17) |
| Vaccine hub | -- -- | -- -- | 59 (18) |
| Opioid treatment/ methadone clinic | -- -- | -- -- | 0 (0) |
| NSP | -- -- | -- -- | 4 (1) |
| Prison | -- -- | -- -- | 17 (5) |
| Other | -- -- | -- -- | 3 (1) |
| Multiple response | -- -- | -- -- | 12 (4) |
| Not reported | -- -- | -- -- | 89 (27) |
| Vaccine hesitancy - if not had three or more doses | | | |
| Haven't got around to it | -- -- | -- -- | 45 (14) |
| Difficulty booking appointment | -- -- | -- -- | 4 (1) |
| Concerns about side effects | -- -- | -- -- | 19 (6) |
| Concerns about vaccine safety | -- -- | -- -- | 26 (8) |
| Not at risk of serious illness | -- -- | -- -- | 21 (6) |
| Other | -- -- | -- -- | 46 (14) |
| Multiple response | -- -- | -- -- | 14 (4) |
| Not reported | -- -- | -- -- | 153 (47) |

HIV antibody prevalence

Table 5.2.1 HIV antibody prevalence by gender and survey year

| Queensland Survey year | Male | | Female | | Total | |
|---------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | 437 | 6 (1.4) | 189 | 2 (1.1) | 627 | 8 (1.3) |
| 2019 | 401 | 11 (2.7) | 157 | 0 (0.0) | 565 | 12 (2.1) |
| 2020 | 284 | 13 (4.6) | 102 | 1 (1.0) | 388 | 14 (3.6) |
| 2021 | 304 | 7 (2.3) | 145 | 0 (0.0) | 455 | 8 (1.8) |
| 2022 | 227 | 7 (3.1) | 91 | 0 (0.0) | 325 | 7 (2.2) |
| X ² p trend | | 0.196 | | 0.205 | | 0.362 |

Table 5.2.2 HIV antibody prevalence by sexual identity, gender and survey year

| Queensland Sexual identity | Male | | Female | | Total | |
|-------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | | | | | | |
| Heterosexual | 365 | 1 (0.3) | 136 | 2 (1.5) | 501 | 3 (0.6) |
| Bisexual | 25 | 2 (8.0) | 36 | 0 (0.0) | 61 | 2 (3.3) |
| Homosexual | 21 | 3 (14.3) | 3 | 0 (0.0) | 25 | 3 (12.0) |
| p value | | <0.001 | | 1.000 | | 0.001 |
| 2019 | | | | | | |
| Heterosexual | 322 | 1 (0.3) | 95 | 0 (0.0) | 419 | 1 (0.2) |
| Bisexual | 24 | 4 (16.7) | 36 | 0 (0.0) | 62 | 4 (6.5) |
| Homosexual | 21 | 6 (28.6) | 5 | 0 (0.0) | 27 | 7 (25.9) |
| p value | | <0.001 | | -- | | <0.001 |
| 2020 | | | | | | |
| Heterosexual | 245 | 2 (0.8) | 69 | 1 (1.5) | 314 | 3 (1.0) |
| Bisexual | 13 | 3 (23.1) | 21 | 0 (0.0) | 36 | 3 (8.3) |
| Homosexual | 9 | 7 (77.8) | 3 | 0 (0.0) | 12 | 7 (58.3) |
| p value | | <0.001 | | 1.000 | | <0.001 |
| 2021 | | | | | | |
| Heterosexual | 267 | 1 (0.4) | 110 | 0 (0.0) | 379 | 1 (0.3) |
| Bisexual | 18 | 0 (0.0) | 24 | 0 (0.0) | 42 | 0 (0.0) |
| Homosexual | 15 | 5 (33.3) | 6 | 0 (0.0) | 24 | 6 (25.0) |
| p value | | <0.001 | | -- | | <0.001 |
| 2022 | | | | | | |
| Heterosexual | 192 | 0 (0.0) | 59 | 0 (0.0) | 252 | 0 (0.0) |
| Bisexual | 13 | 0 (0.0) | 19 | 0 (0.0) | 34 | 0 (0.0) |
| Homosexual | 15 | 7 (46.7) | 5 | 0 (0.0) | 20 | 7 (35.0) |
| p value | | <0.001 | | -- | | <0.001 |

Table 5.2.3 HIV antibody prevalence by age group, gender and survey year

| Queensland Age group | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| <25 years | 21 | 0 (0.0) | 15 | 0 (0.0) | 36 | 0 (0.0) |
| 25-34 years | 75 | 1 (1.3) | 46 | 2 (4.4) | 122 | 3 (2.5) |
| 35-44 years | 158 | 2 (1.3) | 67 | 0 (0.0) | 225 | 2 (0.9) |
| 45+ years | 182 | 3 (1.7) | 61 | 0 (0.0) | 243 | 3 (1.2) |
| p value | | 1.000 | | 0.211 | | 0.671 |
| 2019 | | | | | | |
| <25 years | 18 | 0 (0.0) | 12 | 0 (0.0) | 30 | 0 (0.0) |
| 25-34 years | 62 | 1 (1.6) | 34 | 0 (0.0) | 99 | 2 (2.0) |
| 35-44 years | 147 | 2 (1.4) | 67 | 0 (0.0) | 217 | 2 (0.9) |
| 45+ years | 173 | 8 (4.6) | 44 | 0 (0.0) | 218 | 8 (3.7) |
| p value | | 0.330 | | -- | | 0.244 |
| 2020 | | | | | | |
| <25 years | 11 | 0 (0.0) | 7 | 1 (14.3) | 18 | 1 (5.6) |
| 25-34 years | 42 | 3 (7.1) | 20 | 0 (0.0) | 62 | 3 (4.8) |
| 35-44 years | 103 | 4 (3.9) | 40 | 0 (0.0) | 144 | 4 (2.8) |
| 45+ years | 128 | 6 (4.7) | 35 | 0 (0.0) | 164 | 6 (3.7) |
| p value | | 0.814 | | 0.069 | | 0.621 |
| 2021 | | | | | | |
| <25 years | 14 | 0 (0.0) | 5 | 0 (0.0) | 20 | 0 (0.0) |
| 25-34 years | 49 | 0 (0.0) | 30 | 0 (0.0) | 80 | 0 (0.0) |
| 35-44 years | 108 | 0 (0.0) | 49 | 0 (0.0) | 161 | 1 (0.6) |
| 45+ years | 133 | 7 (5.3) | 61 | 0 (0.0) | 194 | 7 (3.6) |
| p value | | 0.496 | | -- | | 0.120 |
| 2022 | | | | | | |
| <25 years | 9 | 0 (0.0) | 3 | 0 (0.0) | 15 | 0 (0.0) |
| 25-34 years | 36 | 0 (0.0) | 17 | 0 (0.0) | 55 | 0 (0.0) |
| 35-44 years | 71 | 3 (4.2) | 39 | 0 (0.0) | 111 | 3 (2.7) |
| 45+ years | 109 | 4 (3.7) | 31 | 0 (0.0) | 141 | 4 (2.8) |
| p value | | 0.761 | | -- | | 0.775 |

Table 5.2.4 HIV antibody prevalence by last drug injected, gender and survey year

| Queensland Last drug injected | Male | | Female | | Total | |
|----------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Heroin | 56 | 0 (0.0) | 27 | 0 (0.0) | 83 | 0 (0.0) |
| Methamphetamine | 200 | 6 (3.0) | 100 | 2 (2.0) | 301 | 8 (2.7) |
| Other opioids | 130 | 0 (0.0) | 50 | 0 (0.0) | 180 | 0 (0.0) |
| PIEDs | 27 | 0 (0.0) | 1 | 0 (0.0) | 28 | 0 (0.0) |
| Other drugs | 23 | 0 (0.0) | 11 | 0 (0.0) | 34 | 0 (0.0) |
| p value | | 0.218 | | 0.719 | | 0.119 |
| 2019 | | | | | | |
| Heroin | 58 | 2 (3.5) | 31 | 0 (0.0) | 90 | 2 (2.2) |
| Methamphetamine | 189 | 9 (4.8) | 85 | 0 (0.0) | 277 | 10 (3.6) |
| Other opioids | 101 | 0 (0.0) | 31 | 0 (0.0) | 134 | 0 (0.0) |
| PIEDs | 24 | 0 (0.0) | 3 | 0 (0.0) | 27 | 0 (0.0) |
| Other drugs | 29 | 0 (0.0) | 7 | 0 (0.0) | 36 | 0 (0.0) |
| p value | | 0.131 | | -- | | 0.138 |
| 2020 | | | | | | |
| Heroin | 45 | 0 (0.0) | 17 | 0 (0.0) | 62 | 0 (0.0) |
| Methamphetamine | 110 | 11 (10.0) | 44 | 1 (2.3) | 155 | 12 (7.7) |
| Other opioids | 85 | 2 (2.4) | 32 | 0 (0.0) | 118 | 2 (1.7) |
| PIEDs | 29 | 0 (0.0) | 1 | 0 (0.0) | 30 | 0 (0.0) |
| Other drugs | 14 | 0 (0.0) | 8 | 0 (0.0) | 22 | 0 (0.0) |
| p value | | 0.030 | | 1.000 | | 0.025 |
| 2021 | | | | | | |
| Heroin | 48 | 0 (0.0) | 22 | 0 (0.0) | 71 | 0 (0.0) |
| Methamphetamine | 108 | 5 (4.6) | 82 | 0 (0.0) | 193 | 6 (3.1) |
| Other opioids | 94 | 2 (2.1) | 33 | 0 (0.0) | 129 | 2 (1.6) |
| PIEDs | 34 | 0 (0.0) | 0 | 0 (0.0) | 34 | 0 (0.0) |
| Other drugs | 16 | 0 (0.0) | 5 | 0 (0.0) | 21 | 0 (0.0) |
| p value | | 0.496 | | -- | | 0.585 |
| 2022 | | | | | | |
| Heroin | 40 | 0 (0.0) | 21 | 0 (0.0) | 61 | 0 (0.0) |
| Methamphetamine | 87 | 5 (5.8) | 46 | 0 (0.0) | 140 | 5 (3.6) |
| Other opioids | 58 | 1 (1.7) | 15 | 0 (0.0) | 73 | 1 (1.4) |
| PIEDs | 30 | 0 (0.0) | 0 | 0 (0.0) | 30 | 0 (0.0) |
| Other drugs | 11 | 1 (9.1) | 6 | 0 (0.0) | 17 | 1 (5.9) |
| p value | | 0.200 | | -- | | 0.329 |

Table 5.2.5 HIV antibody prevalence by years of injection, gender and survey year

| Queensland Years of injection | Male | | Female | | Total | |
|----------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | | | | | | |
| <3 years | 28 | 1 (3.6) | 13 | 0 (0.0) | 41 | 1 (2.4) |
| 3 to 10 years | 61 | 2 (3.3) | 34 | 0 (0.0) | 95 | 2 (2.1) |
| 11+ years | 348 | 3 (0.9) | 142 | 2 (1.4) | 491 | 5 (1.0) |
| p value | | 0.126 | | 1.000 | | 0.268 |
| 2019 | | | | | | |
| <3 years | 36 | 2 (5.6) | 14 | 0 (0.0) | 50 | 2 (4.0) |
| 3 to 10 years | 59 | 2 (3.4) | 29 | 0 (0.0) | 90 | 3 (3.3) |
| 11+ years | 285 | 7 (2.5) | 110 | 0 (0.0) | 399 | 7 (1.8) |
| p value | | 0.369 | | -- | | 0.302 |
| 2020 | | | | | | |
| <3 years | 24 | 0 (0.0) | 8 | 0 (0.0) | 32 | 0 (0.0) |
| 3 to 10 years | 42 | 5 (11.9) | 18 | 1 (5.6) | 60 | 6 (10.0) |
| 11+ years | 213 | 8 (3.8) | 76 | 0 (0.0) | 291 | 8 (2.8) |
| p value | | 0.053 | | 0.255 | | 0.025 |
| 2021 | | | | | | |
| <3 years | 20 | 0 (0.0) | 6 | 0 (0.0) | 27 | 0 (0.0) |
| 3 to 10 years | 47 | 2 (4.3) | 20 | 0 (0.0) | 67 | 2 (3.0) |
| 11+ years | 228 | 5 (2.2) | 112 | 0 (0.0) | 345 | 6 (1.7) |
| p value | | 0.600 | | -- | | 0.773 |
| 2022 | | | | | | |
| <3 years | 13 | 1 (7.7) | 5 | 0 (0.0) | 21 | 1 (4.8) |
| 3 to 10 years | 28 | 0 (0.0) | 21 | 0 (0.0) | 50 | 0 (0.0) |
| 11+ years | 172 | 6 (3.5) | 62 | 0 (0.0) | 237 | 6 (2.5) |
| p value | | 0.278 | | -- | | 0.328 |

Table 5.2.6 HIV antibody prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| Queensland Receptively shared syringe last month | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | | | | | | |
| No receptive sharing | 337 | 4 (1.2) | 147 | 2 (1.4) | 485 | 6 (1.2) |
| Receptive sharing | 62 | 2 (3.2) | 25 | 0 (0.0) | 87 | 2 (2.3) |
| p value | | 0.236 | | 1.000 | | 0.350 |
| 2019 | | | | | | |
| No receptive sharing | 321 | 9 (2.8) | 130 | 0 (0.0) | 456 | 10 (2.2) |
| Receptive sharing | 40 | 1 (2.5) | 15 | 0 (0.0) | 56 | 1 (1.8) |
| p value | | 1.000 | | -- | | 1.000 |
| 2020 | | | | | | |
| No receptive sharing | 217 | 11 (5.1) | 71 | 1 (1.4) | 290 | 12 (4.1) |
| Receptive sharing | 32 | 1 (3.1) | 15 | 0 (0.0) | 47 | 1 (2.1) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2021 | | | | | | |
| No receptive sharing | 225 | 7 (3.1) | 95 | 0 (0.0) | 324 | 8 (2.5) |
| Receptive sharing | 32 | 0 (0.0) | 19 | 0 (0.0) | 51 | 0 (0.0) |
| p value | | 0.602 | | -- | | 0.605 |
| 2022 | | | | | | |
| No receptive sharing | 180 | 4 (2.2) | 59 | 0 (0.0) | 245 | 4 (1.6) |
| Receptive sharing | 25 | 1 (4.0) | 16 | 0 (0.0) | 41 | 1 (2.4) |
| p value | | 0.482 | | -- | | 0.541 |

Table 5.2.7 HIV antibody prevalence by frequency of drug injection last month, gender and survey year

| Queensland Frequency of injection | Male | | Female | | Total | |
|--------------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | | | | | | |
| Less than daily | 176 | 2 (1.1) | 72 | 2 (2.8) | 248 | 4 (1.6) |
| Daily or more | 226 | 4 (1.8) | 102 | 0 (0.0) | 329 | 4 (1.2) |
| Not last month | 27 | 0 (0.0) | 13 | 0 (0.0) | 40 | 0 (0.0) |
| p value | | 0.797 | | 0.282 | | 0.843 |
| 2019 | | | | | | |
| Less than daily | 163 | 7 (4.3) | 53 | 0 (0.0) | 218 | 7 (3.2) |
| Daily or more | 200 | 3 (1.5) | 93 | 0 (0.0) | 297 | 4 (1.4) |
| Not last month | 37 | 1 (2.7) | 10 | 0 (0.0) | 47 | 1 (2.1) |
| p value | | 0.221 | | -- | | 0.267 |
| 2020 | | | | | | |
| Less than daily | 93 | 7 (7.5) | 41 | 1 (2.4) | 136 | 8 (5.9) |
| Daily or more | 156 | 5 (3.2) | 46 | 0 (0.0) | 202 | 5 (2.5) |
| Not last month | 32 | 1 (3.1) | 14 | 0 (0.0) | 46 | 1 (2.2) |
| p value | | 0.238 | | 0.545 | | 0.235 |
| 2021 | | | | | | |
| Less than daily | 121 | 4 (3.3) | 40 | 0 (0.0) | 162 | 4 (2.5) |
| Daily or more | 136 | 3 (2.2) | 75 | 0 (0.0) | 214 | 4 (1.9) |
| Not last month | 42 | 0 (0.0) | 29 | 0 (0.0) | 73 | 0 (0.0) |
| p value | | 0.670 | | -- | | 0.498 |
| 2022 | | | | | | |
| Less than daily | 100 | 3 (3.0) | 25 | 0 (0.0) | 131 | 3 (2.3) |
| Daily or more | 105 | 2 (1.9) | 50 | 0 (0.0) | 155 | 2 (1.3) |
| Not last month | 21 | 2 (9.5) | 12 | 0 (0.0) | 34 | 2 (5.9) |
| p value | | 0.195 | | -- | | 0.214 |

Table 5.2.8 HIV antibody prevalence by imprisonment last year, gender and survey year

| Queensland Imprisonment last year | Male | | Female | | Total | |
|--------------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | | | | | | |
| No imprisonment | 365 | 5 (1.4) | 165 | 2 (1.2) | 531 | 7 (1.3) |
| Imprisonment | 55 | 1 (1.8) | 16 | 0 (0.0) | 71 | 1 (1.4) |
| p value | | 0.572 | | 1.000 | | 1.000 |
| 2019 | | | | | | |
| No imprisonment | 334 | 11 (3.3) | 133 | 0 (0.0) | 472 | 12 (2.5) |
| Imprisonment | 48 | 0 (0.0) | 15 | 0 (0.0) | 65 | 0 (0.0) |
| p value | | 0.372 | | -- | | 0.377 |
| 2020 | | | | | | |
| No imprisonment | 240 | 10 (4.2) | 91 | 1 (1.1) | 333 | 11 (3.3) |
| Imprisonment | 33 | 3 (9.1) | 6 | 0 (0.0) | 39 | 3 (7.7) |
| p value | | 0.198 | | 1.000 | | 0.172 |
| 2021 | | | | | | |
| No imprisonment | 252 | 7 (2.8) | 126 | 0 (0.0) | 383 | 8 (2.1) |
| Imprisonment | 36 | 0 (0.0) | 9 | 0 (0.0) | 45 | 0 (0.0) |
| p value | | 0.602 | | -- | | 1.000 |
| 2022 | | | | | | |
| No imprisonment | 197 | 7 (3.6) | 82 | 0 (0.0) | 285 | 7 (2.5) |
| Imprisonment | 22 | 0 (0.0) | 7 | 0 (0.0) | 29 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |

Table 5.2.9 HIV antibody prevalence by condom use at last sex with regular partner, gender and survey year

| Queensland Condom use at last sex | Male | | Female | | Total | |
|--------------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | | | | | | |
| No condom use | 145 | 3 (2.1) | 84 | 0 (0.0) | 229 | 3 (1.3) |
| Condom use | 35 | 1 (2.9) | 11 | 0 (0.0) | 47 | 1 (2.1) |
| p value | | 0.582 | | -- | | 0.528 |
| 2019 | | | | | | |
| No condom use | 128 | 6 (4.7) | 65 | 0 (0.0) | 196 | 6 (3.1) |
| Condom use | 32 | 0 (0.0) | 14 | 0 (0.0) | 47 | 0 (0.0) |
| p value | | 0.601 | | -- | | 0.600 |
| 2020 | | | | | | |
| No condom use | 88 | 6 (6.8) | 43 | 0 (0.0) | 131 | 6 (4.6) |
| Condom use | 21 | 1 (4.8) | 5 | 0 (0.0) | 26 | 1 (3.9) |
| p value | | 1.000 | | -- | | 1.000 |
| 2021 | | | | | | |
| No condom use | 106 | 1 (0.9) | 54 | 0 (0.0) | 160 | 1 (0.6) |
| Condom use | 26 | 0 (0.0) | 15 | 0 (0.0) | 42 | 1 (2.4) |
| p value | | 1.000 | | -- | | 0.373 |
| 2022 | | | | | | |
| No condom use | 73 | 1 (1.4) | 41 | 0 (0.0) | 115 | 1 (0.9) |
| Condom use | 18 | 1 (5.6) | 5 | 0 (0.0) | 25 | 1 (4.0) |
| p value | | 0.358 | | -- | | 0.326 |

Table 5.2.10 HIV antibody prevalence by sex work last month, gender and survey year

| Queensland Sex work last month | Male | | Female | | Total | |
|-----------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | | | | | | |
| No sex work | 409 | 6 (1.5) | 168 | 2 (1.2) | 577 | 8 (1.4) |
| Sex work | 14 | 0 (0.0) | 14 | 0 (0.0) | 29 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2019 | | | | | | |
| No sex work | 373 | 9 (2.4) | 140 | 0 (0.0) | 519 | 10 (1.9) |
| Sex work | 13 | 2 (15.4) | 7 | 0 (0.0) | 21 | 2 (9.5) |
| p value | | 0.049 | | -- | | 0.075 |
| 2020 | | | | | | |
| No sex work | 268 | 10 (3.7) | 91 | 1 (1.1) | 361 | 11 (3.1) |
| Sex work | 9 | 3 (33.3) | 4 | 0 (0.0) | 13 | 3 (23.1) |
| p value | | 0.006 | | 1.000 | | 0.010 |
| 2021 | | | | | | |
| No sex work | 285 | 7 (2.5) | 131 | 0 (0.0) | 420 | 7 (1.7) |
| Sex work | 8 | 0 (0.0) | 5 | 0 (0.0) | 14 | 1 (7.1) |
| p value | | 1.000 | | -- | | 0.232 |
| 2022 | | | | | | |
| No sex work | 209 | 7 (3.4) | 85 | 0 (0.0) | 301 | 7 (2.3) |
| Sex work | 5 | 0 (0.0) | 4 | 0 (0.0) | 9 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |

Table 5.2.11 HIV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Queensland Aboriginal and Torres Strait Islander origin | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Non Indigenous | 362 | 3 (0.8) | 143 | 2 (1.4) | 506 | 5 (1.0) |
| Indigenous | 71 | 3 (4.2) | 44 | 0 (0.0) | 115 | 3 (2.6) |
| p value | | 0.058 | | 1.000 | | 0.170 |
| 2019 | | | | | | |
| Non Indigenous | 314 | 8 (2.6) | 129 | 0 (0.0) | 448 | 9 (2.0) |
| Indigenous | 83 | 3 (3.6) | 27 | 0 (0.0) | 112 | 3 (2.7) |
| p value | | 0.705 | | -- | | 0.714 |
| 2020 | | | | | | |
| Non Indigenous | 231 | 9 (3.9) | 82 | 1 (1.2) | 314 | 10 (3.2) |
| Indigenous | 45 | 3 (6.7) | 19 | 0 (0.0) | 64 | 3 (4.7) |
| p value | | 0.421 | | 1.000 | | 0.468 |
| 2021 | | | | | | |
| Non Indigenous | 247 | 6 (2.4) | 114 | 0 (0.0) | 364 | 6 (1.7) |
| Indigenous | 49 | 1 (2.0) | 26 | 0 (0.0) | 78 | 2 (2.6) |
| p value | | 1.000 | | -- | | 0.635 |
| 2022 | | | | | | |
| Non Indigenous | 186 | 7 (3.8) | 73 | 0 (0.0) | 265 | 7 (2.6) |
| Indigenous | 37 | 0 (0.0) | 17 | 0 (0.0) | 55 | 0 (0.0) |
| p value | | 0.604 | | -- | | 0.608 |

Table 5.2.12 HIV antibody prevalence by main language spoken at home by parents, gender and survey year

| Queensland Main language spoken at home by parents | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| English speaking | 425 | 6 (1.4) | 183 | 0 (0.0) | 608 | 6 (1.0) |
| Non-English speaking | 11 | 0 (0.0) | 6 | 2 (33.3) | 18 | 2 (11.1) |
| p value | | 1.000 | | 0.001 | | 0.020 |
| 2019 | | | | | | |
| English speaking | 383 | 11 (2.9) | 153 | 0 (0.0) | 542 | 12 (2.2) |
| Non-English speaking | 17 | 0 (0.0) | 4 | 0 (0.0) | 22 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2020 | | | | | | |
| English speaking | 276 | 13 (4.7) | 101 | 1 (1.0) | 379 | 14 (3.7) |
| Non-English speaking | 8 | 0 (0.0) | 1 | 0 (0.0) | 9 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2021 | | | | | | |
| English speaking | 292 | 7 (2.4) | 143 | 0 (0.0) | 439 | 8 (1.8) |
| Non-English speaking | 8 | 0 (0.0) | 1 | 0 (0.0) | 11 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2022 | | | | | | |
| English speaking | 216 | 7 (3.2) | 90 | 0 (0.0) | 313 | 7 (2.2) |
| Non-English speaking | 11 | 0 (0.0) | 1 | 0 (0.0) | 12 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |

Table 5.2.13 HIV antibody prevalence by region/country of birth, gender and survey year

| Queensland | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Region/country of birth | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Australia | 391 | 5 (1.3) | 175 | 2 (1.1) | 566 | 7 (1.2) |
| Other Oceania | 16 | 0 (0.0) | 3 | 0 (0.0) | 20 | 0 (0.0) |
| Asia | 0 | 0 (0.0) | 0 | 0 (0.0) | 0 | 0 (0.0) |
| UK & Ireland | 17 | 1 (5.9) | 7 | 0 (0.0) | 24 | 1 (4.2) |
| Other | 11 | 0 (0.0) | 4 | 0 (0.0) | 15 | 0 (15.0) |
| p value | | 0.475 | | 1.000 | | 0.550 |
| 2019 | | | | | | |
| Australia | 366 | 11 (3.0) | 145 | 0 (0.0) | 517 | 12 (2.3) |
| Other Oceania | 9 | 0 (0.0) | 2 | 0 (0.0) | 12 | 0 (0.0) |
| Asia | 2 | 0 (0.0) | 2 | 0 (0.0) | 4 | 0 (0.0) |
| UK & Ireland | 8 | 0 (0.0) | 3 | 0 (0.0) | 11 | 0 (0.0) |
| Other | 12 | 0 (0.0) | 5 | 0 (0.0) | 17 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2020 | | | | | | |
| Australia | 252 | 12 (4.8) | 95 | 1 (1.1) | 349 | 13 (3.7) |
| Other Oceania | 11 | 0 (0.0) | 4 | 0 (0.0) | 15 | 0 (0.0) |
| Asia | 1 | 0 (0.0) | 0 | 0 (0.0) | 1 | 0 (0.0) |
| UK & Ireland | 9 | 1 (11.1) | 2 | 0 (0.0) | 11 | 1 (9.1) |
| Other | 7 | 0 (0.0) | 1 | 0 (0.0) | 8 | 0 (0.0) |
| p value | | 0.607 | | 1.000 | | 0.584 |
| 2021 | | | | | | |
| Australia | 273 | 6 (2.2) | 127 | 0 (0.0) | 405 | 7 (1.7) |
| Other Oceania | 7 | 0 (0.0) | 3 | 0 (0.0) | 11 | 0 (0.0) |
| Asia | 2 | 0 (0.0) | 1 | 0 (0.0) | 3 | 0 (0.0) |
| UK & Ireland | 8 | 0 (0.0) | 3 | 0 (0.0) | 11 | 0 (0.0) |
| Other | 10 | 1 (10.0) | 4 | 0 (0.0) | 14 | 1 (7.1) |
| p value | | 0.487 | | -- | | 0.524 |
| 2022 | | | | | | |
| Australia | 200 | 7 (3.5) | 83 | 0 (0.0) | 290 | 7 (2.4) |
| Other Oceania | 7 | 0 (0.0) | 4 | 0 (0.0) | 11 | 0 (0.0) |
| Asia | 0 | 0 (0.0) | 1 | 0 (0.0) | 1 | 0 (0.0) |
| UK & Ireland | 7 | 0 (0.0) | 1 | 0 (0.0) | 8 | 0 (0.0) |
| Other | 6 | 0 (0.0) | 2 | 0 (0.0) | 8 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |

HCV antibody prevalence

Table 5.3.1 HCV antibody prevalence by gender and survey year

| Queensland Survey year | Male | | Female | | Total | |
|---------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | 436 | 182 (42) | 188 | 84 (45) | 625 | 266 (43) |
| 2019 | 401 | 175 (44) | 156 | 57 (37) | 564 | 234 (41) |
| 2020 | 283 | 104 (37) | 102 | 42 (41) | 387 | 146 (38) |
| 2021 | 304 | 120 (39) | 145 | 51 (35) | 455 | 172 (38) |
| 2022 | 227 | 53 (23) | 91 | 30 (33) | 325 | 84 (26) |
| X ² p trend | | <0.001 | | 0.053 | | <0.001 |

Table 5.3.2 HCV antibody prevalence by sexual identity, gender and survey year

| Queensland Sexual identity | Male | | Female | | Total | |
|-------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heterosexual | 364 | 160 (44) | 136 | 61 (45) | 500 | 221 (44) |
| Bisexual | 25 | 8 (32) | 36 | 17 (47) | 61 | 25 (41) |
| Homosexual | 21 | 7 (33) | 3 | 1 (33) | 25 | 8 (32) |
| p value | | 0.340 | | 0.939 | | 0.450 |
| 2019 | | | | | | |
| Heterosexual | 322 | 142 (44) | 94 | 34 (36) | 418 | 178 (43) |
| Bisexual | 24 | 11 (46) | 36 | 14 (39) | 62 | 25 (40) |
| Homosexual | 21 | 6 (29) | 5 | 3 (60) | 27 | 9 (33) |
| p value | | 0.414 | | 0.531 | | 0.620 |
| 2020 | | | | | | |
| Heterosexual | 244 | 92 (38) | 69 | 30 (43) | 313 | 122 (39) |
| Bisexual | 13 | 2 (15) | 21 | 10 (48) | 36 | 12 (33) |
| Homosexual | 9 | 1 (11) | 3 | 0 (0) | 12 | 1 (8) |
| p value | | 0.027 | | 0.559 | | 0.064 |
| 2021 | | | | | | |
| Heterosexual | 267 | 111 (42) | 110 | 39 (35) | 379 | 151 (40) |
| Bisexual | 18 | 5 (28) | 24 | 7 (29) | 42 | 12 (29) |
| Homosexual | 15 | 3 (20) | 6 | 4 (67) | 24 | 7 (29) |
| p value | | 0.238 | | 0.466 | | 0.333 |
| 2022 | | | | | | |
| Heterosexual | 192 | 48 (25) | 59 | 20 (34) | 252 | 69 (27) |
| Bisexual | 13 | 3 (23) | 19 | 5 (26) | 34 | 8 (24) |
| Homosexual | 15 | 2 (13) | 5 | 1 (20) | 20 | 3 (15) |
| p value | | 0.891 | | 0.303 | | 0.144 |

Table 5.3.3 HCV antibody prevalence by age group, gender and survey year

| Queensland Age group | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <25 years | 20 | 2 (10) | 15 | 3 (20) | 35 | 5 (14) |
| 25-34 years | 76 | 23 (30) | 45 | 18 (40) | 122 | 41 (34) |
| 35-44 years | 158 | 59 (37) | 67 | 34 (51) | 225 | 93 (41) |
| 45+ years | 181 | 97 (54) | 61 | 29 (48) | 242 | 126 (52) |
| p value | | <0.001 | | 0.153 | | <0.001 |
| 2019 | | | | | | |
| <25 years | 18 | 5 (28) | 12 | 2 (17) | 30 | 7 (23) |
| 25-34 years | 62 | 16 (26) | 34 | 11 (32) | 99 | 28 (28) |
| 35-44 years | 147 | 66 (45) | 67 | 26 (39) | 217 | 93 (43) |
| 45+ years | 173 | 88 (51) | 43 | 18 (42) | 217 | 106 (49) |
| p value | | 0.003 | | 0.412 | | 0.001 |
| 2020 | | | | | | |
| <25 years | 11 | 3 (27) | 7 | 2 (29) | 18 | 5 (28) |
| 25-34 years | 42 | 3 (7) | 20 | 9 (45) | 62 | 12 (19) |
| 35-44 years | 103 | 42 (41) | 40 | 17 (43) | 144 | 59 (41) |
| 45+ years | 127 | 56 (44) | 35 | 14 (40) | 163 | 70 (43) |
| p value | | <0.001 | | 0.921 | | 0.005 |
| 2021 | | | | | | |
| <25 years | 14 | 2 (14) | 5 | 1 (20) | 20 | 3 (15) |
| 25-34 years | 49 | 14 (29) | 30 | 12 (40) | 80 | 26 (33) |
| 35-44 years | 108 | 47 (44) | 49 | 17 (35) | 161 | 65 (40) |
| 45+ years | 133 | 57 (43) | 61 | 21 (34) | 194 | 78 (40) |
| p value | | 0.055 | | 0.865 | | 0.092 |
| 2022 | | | | | | |
| <25 years | 9 | 1 (11) | 3 | 1 (33) | 15 | 2 (13) |
| 25-34 years | 36 | 5 (14) | 17 | 4 (24) | 55 | 9 (16) |
| 35-44 years | 71 | 16 (23) | 39 | 12 (31) | 111 | 29 (26) |
| 45+ years | 109 | 31 (28) | 31 | 12 (39) | 141 | 43 (31) |
| p value | | 0.285 | | 0.756 | | 0.154 |

Table 5.3.4 HCV antibody prevalence by years since first injection, gender and survey year

| Queensland Years of injection | Male | | Female | | Total | |
|----------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| <3 years | 28 | 1 (4) | 13 | 1 (8) | 41 | 2 (5) |
| 3 to 10 years | 60 | 11 (18) | 33 | 8 (24) | 93 | 19 (20) |
| 11+ years | 333 | 164 (49) | 137 | 73 (53) | 470 | 237 (50) |
| p value | | <0.001 | | <0.001 | | <0.001 |
| 2019 | | | | | | |
| <3 years | 36 | 3 (8) | 14 | 0 (0) | 50 | 3 (6) |
| 3 to 10 years | 59 | 12 (20) | 29 | 7 (24) | 90 | 19 (21) |
| 11+ years | 285 | 151 (53) | 109 | 50 (46) | 398 | 203 (51) |
| p value | | <0.001 | | <0.001 | | <0.001 |
| 2020 | | | | | | |
| <3 years | 24 | 3 (13) | 8 | 0 (0) | 32 | 3 (9) |
| 3 to 10 years | 42 | 2 (5) | 18 | 5 (28) | 60 | 7 (12) |
| 11+ years | 212 | 96 (45) | 76 | 37 (49) | 290 | 133 (46) |
| p value | | <0.001 | | 0.008 | | <0.001 |
| 2021 | | | | | | |
| <3 years | 20 | 0 (0) | 6 | 0 (0) | 27 | 0 (0) |
| 3 to 10 years | 47 | 4 (9) | 20 | 4 (20) | 67 | 8 (12) |
| 11+ years | 228 | 111 (49) | 112 | 45 (40) | 345 | 157 (46) |
| p value | | <0.001 | | 0.043 | | <0.001 |
| 2022 | | | | | | |
| <3 years | 13 | 0 (0) | 5 | 0 (0) | 21 | 0 (0) |
| 3 to 10 years | 28 | 5 (18) | 21 | 3 (14) | 50 | 8 (16) |
| 11+ years | 172 | 44 (26) | 62 | 26 (42) | 237 | 71 (30) |
| p value | | 0.070 | | 0.018 | | 0.001 |

Table 5.3.5 HCV antibody prevalence by re-use of someone else's used needle and syringe last month, years since first injection and survey year

| Queensland Receptively shared syringe last month | <3 years | | 3+ years | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| No receptive sharing | 30 | 1 (3) | 441 | 189 (43) | 483 | 194 (40) |
| Receptive sharing | 7 | 1 (14) | 75 | 45 (60) | 87 | 50 (57) |
| p value | | 0.347 | | 0.006 | | 0.003 |
| 2019 | | | | | | |
| No receptive sharing | 41 | 2 (5) | 401 | 175 (44) | 455 | 183 (40) |
| Receptive sharing | 3 | 0 (0) | 48 | 31 (65) | 56 | 33 (59) |
| p value | | 1.000 | | 0.006 | | 0.007 |
| 2020 | | | | | | |
| No receptive sharing | 22 | 2 (9) | 265 | 104 (39) | 289 | 107 (37) |
| Receptive sharing | 3 | 0 (0) | 43 | 23 (53) | 47 | 24 (51) |
| p value | | 0.770 | | 0.078 | | 0.067 |
| 2021 | | | | | | |
| No receptive sharing | 19 | 0 (0) | 295 | 113 (38) | 324 | 118 (36) |
| Receptive sharing | 1 | 0 (0) | 49 | 26 (53) | 51 | 26 (51) |
| p value | | -- | | 0.051 | | 0.047 |
| 2022 | | | | | | |
| No receptive sharing | 17 | 0 (0) | 217 | 59 (27) | 245 | 63 (26) |
| Receptive sharing | 0 | 0 (0) | 36 | 11 (31) | 41 | 12 (29) |
| p value | | -- | | 0.676 | | 0.632 |

Table 5.3.6 HCV antibody prevalence by last drug injected, years since first injection and survey year

| Queensland Last drug injected | <3 years | | 3+ years | | Total | |
|----------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heroin | 3 | 1 (33) | 74 | 36 (49) | 80 | 40 (50) |
| Methamphetamine | 25 | 1 (4) | 262 | 99 (38) | 297 | 102 (34) |
| Other opioids | 1 | 0 (0) | 172 | 99 (58) | 177 | 101 (57) |
| PIEDs | 7 | 0 (0) | 19 | 0 (0) | 27 | 0 (0) |
| Other drugs | 4 | 0 (0) | 27 | 16 (59) | 33 | 17 (52) |
| p value | | 0.218 | | <0.001 | | <0.001 |
| 2019 | | | | | | |
| Heroin | 1 | 0 (0) | 87 | 55 (63) | 90 | 57 (63) |
| Methamphetamine | 31 | 1 (3) | 230 | 71 (31) | 275 | 75 (27) |
| Other opioids | 4 | 1 (25) | 125 | 78 (62) | 133 | 82 (62) |
| PIEDs | 11 | 0 (0) | 15 | 0 (0) | 27 | 0 (0) |
| Other drugs | 3 | 1 (33) | 29 | 16 (55) | 36 | 18 (50) |
| p value | | 0.102 | | <0.001 | | <0.001 |
| 2020 | | | | | | |
| Heroin | 0 | 0 (0) | 61 | 37 (61) | 61 | 37 (61) |
| Methamphetamine | 14 | 3 (21) | 138 | 29 (21) | 155 | 34 (22) |
| Other opioids | 3 | 0 (0) | 112 | 57 (51) | 116 | 58 (50) |
| PIEDs | 11 | 0 (0) | 18 | 1 (6) | 30 | 1 (3) |
| Other drugs | 3 | 0 (0) | 17 | 13 (76) | 20 | 13 (65) |
| p value | | 0.258 | | <0.001 | | <0.001 |
| 2021 | | | | | | |
| Heroin | 0 | 0 (0) | 71 | 41 (58) | 71 | 41 (58) |
| Methamphetamine | 17 | 0 (0) | 168 | 46 (27) | 190 | 49 (26) |
| Other opioids | 2 | 0 (0) | 120 | 64 (53) | 127 | 65 (51) |
| PIEDs | 6 | 0 (0) | 27 | 3 (11) | 34 | 3 (9) |
| Other drugs | 1 | 0 (0) | 15 | 7 (47) | 20 | 9 (45) |
| p value | | -- | | <0.001 | | <0.001 |
| 2022 | | | | | | |
| Heroin | 1 | 0 (0) | 57 | 26 (46) | 60 | 27 (45) |
| Methamphetamine | 8 | 0 (0) | 121 | 21 (17) | 137 | 24 (18) |
| Other opioids | 2 | 0 (0) | 67 | 25 (37) | 73 | 26 (36) |
| PIEDs | 8 | 0 (0) | 21 | 1 (5) | 30 | 1 (3) |
| Other drugs | 1 | 0 (0) | 14 | 6 (43) | 16 | 6 (38) |
| p value | | -- | | <0.001 | | <0.001 |

Table 5.3.7 HCV antibody prevalence by frequency of drug injection last month, years since first injection and survey year

| Queensland | <3 years | | 3+ years | | Total | |
|-----------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Frequency of injection last month | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Less than daily | 22 | 2 (9) | 217 | 91 (42) | 246 | 96 (39) |
| Daily or more | 15 | 0 (0) | 304 | 146 (48) | 329 | 151 (46) |
| Not last month | 3 | 0 (0) | 34 | 13 (38) | 40 | 13 (33) |
| p value | | 0.577 | | 0.276 | | 0.111 |
| 2019 | | | | | | |
| Less than daily | 25 | 1 (4) | 185 | 74 (40) | 217 | 77 (35) |
| Daily or more | 19 | 1 (5) | 267 | 133 (50) | 297 | 140 (47) |
| Not last month | 6 | 1 (17) | 34 | 13 (38) | 47 | 15 (32) |
| p value | | 0.491 | | 0.083 | | 0.012 |
| 2020 | | | | | | |
| Less than daily | 16 | 0 (0) | 119 | 41 (34) | 136 | 41 (30) |
| Daily or more | 9 | 2 (22) | 190 | 86 (45) | 201 | 90 (45) |
| Not last month | 7 | 1 (14) | 37 | 10 (27) | 46 | 12 (26) |
| p value | | 0.113 | | 0.043 | | 0.006 |
| 2021 | | | | | | |
| Less than daily | 15 | 0 (0) | 143 | 38 (27) | 162 | 40 (25) |
| Daily or more | 5 | 0 (0) | 202 | 102 (51) | 214 | 105 (49) |
| Not last month | 7 | 0 (0) | 61 | 22 (36) | 73 | 24 (33) |
| p value | | -- | | <0.001 | | <0.001 |
| 2022 | | | | | | |
| Less than daily | 13 | 0 (0) | 113 | 28 (25) | 131 | 30 (23) |
| Daily or more | 4 | 0 (0) | 140 | 42 (30) | 155 | 45 (29) |
| Not last month | 3 | 0 (0) | 30 | 9 (30) | 34 | 9 (26) |
| p value | | -- | | 0.631 | | 0.502 |

Table 5.3.8 HCV antibody prevalence by imprisonment last year, years since first injection and survey year

| Queensland | <3 years | | 3+ years | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Imprisonment last year | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| No imprisonment | 39 | 2 (5) | 472 | 210 (44) | 529 | 218 (41) |
| Imprisonment | 2 | 0 (0) | 65 | 35 (54) | 70 | 37 (53) |
| p value | | 1.000 | | 0.156 | | 0.064 |
| 2019 | | | | | | |
| No imprisonment | 46 | 3 (7) | 402 | 170 (42) | 471 | 181 (38) |
| Imprisonment | 3 | 0 (0) | 60 | 41 (68) | 65 | 41 (63) |
| p value | | 1.000 | | <0.001 | | <0.001 |
| 2020 | | | | | | |
| No imprisonment | 31 | 2 (6) | 297 | 112 (38) | 333 | 117 (35) |
| Imprisonment | 1 | 1 (100) | 37 | 17 (46) | 38 | 18 (47) |
| p value | | 0.094 | | 0.372 | | 0.138 |
| 2021 | | | | | | |
| No imprisonment | 25 | 0 (0) | 345 | 124 (36) | 383 | 129 (34) |
| Imprisonment | 0 | 0 (0) | 44 | 31 (70) | 45 | 32 (71) |
| p value | | -- | | <0.001 | | <0.001 |
| 2022 | | | | | | |
| No imprisonment | 17 | 0 (0) | 254 | 70 (28) | 285 | 74 (26) |
| Imprisonment | 3 | 0 (0) | 24 | 6 (25) | 29 | 7 (24) |
| p value | | -- | | 0.788 | | 0.830 |

Table 5.3.9 HCV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Queensland Aboriginal and Torres Strait Islander origin | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Non Indigenous | 362 | 145 (40) | 142 | 63 (44) | 505 | 208 (41) |
| Indigenous | 69 | 35 (51) | 44 | 21 (48) | 113 | 56 (50) |
| p value | | 0.100 | | 0.695 | | 0.104 |
| 2019 | | | | | | |
| Non Indigenous | 314 | 128 (41) | 128 | 42 (33) | 447 | 170 (38) |
| Indigenous | 83 | 45 (54) | 27 | 15 (56) | 112 | 62 (55) |
| p value | | 0.028 | | 0.026 | | 0.001 |
| 2020 | | | | | | |
| Non Indigenous | 231 | 76 (33) | 82 | 33 (40) | 314 | 109 (35) |
| Indigenous | 44 | 26 (59) | 19 | 9 (47) | 63 | 35 (56) |
| p value | | 0.001 | | 0.570 | | 0.002 |
| 2021 | | | | | | |
| Non Indigenous | 247 | 84 (34) | 114 | 37 (32) | 364 | 122 (34) |
| Indigenous | 49 | 33 (67) | 26 | 12 (46) | 78 | 45 (58) |
| p value | | <0.001 | | 0.186 | | <0.001 |
| 2022 | | | | | | |
| Non Indigenous | 186 | 40 (22) | 73 | 24 (33) | 265 | 65 (25) |
| Indigenous | 37 | 11 (30) | 17 | 6 (35) | 55 | 17 (31) |
| p value | | 0.277 | | 0.849 | | 0.324 |

Table 5.3.10 HCV antibody prevalence by main language spoken at home by parents, gender and survey year

| Queensland Main language spoken at home by parents | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| English speaking | 423 | 176 (42) | 182 | 82 (45) | 605 | 258 (43) |
| Non-English speaking | 11 | 5 (45) | 6 | 2 (33) | 18 | 7 (39) |
| p value | | 0.798 | | 0.693 | | 0.751 |
| 2019 | | | | | | |
| English speaking | 383 | 166 (43) | 152 | 55 (36) | 541 | 223 (41) |
| Non-English speaking | 17 | 8 (47) | 4 | 2 (50) | 22 | 10 (45) |
| p value | | 0.762 | | 0.623 | | 0.693 |
| 2020 | | | | | | |
| English speaking | 275 | 99 (36) | 101 | 41 (41) | 378 | 140 (37) |
| Non-English speaking | 8 | 5 (63) | 1 | 1 (100) | 9 | 6 (67) |
| p value | | 0.148 | | 0.412 | | 0.087 |
| 2021 | | | | | | |
| English speaking | 292 | 118 (40) | 143 | 49 (34) | 439 | 167 (38) |
| Non-English speaking | 8 | 2 (25) | 1 | 1 (100) | 11 | 4 (36) |
| p value | | 0.483 | | 0.347 | | 1.000 |
| 2022 | | | | | | |
| English speaking | 216 | 48 (22) | 90 | 30 (33) | 313 | 79 (25) |
| Non-English speaking | 11 | 5 (45) | 1 | 0 (0) | 12 | 5 (42) |
| p value | | 0.076 | | 1.000 | | 0.202 |

Table 5.3.11 HCV antibody prevalence by region/country of birth, gender and survey year

| Queensland | | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | |
| 2018 | | | | | | | |
| Australia | 389 | 164 (42) | 174 | 80 (46) | 563 | 244 (43) | |
| Other Oceania | 16 | 6 (38) | 3 | 1 (33) | 20 | 7 (35) | |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| UK & Ireland | 17 | 9 (53) | 7 | 2 (29) | 24 | 11 (46) | |
| Other | 11 | 2 (18) | 4 | 1 (25) | 15 | 3 (20) | |
| p value | | 0.319 | | 0.694 | | 0.285 | |
| 2019 | | | | | | | |
| Australia | 366 | 163 (45) | 144 | 52 (36) | 516 | 217 (42) | |
| Other Oceania | 9 | 5 (56) | 2 | 1 (50) | 12 | 6 (50) | |
| Asia | 2 | 0 (0) | 2 | 0 (0) | 4 | 0 (0) | |
| UK & Ireland | 8 | 3 (38) | 3 | 2 (67) | 11 | 5 (45) | |
| Other | 12 | 2 (17) | 5 | 2 (40) | 17 | 4 (24) | |
| p value | | 0.213 | | 0.728 | | 0.246 | |
| 2020 | | | | | | | |
| Australia | 251 | 94 (37) | 95 | 41 (43) | 348 | 135 (39) | |
| Other Oceania | 11 | 4 (36) | 4 | 1 (25) | 15 | 5 (33) | |
| Asia | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) | |
| UK & Ireland | 9 | 1 (11) | 2 | 0 (0) | 11 | 1 (9) | |
| Other | 7 | 4 (57) | 1 | 0 (0) | 8 | 4 (50) | |
| p value | | 0.347 | | 0.586 | | 0.230 | |
| 2021 | | | | | | | |
| Australia | 273 | 108 (40) | 127 | 41 (32) | 405 | 150 (37) | |
| Other Oceania | 7 | 3 (43) | 3 | 3 (100) | 11 | 6 (55) | |
| Asia | 2 | 0 (0) | 1 | 1 (100) | 3 | 1 (33) | |
| UK & Ireland | 8 | 3 (38) | 3 | 1 (33) | 11 | 4 (36) | |
| Other | 10 | 5 (50) | 4 | 2 (50) | 14 | 7 (50) | |
| p value | | 0.857 | | 0.049 | | 0.637 | |
| 2022 | | | | | | | |
| Australia | 200 | 44 (22) | 83 | 29 (35) | 290 | 74 (26) | |
| Other Oceania | 7 | 2 (29) | 4 | 1 (25) | 11 | 3 (27) | |
| Asia | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) | |
| UK & Ireland | 7 | 1 (14) | 1 | 0 (0) | 8 | 1 (13) | |
| Other | 6 | 1 (17) | 2 | 0 (0) | 8 | 1 (13) | |
| p value | | 0.953 | | 0.919 | | 0.894 | |

HCV RNA prevalence

Table 5.4.1 HCV RNA prevalence by gender and survey year

| Queensland Survey year | Male | | Female | | Total | |
|---------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | 196 | 43 (22) | 84 | 16 (19) | 280 | 59 (21) |
| 2019* | 371 | 62 (17) | 144 | 16 (11) | 522 | 79 (15) |
| 2020 | 267 | 41 (15) | 101 | 10 (10) | 370 | 51 (14) |
| 2021 | 291 | 32 (11) | 143 | 15 (10) | 440 | 48 (11) |
| 2022 | 223 | 22 (10) | 91 | 9 (10) | 321 | 32 (10) |
| X ² p trend | | <0.001 | | 0.102 | | <0.001 |

* 2018, 2019: Weighted for gender and HCV antibody status

Totals include respondents where gender was reported as other or not reported

Table 5.4.2 HCV RNA prevalence by sexual identity, gender and survey year

| Queensland Sexual identity | Male | | Female | | Total | |
|-------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heterosexual | 172 | 39 (23) | 62 | 12 (19) | 234 | 51 (22) |
| Bisexual | 8 | 1 (13) | 14 | 3 (21) | 22 | 4 (18) |
| Homosexual | 6 | 0 (0) | 3 | 0 (0) | 9 | 0 (0) |
| p value | | 0.332 | | 0.680 | | 0.228 |
| 2019* | | | | | | |
| Heterosexual | 300 | 50 (17) | 86 | 9 (10) | 388 | 61 (16) |
| Bisexual | 23 | 6 (26) | 34 | 4 (12) | 58 | 10 (17) |
| Homosexual | 19 | 1 (5) | 5 | 1 (20) | 25 | 2 (8) |
| p value | | 0.203 | | 0.801 | | 0.554 |
| 2020 | | | | | | |
| Heterosexual | 230 | 34 (15) | 67 | 6 (9) | 297 | 40 (13) |
| Bisexual | 13 | 1 (8) | 22 | 4 (18) | 37 | 5 (14) |
| Homosexual | 9 | 0 (0) | 3 | 0 (0) | 12 | 0 (0) |
| p value | | 0.364 | | 0.400 | | 0.395 |
| 2021 | | | | | | |
| Heterosexual | 256 | 30 (12) | 108 | 11 (10) | 366 | 42 (11) |
| Bisexual | 17 | 2 (12) | 24 | 1 (4) | 41 | 3 (7) |
| Homosexual | 15 | 0 (0) | 6 | 2 (33) | 24 | 2 (8) |
| p value | | 0.372 | | 0.106 | | 0.661 |
| 2022 | | | | | | |
| Heterosexual | 187 | 19 (10) | 59 | 7 (12) | 247 | 27 (11) |
| Bisexual | 13 | 2 (15) | 19 | 1 (5) | 34 | 3 (9) |
| Homosexual | 15 | 1 (7) | 5 | 0 (0) | 20 | 1 (5) |
| p value | | 0.863 | | 0.628 | | 0.833 |

Table 5.4.3 HCV RNA prevalence by age group, gender and survey year

| Queensland Age group | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <25 years | 10 | 2 (20) | 6 | 1 (17) | 16 | 3 (19) |
| 25-34 years | 43 | 5 (12) | 23 | 3 (13) | 66 | 8 (12) |
| 35-44 years | 74 | 18 (24) | 26 | 5 (19) | 100 | 23 (23) |
| 45+ years | 69 | 18 (26) | 28 | 7 (25) | 97 | 25 (26) |
| p value | | 0.407 | | 0.612 | | 0.222 |
| 2019* | | | | | | |
| <25 years | 17 | 2 (12) | 11 | 3 (27) | 28 | 5 (18) |
| 25-34 years | 55 | 5 (9) | 29 | 2 (7) | 87 | 8 (9) |
| 35-44 years | 136 | 28 (21) | 65 | 11 (17) | 204 | 38 (19) |
| 45+ years | 162 | 28 (17) | 39 | 0 (0) | 202 | 28 (14) |
| p value | | 0.288 | | 0.019 | | 0.178 |
| 2020 | | | | | | |
| <25 years | 10 | 1 (10) | 8 | 1 (13) | 18 | 2 (11) |
| 25-34 years | 36 | 1 (3) | 20 | 5 (25) | 56 | 6 (11) |
| 35-44 years | 100 | 24 (24) | 39 | 4 (10) | 140 | 28 (20) |
| 45+ years | 121 | 15 (12) | 34 | 0 (0) | 156 | 15 (10) |
| p value | | 0.011 | | 0.030 | | 0.061 |
| 2021 | | | | | | |
| <25 years | 14 | 0 (0) | 5 | 0 (0) | 20 | 0 (0) |
| 25-34 years | 47 | 9 (19) | 29 | 3 (10) | 77 | 12 (16) |
| 35-44 years | 102 | 16 (16) | 49 | 6 (12) | 155 | 23 (15) |
| 45+ years | 128 | 7 (5) | 60 | 6 (10) | 188 | 13 (7) |
| p value | | 0.011 | | 0.858 | | 0.021 |
| 2022 | | | | | | |
| <25 years | 9 | 1 (11) | 3 | 1 (33) | 15 | 2 (13) |
| 25-34 years | 36 | 3 (8) | 17 | 2 (12) | 55 | 5 (9) |
| 35-44 years | 70 | 9 (13) | 39 | 3 (8) | 110 | 13 (12) |
| 45+ years | 106 | 9 (8) | 31 | 3 (10) | 138 | 12 (9) |
| p value | | 0.716 | | 0.470 | | 0.788 |

Table 5.4.4 HCV RNA prevalence by years since first injection, gender and survey year

| Queensland Years of injection | Male | | Female | | Total | |
|----------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <3 years | 11 | 0 (0) | 7 | 0 (0) | 18 | 0 (0) |
| 3 to 10 years | 33 | 7 (21) | 15 | 1 (7) | 48 | 8 (17) |
| 11+ years | 149 | 36 (24) | 59 | 14 (24) | 208 | 50 (24) |
| p value | | 0.225 | | 0.086 | | 0.052 |
| 2019* | | | | | | |
| <3 years | 33 | 2 (6) | 13 | 1 (8) | 46 | 3 (7) |
| 3 to 10 years | 55 | 6 (11) | 25 | 4 (16) | 82 | 10 (12) |
| 11+ years | 264 | 52 (20) | 102 | 11 (11) | 370 | 64 (17) |
| p value | | 0.055 | | 0.631 | | 0.113 |
| 2020 | | | | | | |
| <3 years | 22 | 0 (0) | 8 | 0 (0) | 30 | 0 (0) |
| 3 to 10 years | 40 | 1 (3) | 19 | 4 (21) | 59 | 5 (8) |
| 11+ years | 200 | 39 (20) | 74 | 6 (8) | 276 | 45 (16) |
| p value | | 0.003 | | 0.150 | | 0.021 |
| 2021 | | | | | | |
| <3 years | 20 | 0 (0) | 6 | 0 (0) | 27 | 0 (0) |
| 3 to 10 years | 47 | 5 (11) | 19 | 0 (0) | 66 | 5 (8) |
| 11+ years | 215 | 27 (13) | 111 | 15 (14) | 331 | 43 (13) |
| p value | | 0.235 | | 0.150 | | 0.071 |
| 2022 | | | | | | |
| <3 years | 13 | 0 (0) | 5 | 0 (0) | 21 | 0 (0) |
| 3 to 10 years | 29 | 5 (17) | 21 | 3 (14) | 51 | 8 (16) |
| 11+ years | 167 | 16 (10) | 62 | 6 (10) | 232 | 23 (10) |
| p value | | 0.255 | | 0.819 | | 0.125 |

Table 5.4.5 HCV RNA prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| Queensland Receptively shared syringe last month | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| No receptive sharing | 150 | 29 (19) | 60 | 12 (20) | 210 | 42 (20) |
| Receptive sharing | 29 | 11 (38) | 15 | 3 (20) | 45 | 14 (31) |
| p value | | 0.039 | | 0.750 | | 0.122 |
| 2019* | | | | | | |
| No receptive sharing | 296 | 50 (17) | 118 | 13 (11) | 418 | 64 (15) |
| Receptive sharing | 37 | 8 (22) | 14 | 2 (14) | 52 | 10 (19) |
| p value | | 0.501 | | 0.649 | | 0.457 |
| 2020 | | | | | | |
| No receptive sharing | 203 | 27 (13) | 69 | 6 (9) | 274 | 33 (12) |
| Receptive sharing | 29 | 12 (41) | 16 | 3 (19) | 45 | 15 (33) |
| p value | | <0.001 | | 0.239 | | <0.001 |
| 2021 | | | | | | |
| No receptive sharing | 214 | 24 (11) | 94 | 10 (11) | 312 | 35 (11) |
| Receptive sharing | 30 | 6 (20) | 19 | 1 (5) | 49 | 7 (14) |
| p value | | 0.170 | | 0.471 | | 0.534 |
| 2022 | | | | | | |
| No receptive sharing | 177 | 16 (9) | 59 | 5 (8) | 242 | 22 (9) |
| Receptive sharing | 25 | 5 (20) | 16 | 2 (13) | 41 | 7 (17) |
| p value | | 0.093 | | 0.637 | | 0.119 |

Table 5.4.6 HCV RNA prevalence by last drug injected, gender and survey year

| Queensland Last drug injected | Male | | Female | | Total | |
|----------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heroin | 22 | 6 (27) | 9 | 2 (22) | 31 | 8 (26) |
| Methamphetamine | 87 | 21 (24) | 47 | 9 (19) | 134 | 30 (22) |
| Other opioids | 58 | 10 (17) | 21 | 4 (19) | 79 | 15 (19) |
| PIEDs | 13 | 0 (0) | 0 | 0 (0) | 13 | 0 (0) |
| Other drugs | 13 | 4 (31) | 5 | 0 (0) | 18 | 4 (22) |
| p value | | 0.271 | | 0.677 | | 0.420 |
| 2019* | | | | | | |
| Heroin | 55 | 10 (18) | 28 | 2 (7) | 83 | 12 (14) |
| Methamphetamine | 175 | 28 (16) | 76 | 7 (9) | 254 | 35 (14) |
| Other opioids | 92 | 19 (21) | 30 | 3 (10) | 124 | 23 (19) |
| PIEDs | 21 | 0 (0) | 3 | 0 (0) | 24 | 0 (0) |
| Other drugs | 27 | 6 (22) | 7 | 2 (29) | 34 | 8 (24) |
| p value | | 0.208 | | 0.525 | | 0.105 |
| 2020 | | | | | | |
| Heroin | 44 | 7 (16) | 18 | 1 (6) | 62 | 8 (13) |
| Methamphetamine | 103 | 15 (15) | 43 | 3 (7) | 147 | 18 (12) |
| Other opioids | 78 | 15 (19) | 30 | 1 (3) | 109 | 16 (15) |
| PIEDs | 27 | 0 (0) | 1 | 0 (0) | 28 | 0 (0) |
| Other drugs | 11 | 3 (27) | 8 | 5 (63) | 19 | 8 (42) |
| p value | | 0.130 | | <0.001 | | 0.001 |
| 2021 | | | | | | |
| Heroin | 46 | 4 (9) | 22 | 2 (9) | 69 | 7 (10) |
| Methamphetamine | 103 | 10 (10) | 80 | 6 (8) | 186 | 16 (9) |
| Other opioids | 86 | 13 (15) | 33 | 7 (21) | 121 | 20 (17) |
| PIEDs | 34 | 2 (6) | 0 | 0 (0) | 34 | 2 (6) |
| Other drugs | 15 | 2 (13) | 5 | 0 (0) | 20 | 2 (10) |
| p value | | 0.571 | | 0.149 | | 0.208 |
| 2022 | | | | | | |
| Heroin | 38 | 4 (11) | 20 | 3 (15) | 58 | 7 (12) |
| Methamphetamine | 86 | 10 (12) | 44 | 3 (7) | 137 | 14 (10) |
| Other opioids | 57 | 7 (12) | 15 | 3 (20) | 72 | 10 (14) |
| PIEDs | 30 | 0 (0) | 0 | 0 (0) | 30 | 0 (0) |
| Other drugs | 11 | 1 (9) | 5 | 0 (0) | 16 | 1 (6) |
| p value | | 0.410 | | 0.386 | | 0.291 |

Table 5.4.7 HCV RNA prevalence by frequency of drug injection last month, gender and survey year

| Queensland Frequency of injection last month | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| Less than daily | 84 | 19 (23) | 30 | 8 (27) | 114 | 27 (24) |
| Daily or more | 95 | 21 (22) | 45 | 7 (16) | 140 | 29 (21) |
| Not last month | 14 | 2 (14) | 6 | 0 (0) | 20 | 2 (10) |
| p value | | 0.753 | | 0.181 | | 0.354 |
| 2019* | | | | | | |
| Less than daily | 152 | 21 (14) | 46 | 4 (9) | 200 | 26 (13) |
| Daily or more | 182 | 38 (21) | 88 | 11 (13) | 273 | 48 (18) |
| Not last month | 36 | 4 (11) | 10 | 0 (0) | 46 | 4 (9) |
| p value | | 0.142 | | 0.469 | | 0.162 |
| 2020 | | | | | | |
| Less than daily | 88 | 12 (14) | 40 | 2 (5) | 130 | 14 (11) |
| Daily or more | 144 | 27 (19) | 46 | 7 (15) | 190 | 34 (18) |
| Not last month | 32 | 1 (3) | 14 | 1 (7) | 46 | 2 (4) |
| p value | | 0.074 | | 0.269 | | 0.027 |
| 2021 | | | | | | |
| Less than daily | 115 | 9 (8) | 40 | 2 (5) | 156 | 11 (7) |
| Daily or more | 129 | 21 (16) | 73 | 9 (12) | 205 | 31 (15) |
| Not last month | 42 | 1 (2) | 29 | 4 (14) | 73 | 5 (7) |
| p value | | 0.017 | | 0.392 | | 0.025 |
| 2022 | | | | | | |
| Less than daily | 100 | 6 (6) | 25 | 1 (4) | 131 | 8 (6) |
| Daily or more | 102 | 15 (15) | 50 | 6 (12) | 152 | 21 (14) |
| Not last month | 21 | 1 (5) | 12 | 2 (17) | 34 | 3 (9) |
| p value | | 0.103 | | 0.316 | | 0.100 |

Table 5.4.8 HCV RNA prevalence by imprisonment last year, gender and survey year

| Queensland Imprisonment last year | Male | | Female | | Total | |
|--------------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| No imprisonment | 162 | 35 (22) | 74 | 14 (19) | 236 | 50 (21) |
| Imprisonment | 27 | 7 (26) | 7 | 1 (14) | 35 | 8 (23) |
| p value | | 0.759 | | 0.621 | | 0.911 |
| 2019* | | | | | | |
| No imprisonment | 310 | 43 (14) | 121 | 11 (9) | 435 | 55 (13) |
| Imprisonment | 45 | 14 (31) | 14 | 3 (21) | 61 | 18 (30) |
| p value | | 0.004 | | 0.164 | | <0.001 |
| 2020 | | | | | | |
| No imprisonment | 228 | 30 (13) | 90 | 9 (10) | 320 | 39 (12) |
| Imprisonment | 30 | 9 (30) | 6 | 1 (17) | 36 | 10 (28) |
| p value | | 0.015 | | 0.605 | | 0.010 |
| 2021 | | | | | | |
| No imprisonment | 243 | 21 (9) | 124 | 13 (10) | 372 | 34 (9) |
| Imprisonment | 34 | 10 (29) | 9 | 2 (22) | 43 | 12 (28) |
| p value | | <0.001 | | 0.282 | | <0.001 |
| 2022 | | | | | | |
| No imprisonment | 194 | 16 (8) | 82 | 6 (7) | 282 | 23 (8) |
| Imprisonment | 22 | 5 (23) | 7 | 2 (29) | 29 | 7 (24) |
| p value | | 0.030 | | 0.119 | | 0.006 |

Table 5.4.9 HCV RNA prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Queensland | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| Aboriginal and Torres Strait Islander origin | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| Non Indigenous | 159 | 34 (21) | 65 | 11 (17) | 224 | 45 (20) |
| Indigenous | 33 | 9 (27) | 18 | 4 (22) | 52 | 14 (27) |
| p value | | 0.497 | | 0.496 | | 0.367 |
| 2019* | | | | | | |
| Non Indigenous | 290 | 39 (13) | 118 | 9 (8) | 412 | 48 (12) |
| Indigenous | 78 | 23 (29) | 26 | 6 (23) | 105 | 30 (29) |
| p value | | <0.001 | | 0.018 | | <0.001 |
| 2020 | | | | | | |
| Non Indigenous | 215 | 28 (13) | 81 | 8 (10) | 297 | 36 (12) |
| Indigenous | 44 | 13 (30) | 19 | 2 (11) | 63 | 15 (24) |
| p value | | 0.006 | | 0.932 | | 0.016 |
| 2021 | | | | | | |
| Non Indigenous | 236 | 18 (8) | 112 | 12 (11) | 351 | 31 (9) |
| Indigenous | 47 | 14 (30) | 26 | 3 (12) | 76 | 17 (22) |
| p value | | <0.001 | | 0.903 | | 0.001 |
| 2022 | | | | | | |
| Non Indigenous | 182 | 15 (8) | 73 | 6 (8) | 261 | 22 (8) |
| Indigenous | 37 | 6 (16) | 17 | 3 (18) | 55 | 9 (16) |
| p value | | 0.133 | | 0.363 | | 0.072 |

Table 5.4.10 HCV RNA prevalence by main language spoken at home by parents, gender and survey year

| Queensland | Male | | Female | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| Main language spoken at home by parents | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| English speaking | 188 | 42 (22) | 80 | 16 (20) | 268 | 58 (22) |
| Non-English speaking | 6 | 1 (17) | 4 | 0 (0) | 10 | 1 (10) |
| p value | | 0.661 | | 0.999 | | 0.349 |
| 2019* | | | | | | |
| English speaking | 354 | 59 (17) | 141 | 15 (11) | 501 | 75 (15) |
| Non-English speaking | 16 | 3 (19) | 3 | 1 (33) | 20 | 4 (20) |
| p value | | 0.843 | | 0.190 | | 0.521 |
| 2020 | | | | | | |
| English speaking | 259 | 39 (15) | 100 | 10 (10) | 361 | 49 (14) |
| Non-English speaking | 8 | 2 (25) | 1 | 0 (0) | 9 | 2 (22) |
| p value | | 0.442 | | 0.739 | | 0.457 |
| 2021 | | | | | | |
| English speaking | 279 | 31 (11) | 142 | 15 (11) | 425 | 46 (11) |
| Non-English speaking | 8 | 1 (13) | 1 | 0 (0) | 11 | 2 (18) |
| p value | | 0.902 | | 0.731 | | 0.441 |
| 2022 | | | | | | |
| English speaking | 212 | 19 (9) | 90 | 9 (10) | 309 | 29 (9) |
| Non-English speaking | 11 | 3 (27) | 1 | 0 (0) | 12 | 3 (25) |
| p value | | 0.082 | | 1.000 | | 0.106 |

Table 5.4.11 HCV RNA prevalence by region/country of birth, gender and survey year

| Queensland | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Australia | 180 | 40 (22) | 75 | 15 (20) | 255 | 55 (22) |
| Other Oceania | 6 | 1 (17) | 1 | 0 (0) | 7 | 1 (14) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 5 | 1 (20) | 4 | 1 (25) | 9 | 2 (22) |
| Other | 3 | 1 (33) | 4 | 0 (0) | 7 | 1 (14) |
| p value | | 0.952 | | 0.727 | | 0.908 |
| 2019* | | | | | | |
| Australia | 338 | 61 (18) | 132 | 15 (11) | 476 | 77 (16) |
| Other Oceania | 9 | 1 (11) | 2 | 0 (0) | 12 | 1 (8) |
| Asia | 2 | 0 (0) | 2 | 0 (0) | 4 | 0 (0) |
| UK & Ireland | 8 | 0 (0) | 3 | 0 (0) | 11 | 0 (0) |
| Other | 10 | 0 (0) | 5 | 1 (20) | 15 | 1 (7) |
| p value | | 0.323 | | 0.853 | | 0.377 |
| 2020 | | | | | | |
| Australia | 237 | 35 (15) | 94 | 10 (11) | 333 | 45 (14) |
| Other Oceania | 10 | 3 (30) | 4 | 0 (0) | 14 | 3 (21) |
| Asia | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| UK & Ireland | 8 | 0 (0) | 2 | 0 (0) | 10 | 0 (0) |
| Other | 7 | 3 (43) | 1 | 0 (0) | 8 | 3 (38) |
| p value | | 0.148 | | 0.843 | | 0.229 |
| 2021 | | | | | | |
| Australia | 261 | 27 (10) | 126 | 13 (10) | 392 | 41 (10) |
| Other Oceania | 6 | 1 (17) | 3 | 1 (33) | 10 | 2 (20) |
| Asia | 2 | 0 (0) | 1 | 0 (0) | 3 | 0 (0) |
| UK & Ireland | 8 | 1 (13) | 3 | 0 (0) | 11 | 1 (9) |
| Other | 10 | 3 (30) | 4 | 1 (25) | 14 | 4 (29) |
| p value | | 0.379 | | 0.576 | | 0.224 |
| 2022 | | | | | | |
| Australia | 197 | 16 (8) | 83 | 9 (11) | 287 | 26 (9) |
| Other Oceania | 7 | 1 (14) | 4 | 0 (0) | 11 | 1 (9) |
| Asia | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| UK & Ireland | 7 | 0 (0) | 1 | 0 (0) | 8 | 0 (0) |
| Other | 6 | 1 (17) | 2 | 0 (0) | 8 | 1 (13) |
| p value | | 0.437 | | 1.000 | | 0.858 |

South Australia

Table 6.1.1 Number (percentage) of respondents by demographic characteristics and survey year

| South Australia | 2018 | 2019 | 2020* | 2021* | 2022 |
|---|----------|----------|----------|----------|----------|
| Number of sites | 7 | 8 | 7 | 8 | 8 |
| N° surveyed | N=321 | N=326 | N=20 | N=210 | N=279 |
| Response rate (%) | 53% | 57% | -- | 50% | 49% |
| Gender (%) | | | | | |
| Male | 197 (61) | 191 (59) | 13 (65) | 133 (63) | 168 (60) |
| Female | 123 (38) | 134 (41) | 7 (35) | 70 (33) | 102 (37) |
| Other | 1 (<1) | 1 (<1) | 0 (0) | 7 (3) | 9 (3) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Sexual identity (%) | | | | | |
| Heterosexual | 248 (77) | 257 (79) | 16 (80) | 166 (79) | 219 (78) |
| Bisexual | 30 (9) | 27 (8) | 1 (5) | 23 (11) | 28 (10) |
| Gay/Lesbian | 20 (6) | 19 (6) | 0 (0) | 10 (5) | 11 (4) |
| Other | -- -- | -- -- | 1 (5) | 9 (4) | 17 (6) |
| Not reported | 23 (7) | 23 (7) | 2 (10) | 2 (1) | 4 (1) |
| Aboriginal and Torres Strait Islander origin (%) | | | | | |
| No | 264 (82) | 251 (77) | 17 (85) | 138 (66) | 200 (72) |
| Yes | 51 (16) | 67 (21) | 3 (15) | 55 (26) | 75 (27) |
| Not reported | 6 (2) | 8 (2) | 0 (0) | 17 (8) | 4 (1) |
| Region/country of birth (%) | | | | | |
| Australia | 277 (86) | 285 (87) | 17 (85) | 181 (86) | 238 (85) |
| Other Oceania | 4 (1) | 5 (2) | 0 (0) | 3 (1) | 4 (1) |
| Asia | 6 (2) | 6 (2) | 0 (0) | 3 (1) | 10 (4) |
| UK & Ireland | 21 (7) | 20 (6) | 3 (15) | 12 (6) | 13 (5) |
| Other | 11 (3) | 7 (2) | 0 (0) | 5 (2) | 6 (2) |
| Not reported | 2 (1) | 3 (1) | 0 (0) | 6 (3) | 8 (3) |
| Main language spoken at home by parents (%) | | | | | |
| English | 293 (91) | 312 (96) | 20 (100) | 200 (95) | 259 (93) |
| Non-English | 25 (8) | 12 (4) | 0 (0) | 7 (3) | 16 (6) |
| Not reported | 3 (1) | 2 (1) | 0 (0) | 3 (1) | 4 (1) |
| Age and time since first injection (years) | | | | | |
| <i>Median age</i> | 43 | 44 | 48 | 45 | 48 |
| <i>Age range</i> | 18-71 | 19-72 | 32-62 | 19-78 | 22-68 |
| Age group (%) | | | | | |
| <25 years | 10 (3) | 8 (2) | 0 (0) | 3 (1) | 4 (1) |
| 25+ years | 310 (97) | 315 (97) | 20 (100) | 206 (98) | 269 (96) |
| Not reported | 1 (<1) | 3 (1) | 0 (0) | 1 (<1) | 6 (2) |
| Median age first injection | | | | | |
| <i>Median age first injection</i> | 18 | 18 | 21 | 19 | 20 |
| <i>Age range</i> | 10-64 | 12-67 | 12-42 | 12-46 | 10-54 |
| <i>N° not reported</i> | 11 | 23 | 0 | 10 | 15 |
| Median yrs since first injection | | | | | |
| <i>Median yrs since first injection</i> | 23 | 24 | 27 | 25 | 27 |
| <i>Range</i> | 1-49 | <1-52 | <1-45 | <1-62 | <1-51 |
| Years since first injection | | | | | |
| <3 years | 8 (2) | 8 (2) | 2 (10) | 3 (1) | 11 (4) |
| 3+ years | 302 (94) | 295 (90) | 18 (90) | 196 (93) | 252 (90) |
| Not reported | 11 (3) | 23 (7) | 0 (0) | 11 (5) | 16 (6) |
| Imprisonment last year (%) | | | | | |
| No | 261 (81) | 267 (82) | 19 (95) | 176 (84) | 234 (84) |
| Yes | 33 (10) | 32 (10) | 1 (5) | 20 (10) | 37 (13) |
| Not reported | 27 (8) | 27 (8) | 0 (0) | 14 (7) | 8 (3) |
| N° in prison | | | | | |
| <i>N° in prison</i> | N=33 | N=32 | N=1 | N=20 | N=37 |
| <i>Injected in prison</i> | 9 (27) | 6 (19) | 0 (0) | 2 (10) | 10 (27) |

* Implementation of the 2020 and 2021 ANSPS was impacted by the COVID-19 pandemic, see page 1 for more information

Table 6.1.2 Number (percentage) of respondents by last drug injected, frequency of injecting and survey year

| South Australia | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|----------|----------|---------|----------|----------|
| N° surveyed | N=321 | N=326 | N=20 | N=210 | N=279 |
| Drug last injected (%) | | | | | |
| Cocaine | 2 (1) | 0 (0) | 0 (0) | 2 (1) | 0 (0) |
| Methamphetamine | 183 (57) | 175 (54) | 12 (60) | 115 (55) | 134 (48) |
| Heroin | 98 (31) | 79 (24) | 5 (25) | 57 (27) | 83 (30) |
| Pharm. opioids | 5 (2) | 10 (3) | 1 (5) | 5 (2) | 8 (3) |
| Methadone | 16 (5) | 33 (10) | 1 (5) | 9 (4) | 7 (3) |
| Buprenorphine | 2 (1) | 1 (<1) | 0 (0) | 1 (<1) | 6 (2) |
| Buprenorphine/naloxone | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 4 (1) |
| PIEDs | 0 (0) | 2 (1) | 0 (0) | 5 (2) | 2 (1) |
| More than one | 13 (4) | 17 (5) | 1 (5) | 16 (8) | 30 (11) |
| Other | 0 (0) | 2 (1) | 0 (0) | 0 (0) | 3 (1) |
| Not reported | 2 (1) | 7 (2) | 0 (0) | 0 (0) | 2 (1) |
| Ever injected opioids | | | | | |
| No | 84 (26) | 85 (26) | 7 (35) | 84 (40) | 71 (25) |
| Yes | 234 (73) | 237 (73) | 13 (65) | 126 (60) | 205 (73) |
| Not reported | 3 (1) | 4 (1) | 0 (0) | 0 (0) | 3 (1) |
| Frequency of injection last month (%) | | | | | |
| Not last month | 15 (5) | 40 (12) | 1 (5) | 38 (18) | 19 (7) |
| Less than weekly | 56 (17) | 49 (15) | 5 (25) | 43 (20) | 29 (10) |
| Weekly not daily | 94 (29) | 83 (25) | 4 (20) | 40 (19) | 68 (24) |
| Daily or more | 151 (47) | 143 (44) | 10 (50) | 83 (40) | 157 (56) |
| Not reported | 5 (2) | 11 (3) | 0 (0) | 6 (3) | 6 (2) |
| Present when someone injected for the first time in the last 12 months (%) | | | | | |
| No | 252 (79) | 255 (78) | 19 (95) | 175 (83) | 229 (82) |
| Yes | 57 (18) | 53 (16) | 1 (5) | 26 (12) | 44 (16) |
| Not reported | 12 (4) | 18 (6) | 0 (0) | 9 (4) | 6 (2) |
| Overdosed in last 12 months | | | | | |
| No | 258 (80) | 282 (87) | 18 (90) | 185 (88) | 254 (91) |
| Yes | 49 (15) | 26 (8) | 2 (10) | 15 (7) | 20 (7) |
| Not reported | 14 (4) | 18 (6) | 0 (0) | 10 (5) | 5 (2) |

Table 6.1.3 Number (percentage) of respondents by injecting behaviour in the last month and survey year

| South Australia | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|----------|----------|----------|----------|----------|
| N° injected last month | N=301 | N=275 | N=19 | N=166 | N=254 |
| Places injected last month (%) | | | | | |
| All private | 161 (53) | 156 (57) | 13 (68) | 86 (52) | 125 (49) |
| Any public | 136 (45) | 114 (41) | 6 (32) | 80 (48) | 129 (51) |
| Not reported | 4 (1) | 5 (2) | 0 (0) | 0 (0) | 0 (0) |
| Use of new and sterile needles and syringes last month (%) | | | | | |
| All injections | 232 (77) | 218 (79) | 18 (95) | 135 (81) | 202 (80) |
| Most of the time | 59 (20) | 41 (15) | 1 (5) | 23 (14) | 41 (16) |
| Half of the time | 3 (1) | 5 (2) | 0 (0) | 2 (1) | 2 (1) |
| Some of the time | 1 (<1) | 2 (1) | 0 (0) | 3 (2) | 3 (1) |
| Not last month | 0 (0) | 2 (1) | 0 (0) | 1 (1) | 0 (0) |
| Not reported | 6 (2) | 7 (3) | 0 (0) | 2 (1) | 6 (2) |
| Re-used someone else's used needle & syringe last month (%) | | | | | |
| None | 258 (86) | 224 (81) | 19 (100) | 141 (85) | 210 (83) |
| Once | 18 (6) | 19 (7) | 0 (0) | 7 (4) | 13 (5) |
| Twice | 10 (3) | 15 (5) | 0 (0) | 6 (4) | 16 (6) |
| 3-5 times | 5 (2) | 5 (2) | 0 (0) | 5 (3) | 6 (2) |
| >5 times | 6 (2) | 5 (2) | 0 (0) | 4 (2) | 7 (3) |
| Not reported | 4 (1) | 7 (3) | 0 (0) | 3 (2) | 2 (1) |
| Number of people needle & syringe was re-used after last month (%) | | | | | |
| None | 258 (86) | 224 (81) | 19 (100) | 141 (85) | 210 (83) |
| One | 21 (7) | 18 (7) | 0 (0) | 10 (6) | 19 (7) |
| Two | 3 (1) | 5 (2) | 0 (0) | 3 (2) | 8 (3) |
| Three to five | 0 (0) | 4 (1) | 0 (0) | 1 (1) | 1 (<1) |
| More than five | 0 (0) | 3 (1) | 0 (0) | 1 (1) | 3 (1) |
| Don't know | 7 (2) | 8 (3) | 0 (0) | 5 (3) | 9 (4) |
| Not reported | 12 (4) | 13 (5) | 0 (0) | 5 (3) | 4 (2) |
| Relationship to people needle & syringe was used after last month (%) [more than one could be selected] | | | | | |
| Regular sex partner | 17 (6) | 24 (9) | 0 (0) | 9 (5) | 17 (7) |
| Casual sex partner | 1 (<1) | 1 (<1) | 0 (0) | 2 (1) | 2 (1) |
| Close friend | 9 (3) | 8 (3) | 0 (0) | 9 (5) | 15 (6) |
| Acquaintance | 4 (1) | 6 (2) | 0 (0) | 1 (1) | 7 (3) |
| Other | 1 (<1) | 4 (1) | 0 (0) | 1 (1) | 2 (1) |
| Equipment used after someone else last month (%) [more than one could be selected] | | | | | |
| Spoon | 41 (14) | 45 (16) | 1 (5) | 29 (17) | 43 (17) |
| Water | 62 (21) | 56 (20) | 1 (5) | 36 (22) | 56 (22) |
| Filter | 23 (8) | 24 (9) | 0 (0) | 11 (7) | 21 (8) |
| Drug mix | 32 (11) | 22 (8) | 1 (5) | 11 (7) | 18 (7) |
| None | 208 (69) | 184 (67) | 16 (84) | 107 (64) | 167 (66) |
| Injected by someone after that person injected themselves or others last month (%) | | | | | |
| No | 250 (83) | 231 (84) | 18 (95) | 138 (83) | 196 (77) |
| Yes | 44 (15) | 38 (14) | 1 (5) | 25 (15) | 55 (22) |
| Not reported | 7 (2) | 6 (2) | 0 (0) | 3 (2) | 3 (1) |
| Source of needle acquisition | | | | | |
| Needle Syringe Program | 266 (88) | 238 (87) | 16 (84) | 142 (86) | 228 (90) |
| Chemist/Pharmacy | 60 (20) | 44 (16) | 3 (16) | 25 (15) | 33 (13) |
| Personal sources | 53 (18) | 43 (16) | 4 (21) | 18 (11) | 40 (16) |
| Dispensing/Vending Machine | 56 (19) | 55 (20) | 3 (16) | 36 (22) | 35 (14) |
| Other sources | 4 (1) | 3 (1) | 0 (0) | 3 (2) | 2 (1) |

Table 6.1.4 Number (percentage) of respondents by treatment for drug use and survey year

| South Australia | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|----------|----------|---------|----------|----------|
| N° surveyed | N=321 | N=326 | N=20 | N=210 | N=279 |
| Ever any treatment/therapy for drug use (%) | | | | | |
| No | 104 (32) | 109 (33) | 6 (30) | 87 (41) | 93 (33) |
| Yes | 216 (67) | 214 (66) | 14 (70) | 123 (59) | 185 (66) |
| Not reported | 1 (<1) | 3 (1) | 0 (0) | 0 (0) | 1 (<1) |
| History of methadone maintenance treatment (%) | | | | | |
| Currently | 68 (21) | 86 (26) | 4 (20) | 34 (16) | 53 (19) |
| Previously | 81 (25) | 49 (15) | 2 (10) | 44 (21) | 66 (24) |
| Never | 160 (50) | 181 (56) | 13 (65) | 120 (57) | 149 (53) |
| Not reported | 12 (4) | 10 (3) | 1 (5) | 12 (6) | 11 (4) |
| History of buprenorphine (Subutex®) treatment (%) | | | | | |
| Currently | 13 (4) | 10 (3) | 2 (10) | 8 (4) | 17 (6) |
| Previously | 86 (27) | 70 (21) | 6 (30) | 56 (27) | 76 (27) |
| Never | 214 (67) | 240 (74) | 12 (60) | 138 (66) | 175 (63) |
| Not reported | 8 (2) | 6 (2) | 0 (0) | 8 (4) | 11 (4) |
| History of buprenorphine-naloxone (Suboxone®) treatment (%) | | | | | |
| Currently | 22 (7) | 21 (6) | 2 (10) | 24 (11) | 18 (6) |
| Previously | 69 (22) | 57 (17) | 5 (25) | 44 (21) | 75 (27) |
| Never | 222 (69) | 240 (74) | 13 (65) | 142 (68) | 185 (66) |
| Not reported | 8 (2) | 8 (2) | 0 (0) | 0 (0) | 1 (<1) |

Table 6.1.5 Number (percentage) of respondents by testing for HIV and HCV, HCV treatment by survey year

| South Australia | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|----------|----------|---------|----------|----------|
| N° surveyed | N=321 | N=326 | N=20 | N=210 | N=279 |
| Previous HIV test (%) | | | | | |
| Yes, ever | 256 (80) | 241 (74) | 16 (80) | 136 (65) | 186 (67) |
| Yes, last year | 149 (46) | 124 (38) | 8 (40) | 59 (28) | 94 (34) |
| >1 year ago | 107 (33) | 117 (36) | 8 (40) | 77 (37) | 92 (33) |
| Never tested | 52 (16) | 60 (18) | 4 (20) | 63 (30) | 79 (28) |
| Not reported | 13 (4) | 25 (8) | 0 (0) | 11 (5) | 14 (5) |
| Previous HCV test (%) | | | | | |
| Yes, ever | 241 (75) | 241 (74) | 17 (85) | 139 (66) | 203 (73) |
| Yes, last year | 145 (45) | 150 (46) | 8 (40) | 66 (31) | 123 (44) |
| >1 year ago | 96 (30) | 91 (28) | 9 (45) | 73 (35) | 80 (29) |
| Never tested | 45 (14) | 48 (15) | 2 (10) | 51 (24) | 52 (19) |
| Not reported | 35 (11) | 37 (11) | 1 (5) | 20 (10) | 24 (9) |
| Lifetime treatment for HCV (%)# | | | | | |
| N° self-reported HCV diagnosis | N=68 | N=56 | N=3 | N=23 | N=54 |
| Antiviral treatment | 44 (65) | 41 (73) | 3 (100) | 18 (78) | 40 (74) |
| No antiviral treatment | 23 (34) | 15 (27) | 0 (0) | 5 (22) | 14 (26) |
| Not reported | 1 (1) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Treatment for HCV in past 12 months (%)** | | | | | |
| N° self-reported HCV diagnosis | N=53 | N=38 | N=1 | N=11 | N=21 |
| Antiviral treatment | 22 (42) | 21 (55) | 1 (100) | 3 (27) | 5 (24) |
| No antiviral treatment | 30 (57) | 17 (45) | 0 (0) | 8 (73) | 16 (76) |
| Not reported | 1 (2) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |

among people who tested HCV antibody positive and did not report spontaneous clearance

* excludes people who reported treatment induced clearance more than 12 months previously

Table 6.1.6 Number (percentage) of respondents by sexual behaviour in the month prior to survey and survey year

| South Australia | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|----------|----------|----------|----------|----------|
| N° surveyed | N=321 | N=326 | N=20 | N=210 | N=279 |
| Sex with a regular partner last month (%) | | | | | |
| No | 152 (47) | 165 (51) | 14 (70) | 116 (55) | 148 (53) |
| Yes | 145 (45) | 134 (41) | 6 (30) | 83 (40) | 120 (43) |
| Not reported | 24 (7) | 27 (8) | 0 (0) | 11 (5) | 11 (4) |
| Condom used with regular partner last month (%) | | | | | |
| N° surveyed | N=145 | N=134 | N=6 | N=83 | N=120 |
| Never | 114 (79) | 89 (66) | 4 (67) | 58 (70) | 103 (86) |
| Sometimes | 7 (5) | 18 (13) | 0 (0) | 17 (20) | 9 (8) |
| Every time | 16 (11) | 19 (14) | 0 (0) | 5 (6) | 4 (3) |
| Not reported | 8 (6) | 8 (6) | 2 (33) | 3 (4) | 4 (3) |
| Sex with other partner(s) last month (%) | | | | | |
| No | 257 (80) | 254 (78) | 18 (90) | 165 (79) | 231 (83) |
| Yes | 38 (12) | 43 (13) | 2 (10) | 33 (16) | 35 (13) |
| Not reported | 26 (8) | 29 (9) | 0 (0) | 12 (6) | 13 (5) |
| Condom used with other partner(s) last month (%) | | | | | |
| N° surveyed | N=38 | N=43 | N=2 | N=33 | N=35 |
| Never | 7 (18) | 21 (49) | 1 (50) | 17 (52) | 23 (66) |
| Sometimes | 14 (37) | 10 (23) | 1 (50) | 5 (15) | 6 (17) |
| Every time | 17 (45) | 12 (28) | 0 (0) | 9 (27) | 5 (14) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 2 (6) | 1 (3) |
| Sex work last month (%) | | | | | |
| No | 286 (89) | 279 (86) | 20 (100) | 186 (89) | 244 (87) |
| Yes | 15 (5) | 12 (4) | 0 (0) | 11 (5) | 19 (7) |
| Not reported | 20 (6) | 35 (11) | 0 (0) | 13 (6) | 16 (6) |
| Condom used at last sex work (%) | | | | | |
| Yes | 10 (67) | 7 (58) | 0 (0) | 8 (73) | 14 (74) |

Table 6.1.7 Number (percentage) of respondents by COVID-19 related impacts, 2020-2022

| South Australia | 2020 | 2021 | 2022 |
|---|---------|----------|----------|
| N° surveyed | N=20 | N=210 | N=279 |
| COVID-19 test | | | |
| Yes | 4 (20) | 70 (33) | 227 (81) |
| No | 15 (75) | 93 (44) | 51 (18) |
| Don't know | 1 (5) | 15 (7) | -- -- |
| Not reported | 0 (0) | 32 (15) | 1 (<1) |
| Self-reported positive COVID-19 test result | | | |
| Yes | 0 (0) | 1 (<1) | 97 (43) |
| Yes, last 12 months | -- -- | -- -- | 77 (34) |
| Yes, more than a year ago | -- -- | -- -- | 20 (9) |
| COVID-19 vaccine uptake | | | |
| ≥1 dose | -- -- | 93 (44) | 206 (74) |
| COVID-19 vaccine doses | | | |
| One dose | -- -- | 44 (21) | 26 (9) |
| Two doses | -- -- | 49 (23) | 53 (19) |
| Three or more doses | -- -- | -- -- | 127 (46) |
| No doses | -- -- | 94 (45) | 69 (25) |
| Not reported | -- -- | 23 (11) | 4 (1) |
| Vaccine type * | | | N=206 |
| Astra Zeneca | -- -- | 11 (5) | 49 (24) |
| Pfizer | -- -- | 48 (23) | 132 (64) |
| Don't know | -- -- | 20 (10) | 27 (13) |
| Other | -- -- | -- -- | 19 (9) |
| Not reported | -- -- | 131 (62) | -- -- |
| * 2021 only one response permitted, 2022 multiple responses permitted | | | |
| Last vaccine location | | | N=279 |
| GP or medical centre | -- -- | -- -- | 83 (30) |
| Pharmacy/ chemist | -- -- | -- -- | 47 (17) |
| Vaccine hub | -- -- | -- -- | 34 (12) |
| Opioid treatment/ methadone clinic | -- -- | -- -- | 4 (1) |
| NSP | -- -- | -- -- | 4 (1) |
| Prison | -- -- | -- -- | 17 (6) |
| Other | -- -- | -- -- | 7 (3) |
| Multiple response | -- -- | -- -- | 14 (5) |
| Not reported | -- -- | -- -- | 69 (25) |
| Vaccine hesitancy - if not had three or more doses | | | |
| Haven't got around to it | -- -- | -- -- | 66 (24) |
| Difficulty booking appointment | -- -- | -- -- | 7 (3) |
| Concerns about side effects | -- -- | -- -- | 17 (6) |
| Concerns about vaccine safety | -- -- | -- -- | 18 (6) |
| Not at risk of serious illness | -- -- | -- -- | 14 (5) |
| Other | -- -- | -- -- | 26 (9) |
| Multiple response | -- -- | -- -- | 15 (5) |
| Not reported | -- -- | -- -- | 116 (42) |

HIV antibody prevalence

Table 6.2.1 HIV antibody prevalence by gender and survey year

| South Australia | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Survey year | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | 193 | 2 (1.0) | 122 | 0 (0.0) | 315 | 2 (0.6) |
| 2019 | 177 | 4 (2.3) | 126 | 0 (0.0) | 304 | 4 (1.3) |
| 2020 | 13 | 0 (0.0) | 7 | 0 (0.0) | 20 | 0 (0.0) |
| 2021 | 116 | 0 (0.0) | 67 | 0 (0.0) | 188 | 0 (0.0) |
| 2022 | 163 | 1 (0.6) | 102 | 0 (0.0) | 273 | 1 (0.4) |
| X ² p trend | | 0.262 | | -- | | 0.262 |

HCV antibody prevalence

Table 6.3.1 HCV antibody prevalence by gender and survey year

| South Australia | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Survey year | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | 187 | 89 (48) | 122 | 34 (28) | 309 | 123 (40) |
| 2019 | 175 | 82 (47) | 126 | 43 (34) | 302 | 125 (41) |
| 2020 | 13 | 2 (15) | 7 | 2 (29) | 20 | 4 (20) |
| 2021 | 116 | 36 (31) | 67 | 20 (30) | 188 | 57 (30) |
| 2022 | 162 | 54 (33) | 102 | 36 (35) | 272 | 93 (34) |
| X ² p trend | | <0.001 | | 0.410 | | 0.021 |

Table 6.3.2 HCV antibody prevalence by sexual identity, gender and survey year

| South Australia | Male | | Female | | Total | |
|-----------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Sexual identity | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heterosexual | 161 | 81 (50) | 76 | 22 (29) | 237 | 103 (43) |
| Bisexual | 4 | 0 (0) | 26 | 7 (27) | 30 | 7 (23) |
| Homosexual | 9 | 2 (22) | 11 | 2 (18) | 20 | 4 (20) |
| p value | | 0.045 | | 0.847 | | 0.019 |
| 2019 | | | | | | |
| Heterosexual | 147 | 73 (50) | 92 | 33 (36) | 239 | 106 (44) |
| Bisexual | 6 | 2 (33) | 17 | 5 (29) | 24 | 7 (29) |
| Homosexual | 8 | 0 (0) | 10 | 1 (10) | 18 | 1 (6) |
| p value | | 0.009 | | 0.256 | | 0.001 |
| 2020 | | | | | | |
| Heterosexual | 11 | 2 (18) | 5 | 1 (20) | 16 | 3 (19) |
| Bisexual | 0 | 0 (0) | 1 | 1 (100) | 1 | 1 (100) |
| Homosexual | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | 1.000 | | 0.333 | | 0.405 |
| 2021 | | | | | | |
| Heterosexual | 100 | 31 (31) | 49 | 13 (27) | 149 | 44 (30) |
| Bisexual | 7 | 4 (57) | 14 | 4 (29) | 22 | 8 (36) |
| Homosexual | 7 | 1 (14) | 0 | 0 (0) | 8 | 1 (13) |
| p value | | 0.361 | | 0.749 | | 0.828 |
| 2022 | | | | | | |
| Heterosexual | 135 | 50 (37) | 75 | 26 (35) | 213 | 78 (37) |
| Bisexual | 11 | 4 (36) | 16 | 7 (44) | 28 | 11 (39) |
| Homosexual | 7 | 0 (0) | 4 | 1 (25) | 11 | 1 (9) |
| p value | | 0.158 | | 0.780 | | 0.287 |

Table 6.3.3 HCV antibody prevalence by age group, gender and survey year

| South Australia Age group | Male | | Female | | Total | |
|------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <25 years | 4 | 0 (0) | 6 | 0 (0) | 10 | 0 (0) |
| 25-34 years | 33 | 9 (27) | 20 | 4 (20) | 53 | 13 (25) |
| 35-44 years | 63 | 31 (49) | 48 | 15 (31) | 111 | 46 (41) |
| 45+ years | 87 | 49 (56) | 47 | 15 (32) | 134 | 64 (48) |
| p value | | 0.006 | | 0.368 | | 0.001 |
| 2019 | | | | | | |
| <25 years | 4 | 1 (25) | 3 | 1 (33) | 7 | 2 (29) |
| 25-34 years | 21 | 7 (33) | 23 | 4 (17) | 44 | 11 (25) |
| 35-44 years | 57 | 20 (35) | 45 | 15 (33) | 102 | 35 (34) |
| 45+ years | 91 | 54 (59) | 55 | 23 (42) | 147 | 77 (52) |
| p value | | 0.008 | | 0.195 | | 0.002 |
| 2020 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 35-44 years | 4 | 0 (0) | 2 | 0 (0) | 6 | 0 (0) |
| 45+ years | 8 | 2 (25) | 5 | 2 (40) | 13 | 4 (31) |
| p value | | 0.590 | | 1.000 | | 0.404 |
| 2021 | | | | | | |
| <25 years | 3 | 1 (33) | 0 | 0 (0) | 3 | 1 (33) |
| 25-34 years | 15 | 2 (13) | 9 | 2 (22) | 24 | 4 (17) |
| 35-44 years | 37 | 16 (43) | 28 | 9 (32) | 67 | 25 (37) |
| 45+ years | 61 | 17 (28) | 30 | 9 (30) | 94 | 27 (29) |
| p value | | 0.142 | | 0.936 | | 0.232 |
| 2022 | | | | | | |
| <25 years | 4 | 0 (0) | 0 | 0 (0) | 4 | 0 (0) |
| 25-34 years | 14 | 1 (7) | 10 | 2 (20) | 24 | 3 (13) |
| 35-44 years | 42 | 11 (26) | 27 | 8 (30) | 69 | 19 (28) |
| 45+ years | 98 | 41 (42) | 64 | 25 (39) | 169 | 68 (40) |
| p value | | 0.014 | | 0.409 | | 0.008 |

Table 6.3.4 HCV antibody prevalence by years since first injection, gender and survey year

| South Australia Years of injection | Male | | Female | | Total | |
|---------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <3 years | 3 | 0 (0) | 5 | 0 (0) | 8 | 0 (0) |
| 3 to 10 years | 24 | 3 (13) | 19 | 2 (11) | 43 | 5 (12) |
| 11+ years | 154 | 82 (53) | 96 | 32 (33) | 250 | 114 (46) |
| p value | | <0.001 | | 0.053 | | <0.001 |
| 2019 | | | | | | |
| <3 years | 3 | 0 (0) | 4 | 0 (0) | 7 | 0 (0) |
| 3 to 10 years | 16 | 2 (13) | 20 | 7 (35) | 36 | 9 (25) |
| 11+ years | 145 | 74 (51) | 92 | 33 (36) | 238 | 107 (45) |
| p value | | 0.002 | | 0.466 | | 0.005 |
| 2020 | | | | | | |
| <3 years | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| 3 to 10 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 11+ years | 10 | 2 (20) | 7 | 2 (29) | 17 | 4 (24) |
| p value | | 1.000 | | -- | | 1.000 |
| 2021 | | | | | | |
| <3 years | 1 | 1 (100) | 1 | 0 (0) | 2 | 1 (50) |
| 3 to 10 years | 10 | 3 (30) | 9 | 0 (0) | 19 | 3 (16) |
| 11+ years | 98 | 28 (29) | 54 | 19 (35) | 157 | 48 (31) |
| p value | | 0.405 | | 0.064 | | 0.256 |
| 2022 | | | | | | |
| <3 years | 9 | 0 (0) | 2 | 1 (50) | 11 | 1 (9) |
| 3 to 10 years | 14 | 2 (14) | 10 | 1 (10) | 25 | 4 (16) |
| 11+ years | 128 | 49 (38) | 86 | 33 (38) | 220 | 83 (38) |
| p value | | 0.013 | | 0.139 | | 0.016 |

Table 6.3.5 HCV antibody prevalence by re-use of someone else's used needle and syringe last month, years since first injection and survey year

| South Australia Receptively shared syringe last month | <3 years | | 3+ years | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| No receptive sharing | 4 | 0 (0) | 240 | 94 (39) | 249 | 97 (39) |
| Receptive sharing | 3 | 0 (0) | 32 | 18 (56) | 37 | 19 (51) |
| p value | | -- | | 0.065 | | 0.152 |
| 2019 | | | | | | |
| No receptive sharing | 5 | 0 (0) | 193 | 83 (43) | 209 | 90 (43) |
| Receptive sharing | 0 | 0 (0) | 35 | 12 (34) | 38 | 13 (34) |
| p value | | -- | | 0.336 | | 0.309 |
| 2020 | | | | | | |
| No receptive sharing | 2 | 0 (0) | 17 | 4 (24) | 19 | 4 (21) |
| Receptive sharing | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No receptive sharing | 2 | 1 (50) | 118 | 40 (34) | 127 | 44 (35) |
| Receptive sharing | 0 | 0 (0) | 17 | 4 (24) | 19 | 5 (26) |
| p value | | -- | | 0.581 | | 0.473 |
| 2022 | | | | | | |
| No receptive sharing | 9 | 1 (11) | 190 | 65 (34) | 205 | 69 (34) |
| Receptive sharing | 1 | 0 (0) | 36 | 14 (39) | 41 | 14 (34) |
| p value | | 1.000 | | 0.589 | | 0.952 |

Table 6.3.6 HCV antibody prevalence by last drug injected, years since first injection and survey year

| South Australia Last drug injected | <3 years | | 3+ years | | Total | |
|---------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heroin | 1 | 0 (0) | 92 | 48 (52) | 95 | 50 (53) |
| Methamphetamine | 7 | 0 (0) | 163 | 53 (33) | 174 | 54 (31) |
| Other opioids | 0 | 0 (0) | 21 | 11 (52) | 21 | 11 (52) |
| Other drugs | 0 | 0 (0) | 12 | 4 (33) | 13 | 5 (38) |
| p value | | -- | | 0.012 | | 0.004 |
| 2019 | | | | | | |
| Heroin | 0 | 0 (0) | 69 | 43 (62) | 72 | 46 (64) |
| Methamphetamine | 6 | 0 (0) | 141 | 40 (28) | 157 | 44 (28) |
| Other opioids | 0 | 0 (0) | 40 | 19 (48) | 41 | 20 (49) |
| Other drugs | 1 | 0 (0) | 17 | 10 (59) | 20 | 10 (50) |
| p value | | -- | | <0.001 | | <0.001 |
| 2020 | | | | | | |
| Heroin | 0 | 0 (0) | 5 | 2 (40) | 5 | 2 (40) |
| Methamphetamine | 2 | 0 (0) | 10 | 2 (20) | 12 | 2 (17) |
| Other opioids | 0 | 0 (0) | 2 | 0 (0) | 2 | 0 (0) |
| Other drugs | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | 0.615 | | 0.555 |
| 2021 | | | | | | |
| Heroin | 0 | 0 (0) | 47 | 20 (43) | 51 | 22 (43) |
| Methamphetamine | 2 | 1 (50) | 92 | 20 (22) | 99 | 23 (23) |
| Other opioids | 0 | 0 (0) | 13 | 4 (31) | 14 | 5 (36) |
| Other drugs | 0 | 0 (0) | 19 | 5 (26) | 19 | 5 (26) |
| p value | | -- | | 0.083 | | 0.083 |
| 2022 | | | | | | |
| Heroin | 0 | 0 (0) | 75 | 37 (49) | 79 | 40 (51) |
| Methamphetamine | 7 | 1 (14) | 120 | 31 (26) | 131 | 32 (24) |
| Other opioids | 1 | 0 (0) | 19 | 9 (47) | 22 | 9 (41) |
| Other drugs | 2 | 0 (0) | 27 | 8 (30) | 32 | 8 (25) |
| p value | | 0.788 | | 0.005 | | 0.001 |

Table 6.3.7 HCV antibody prevalence by frequency of drug injection last month, years since first injection and survey year

| South Australia | <3 years | | 3+ years | | Total | |
|-----------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| Frequency of injection last month | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| Less than daily | 2 | 0 (0) | 141 | 56 (40) | 147 | 58 (39) |
| Daily or more | 5 | 0 (0) | 135 | 58 (43) | 143 | 60 (42) |
| Not last month | 1 | 0 (0) | 13 | 3 (23) | 14 | 3 (21) |
| p value | | -- | | 0.392 | | 0.335 |
| 2019 | | | | | | |
| Less than daily | 3 | 0 (0) | 116 | 44 (38) | 124 | 47 (38) |
| Daily or more | 3 | 0 (0) | 118 | 52 (44) | 130 | 57 (44) |
| Not last month | 1 | 0 (0) | 33 | 16 (48) | 38 | 16 (42) |
| p value | | -- | | 0.457 | | 0.624 |
| 2020 | | | | | | |
| Less than daily | 2 | 0 (0) | 7 | 2 (29) | 9 | 2 (22) |
| Daily or more | 0 | 0 (0) | 10 | 2 (20) | 10 | 2 (20) |
| Not last month | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | 1.000 | | 1.000 |
| 2021 | | | | | | |
| Less than daily | 1 | 0 (0) | 72 | 17 (24) | 75 | 18 (24) |
| Daily or more | 1 | 1 (100) | 66 | 29 (44) | 74 | 33 (45) |
| Not last month | 0 | 0 (0) | 33 | 3 (9) | 34 | 4 (12) |
| p value | | 1.000 | | 0.001 | | 0.001 |
| 2022 | | | | | | |
| Less than daily | 8 | 1 (13) | 84 | 27 (32) | 94 | 28 (30) |
| Daily or more | 2 | 0 (0) | 143 | 53 (37) | 154 | 56 (36) |
| Not last month | 1 | 0 (0) | 15 | 5 (33) | 18 | 5 (28) |
| p value | | 1.000 | | 0.747 | | 0.493 |

Table 6.3.8 HCV antibody prevalence by imprisonment last year, years since first injection and survey year

| South Australia | <3 years | | 3+ years | | Total | |
|------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| Imprisonment last year | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| No imprisonment | 6 | 0 (0) | 241 | 93 (39) | 251 | 96 (38) |
| Imprisonment | 2 | 0 (0) | 29 | 16 (55) | 31 | 16 (52) |
| p value | | -- | | 0.086 | | 0.151 |
| 2019 | | | | | | |
| No imprisonment | 5 | 0 (0) | 222 | 92 (41) | 246 | 101 (41) |
| Imprisonment | 1 | 0 (0) | 30 | 15 (50) | 32 | 15 (47) |
| p value | | -- | | 0.373 | | 0.530 |
| 2020 | | | | | | |
| No imprisonment | 2 | 0 (0) | 17 | 4 (24) | 19 | 4 (21) |
| Imprisonment | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | 1.000 | | 1.000 |
| 2021 | | | | | | |
| No imprisonment | 1 | 0 (0) | 149 | 43 (29) | 156 | 47 (30) |
| Imprisonment | 0 | 0 (0) | 16 | 1 (6) | 19 | 1 (5) |
| p value | | -- | | 0.072 | | 0.027 |
| 2022 | | | | | | |
| No imprisonment | 10 | 1 (10) | 207 | 76 (37) | 228 | 80 (35) |
| Imprisonment | 1 | 0 (0) | 31 | 7 (23) | 36 | 8 (22) |
| p value | | 1.000 | | 0.124 | | 0.128 |

Table 6.3.9 HCV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| South Australia | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Aboriginal and Torres Strait Islander origin | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Non Indigenous | 154 | 65 (42) | 102 | 31 (30) | 256 | 96 (38) |
| Indigenous | 30 | 22 (73) | 17 | 3 (18) | 47 | 25 (53) |
| p value | | 0.002 | | 0.389 | | 0.043 |
| 2019 | | | | | | |
| Non Indigenous | 133 | 65 (49) | 98 | 33 (34) | 231 | 98 (42) |
| Indigenous | 39 | 14 (36) | 24 | 9 (38) | 64 | 23 (36) |
| p value | | 0.153 | | 0.724 | | 0.351 |
| 2020 | | | | | | |
| Non Indigenous | 10 | 2 (20) | 7 | 2 (29) | 17 | 4 (24) |
| Indigenous | 3 | 0 (0) | 0 | 0 (0) | 3 | 0 (0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2021 | | | | | | |
| Non Indigenous | 78 | 20 (26) | 43 | 12 (28) | 124 | 32 (26) |
| Indigenous | 28 | 12 (43) | 18 | 6 (33) | 48 | 19 (40) |
| p value | | 0.089 | | 0.672 | | 0.076 |
| 2022 | | | | | | |
| Non Indigenous | 115 | 40 (35) | 73 | 27 (37) | 194 | 70 (36) |
| Indigenous | 45 | 13 (29) | 28 | 9 (32) | 74 | 22 (30) |
| p value | | 0.476 | | 0.649 | | 0.327 |

Table 6.3.10 HCV antibody prevalence by main language spoken at home by parents, gender and survey year

| South Australia | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Main language spoken at home by parents | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| English speaking | 167 | 81 (49) | 119 | 34 (29) | 286 | 115 (40) |
| Non-English speaking | 17 | 8 (47) | 3 | 0 (0) | 20 | 8 (40) |
| p value | | 0.910 | | 0.559 | | 0.985 |
| 2019 | | | | | | |
| English speaking | 165 | 79 (48) | 124 | 42 (34) | 290 | 121 (42) |
| Non-English speaking | 9 | 3 (33) | 2 | 1 (50) | 11 | 4 (36) |
| p value | | 0.503 | | 1.000 | | 1.000 |
| 2020 | | | | | | |
| English speaking | 13 | 2 (15) | 7 | 2 (29) | 20 | 4 (20) |
| Non-English speaking | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| English speaking | 110 | 35 (32) | 63 | 19 (30) | 178 | 55 (31) |
| Non-English speaking | 3 | 1 (33) | 4 | 1 (25) | 7 | 2 (29) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2022 | | | | | | |
| English speaking | 150 | 49 (33) | 95 | 34 (36) | 252 | 86 (34) |
| Non-English speaking | 10 | 5 (50) | 5 | 1 (20) | 16 | 6 (38) |
| p value | | 0.262 | | 0.655 | | 0.783 |

Table 6.3.11 HCV antibody prevalence by region/country of birth, gender and survey year

| South Australia | | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | |
| 2018 | | | | | | | |
| Australia | 159 | 75 (47) | 110 | 30 (27) | 269 | 105 (39) | |
| Other Oceania | 2 | 0 (0) | 2 | 1 (50) | 4 | 1 (25) | |
| Asia | 6 | 4 (67) | 0 | 0 (0) | 6 | 4 (67) | |
| UK & Ireland | 12 | 7 (58) | 8 | 3 (38) | 20 | 10 (50) | |
| Other | 7 | 3 (43) | 2 | 0 (0) | 9 | 3 (33) | |
| p value | | 0.592 | | 0.660 | | 0.547 | |
| 2019 | | | | | | | |
| Australia | 152 | 69 (45) | 113 | 38 (34) | 266 | 107 (40) | |
| Other Oceania | 2 | 1 (50) | 2 | 0 (0) | 4 | 1 (25) | |
| Asia | 5 | 1 (20) | 1 | 1 (100) | 6 | 2 (33) | |
| UK & Ireland | 10 | 7 (70) | 8 | 3 (38) | 18 | 10 (56) | |
| Other | 4 | 2 (50) | 2 | 1 (50) | 6 | 3 (50) | |
| p value | | 0.426 | | 0.600 | | 0.656 | |
| 2020 | | | | | | | |
| Australia | 12 | 2 (17) | 5 | 1 (20) | 17 | 3 (18) | |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| UK & Ireland | 1 | 0 (0) | 2 | 1 (50) | 3 | 1 (33) | |
| Other | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| p value | | 1.000 | | 1.000 | | 0.509 | |
| 2021 | | | | | | | |
| Australia | 99 | 29 (29) | 58 | 14 (24) | 162 | 44 (27) | |
| Other Oceania | 1 | 1 (100) | 2 | 2 (100) | 3 | 3 (100) | |
| Asia | 2 | 1 (50) | 1 | 1 (100) | 3 | 2 (67) | |
| UK & Ireland | 7 | 2 (29) | 3 | 2 (67) | 10 | 4 (40) | |
| Other | 3 | 0 (0) | 2 | 1 (50) | 5 | 1 (20) | |
| p value | | 0.419 | | 0.013 | | 0.029 | |
| 2022 | | | | | | | |
| Australia | 140 | 44 (31) | 86 | 29 (34) | 233 | 76 (33) | |
| Other Oceania | 2 | 2 (100) | 2 | 1 (50) | 4 | 3 (75) | |
| Asia | 5 | 1 (20) | 3 | 1 (33) | 8 | 2 (25) | |
| UK & Ireland | 5 | 2 (40) | 8 | 5 (63) | 13 | 7 (54) | |
| Other | 3 | 1 (33) | 2 | 0 (0) | 6 | 1 (17) | |
| p value | | 0.340 | | 0.412 | | 0.161 | |

HCV RNA prevalence

Table 6.4.1 HCV RNA prevalence by gender and survey year

| South Australia | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Survey year | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | 87 | 21 (24) | 56 | 8 (14) | 143 | 29 (20) |
| 2019* | 117 | 18 (15) | 85 | 5 (6) | 203 | 23 (11) |
| 2020 | 9 | 1 (11) | 5 | 0 (0) | 14 | 1 (7) |
| 2021 | 87 | 16 (18) | 54 | 5 (9) | 146 | 22 (15) |
| 2022 | 149 | 19 (13) | 87 | 13 (15) | 243 | 32 (13) |
| X ² p trend | | 0.083 | | 0.421 | | 0.307 |

* 2018, 2019: Weighted for gender and HCV antibody status

Totals include respondents where gender was reported as other or not reported

Table 6.4.2 HCV RNA prevalence by sexual identity, gender and survey year

| South Australia | Male | | Female | | Total | |
|-----------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Sexual identity | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heterosexual | 66 | 17 (26) | 35 | 6 (17) | 100 | 23 (23) |
| Bisexual | 4 | 0 (0) | 12 | 1 (8) | 16 | 1 (6) |
| Homosexual | 8 | 0 (0) | 5 | 0 (0) | 14 | 0 (0) |
| p value | | 0.178 | | 0.369 | | 0.037 |
| 2019* | | | | | | |
| Heterosexual | 96 | 15 (16) | 58 | 4 (7) | 155 | 19 (12) |
| Bisexual | 5 | 1 (20) | 12 | 0 (0) | 18 | 1 (6) |
| Homosexual | 8 | 0 (0) | 9 | 0 (0) | 17 | 0 (0) |
| p value | | 0.492 | | 0.444 | | 0.221 |
| 2020 | | | | | | |
| Heterosexual | 7 | 1 (14) | 4 | 0 (0) | 11 | 1 (9) |
| Bisexual | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| Homosexual | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | 0.753 |
| 2021 | | | | | | |
| Heterosexual | 76 | 14 (18) | 38 | 3 (8) | 114 | 17 (15) |
| Bisexual | 4 | 2 (50) | 13 | 1 (8) | 18 | 3 (17) |
| Homosexual | 5 | 0 (0) | 0 | 0 (0) | 6 | 0 (0) |
| p value | | 0.156 | | 0.981 | | 0.576 |
| 2022 | | | | | | |
| Heterosexual | 122 | 17 (14) | 63 | 10 (16) | 188 | 27 (14) |
| Bisexual | 11 | 2 (18) | 14 | 2 (14) | 25 | 4 (16) |
| Homosexual | 7 | 0 (0) | 3 | 0 (0) | 10 | 0 (0) |
| p value | | 0.859 | | 1.000 | | 0.865 |

Table 6.4.3 HCV RNA prevalence by age group, gender and survey year

| South Australia Age group | Male | | Female | | Total | |
|------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <25 years | 1 | 0 (0) | 4 | 0 (0) | 6 | 0 (0) |
| 25-34 years | 23 | 5 (22) | 14 | 3 (21) | 37 | 8 (22) |
| 35-44 years | 30 | 7 (23) | 23 | 5 (22) | 53 | 12 (23) |
| 45+ years | 33 | 9 (27) | 14 | 0 (0) | 47 | 9 (19) |
| p value | | 0.903 | | 0.155 | | 0.630 |
| 2019* | | | | | | |
| <25 years | 2 | 0 (0) | 3 | 0 (0) | 5 | 0 (0) |
| 25-34 years | 17 | 2 (12) | 16 | 2 (13) | 33 | 4 (12) |
| 35-44 years | 37 | 3 (8) | 27 | 1 (4) | 63 | 4 (6) |
| 45+ years | 60 | 13 (22) | 39 | 2 (5) | 100 | 15 (15) |
| p value | | 0.307 | | 0.680 | | 0.322 |
| 2020 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 35-44 years | 3 | 0 (0) | 1 | 0 (0) | 4 | 0 (0) |
| 45+ years | 6 | 1 (17) | 4 | 0 (0) | 10 | 1 (10) |
| p value | | 0.501 | | -- | | 0.525 |
| 2021 | | | | | | |
| <25 years | 3 | 1 (33) | 0 | 0 (0) | 3 | 1 (33) |
| 25-34 years | 11 | 2 (18) | 9 | 1 (11) | 20 | 3 (15) |
| 35-44 years | 28 | 5 (18) | 20 | 1 (5) | 50 | 6 (12) |
| 45+ years | 45 | 8 (18) | 25 | 3 (12) | 73 | 12 (16) |
| p value | | 0.927 | | 0.707 | | 0.739 |
| 2022 | | | | | | |
| <25 years | 4 | 0 (0) | 0 | 0 (0) | 4 | 0 (0) |
| 25-34 years | 14 | 0 (0) | 9 | 1 (11) | 23 | 1 (4) |
| 35-44 years | 41 | 4 (10) | 24 | 3 (13) | 65 | 7 (11) |
| 45+ years | 86 | 14 (16) | 53 | 9 (17) | 145 | 23 (16) |
| p value | | 0.358 | | 0.903 | | 0.428 |

Table 6.4.4 HCV RNA prevalence by years since first injection, gender and survey year

| South Australia Years of injection | Male | | Female | | Total | |
|---------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <3 years | 1 | 0 (0) | 2 | 0 (0) | 3 | 0 (0) |
| 3 to 10 years | 10 | 0 (0) | 12 | 1 (8) | 22 | 1 (5) |
| 11+ years | 72 | 19 (26) | 42 | 7 (17) | 114 | 26 (23) |
| p value | | 0.196 | | 0.565 | | 0.074 |
| 2019* | | | | | | |
| <3 years | 2 | 0 (0) | 3 | 0 (0) | 5 | 0 (0) |
| 3 to 10 years | 10 | 0 (0) | 15 | 3 (20) | 25 | 3 (12) |
| 11+ years | 97 | 18 (19) | 62 | 1 (2) | 160 | 19 (12) |
| p value | | 0.291 | | 0.016 | | 0.719 |
| 2020 | | | | | | |
| <3 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 3 to 10 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 11+ years | 8 | 1 (13) | 5 | 0 (0) | 13 | 1 (8) |
| p value | | 0.708 | | -- | | 0.773 |
| 2021 | | | | | | |
| <3 years | 1 | 1 (100) | 1 | 0 (0) | 2 | 1 (50) |
| 3 to 10 years | 8 | 3 (38) | 8 | 1 (13) | 16 | 4 (25) |
| 11+ years | 73 | 11 (15) | 42 | 4 (10) | 120 | 16 (13) |
| p value | | 0.031 | | 0.915 | | 0.183 |
| 2022 | | | | | | |
| <3 years | 9 | 0 (0) | 2 | 0 (0) | 11 | 0 (0) |
| 3 to 10 years | 14 | 1 (7) | 10 | 1 (10) | 25 | 2 (8) |
| 11+ years | 116 | 17 (15) | 71 | 12 (17) | 192 | 29 (15) |
| p value | | 0.675 | | 1.000 | | 0.415 |

Table 6.4.5 HCV RNA prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| South Australia Receptively shared syringe last month | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| No receptive sharing | 70 | 16 (23) | 42 | 6 (14) | 112 | 22 (20) |
| Receptive sharing | 8 | 4 (50) | 11 | 1 (9) | 20 | 5 (25) |
| p value | | 0.127 | | 0.519 | | 0.611 |
| 2019* | | | | | | |
| No receptive sharing | 88 | 15 (17) | 58 | 4 (7) | 147 | 19 (13) |
| Receptive sharing | 12 | 2 (17) | 12 | 0 (0) | 23 | 2 (9) |
| p value | | 0.983 | | 0.347 | | 0.552 |
| 2020 | | | | | | |
| No receptive sharing | 8 | 1 (13) | 5 | 0 (0) | 13 | 1 (8) |
| Receptive sharing | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No receptive sharing | 62 | 16 (26) | 33 | 4 (12) | 98 | 20 (20) |
| Receptive sharing | 7 | 0 (0) | 4 | 0 (0) | 13 | 1 (8) |
| p value | | 0.125 | | 0.461 | | 0.271 |
| 2022 | | | | | | |
| No receptive sharing | 112 | 17 (15) | 64 | 6 (9) | 181 | 23 (13) |
| Receptive sharing | 20 | 1 (5) | 16 | 5 (31) | 37 | 6 (16) |
| p value | | 0.307 | | 0.023 | | 0.567 |

Table 6.4.6 HCV RNA prevalence by last drug injected, gender and survey year

| South Australia Last drug injected | Male | | Female | | Total | |
|---------------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heroin | 28 | 5 (18) | 16 | 2 (13) | 44 | 7 (16) |
| Methamphetamine | 44 | 11 (25) | 35 | 4 (11) | 79 | 15 (19) |
| Other opioids | 8 | 3 (38) | 3 | 2 (67) | 11 | 5 (45) |
| Other drugs | 4 | 1 (25) | 2 | 0 (0) | 6 | 1 (17) |
| p value | | 0.810 | | 0.065 | | 0.254 |
| 2019* | | | | | | |
| Heroin | 24 | 5 (21) | 23 | 2 (9) | 48 | 7 (15) |
| Methamphetamine | 65 | 8 (12) | 43 | 2 (5) | 109 | 10 (9) |
| Other opioids | 18 | 3 (17) | 10 | 1 (10) | 29 | 4 (14) |
| Other drugs | 6 | 2 (33) | 5 | 0 (0) | 11 | 2 (18) |
| p value | | 0.529 | | 0.826 | | 0.671 |
| 2020 | | | | | | |
| Heroin | 3 | 0 (0) | 1 | 0 (0) | 4 | 0 (0) |
| Methamphetamine | 4 | 1 (25) | 3 | 0 (0) | 7 | 1 (14) |
| Other opioids | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) |
| Other drugs | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | 0.704 | | -- | | 0.783 |
| 2021 | | | | | | |
| Heroin | 21 | 5 (24) | 14 | 2 (14) | 37 | 7 (19) |
| Methamphetamine | 51 | 9 (18) | 29 | 1 (3) | 82 | 10 (12) |
| Other opioids | 8 | 1 (13) | 3 | 0 (0) | 12 | 2 (17) |
| Other drugs | 7 | 1 (14) | 4 | 1 (25) | 11 | 2 (18) |
| p value | | 0.875 | | 0.334 | | 0.784 |
| 2022 | | | | | | |
| Heroin | 37 | 5 (14) | 30 | 7 (23) | 69 | 12 (17) |
| Methamphetamine | 70 | 10 (14) | 47 | 4 (9) | 121 | 14 (12) |
| Other opioids | 16 | 0 (0) | 4 | 0 (0) | 21 | 0 (0) |
| Other drugs | 19 | 3 (16) | 5 | 2 (40) | 24 | 5 (21) |
| p value | | 0.444 | | 0.098 | | 0.124 |

Table 6.4.7 HCV RNA prevalence by frequency of drug injection last month, gender and survey year

| South Australia | | Male | | Female | | Total | |
|-----------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Frequency of injection last month | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | |
| 2018* | | | | | | | |
| Less than daily | 40 | 6 (15) | 30 | 3 (10) | 70 | 9 (13) | |
| Daily or more | 42 | 14 (33) | 23 | 4 (17) | 66 | 19 (29) | |
| Not last month | 3 | 1 (33) | 2 | 0 (0) | 5 | 1 (20) | |
| p value | | 0.165 | | 0.618 | | 0.104 | |
| 2019* | | | | | | | |
| Less than daily | 51 | 5 (10) | 43 | 1 (2) | 95 | 6 (6) | |
| Daily or more | 51 | 12 (24) | 29 | 3 (10) | 80 | 15 (19) | |
| Not last month | 12 | 1 (8) | 10 | 1 (10) | 23 | 2 (9) | |
| p value | | 0.128 | | 0.330 | | 0.033 | |
| 2020 | | | | | | | |
| Less than daily | 3 | 0 (0) | 2 | 0 (0) | 5 | 0 (0) | |
| Daily or more | 5 | 1 (20) | 3 | 0 (0) | 8 | 1 (13) | |
| Not last month | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) | |
| p value | | 0.638 | | -- | | 0.668 | |
| 2021 | | | | | | | |
| Less than daily | 37 | 6 (16) | 25 | 2 (8) | 65 | 8 (12) | |
| Daily or more | 32 | 10 (31) | 14 | 2 (14) | 48 | 13 (27) | |
| Not last month | 18 | 0 (0) | 11 | 0 (0) | 29 | 0 (0) | |
| p value | | 0.021 | | 0.426 | | 0.004 | |
| 2022 | | | | | | | |
| Less than daily | 56 | 7 (13) | 33 | 4 (12) | 90 | 11 (12) | |
| Daily or more | 78 | 11 (14) | 47 | 7 (15) | 130 | 18 (14) | |
| Not last month | 10 | 0 (0) | 6 | 2 (33) | 17 | 2 (12) | |
| p value | | 0.678 | | 0.347 | | 0.953 | |

Table 6.4.8 HCV RNA prevalence by imprisonment last year, gender and survey year

| South Australia | | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Imprisonment last year | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | |
| 2018* | | | | | | | |
| No imprisonment | 70 | 14 (20) | 49 | 7 (14) | 119 | 21 (18) | |
| Imprisonment | 8 | 5 (63) | 3 | 0 (0) | 12 | 5 (42) | |
| p value | | 0.010 | | 0.413 | | 0.035 | |
| 2019* | | | | | | | |
| No imprisonment | 86 | 13 (15) | 71 | 5 (7) | 158 | 18 (11) | |
| Imprisonment | 17 | 2 (12) | 8 | 0 (0) | 24 | 2 (8) | |
| p value | | 0.750 | | 0.440 | | 0.639 | |
| 2020 | | | | | | | |
| No imprisonment | 9 | 1 (11) | 5 | 0 (0) | 14 | 1 (7) | |
| Imprisonment | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| p value | | -- | | -- | | -- | |
| 2021 | | | | | | | |
| No imprisonment | 71 | 11 (15) | 46 | 2 (4) | 122 | 14 (11) | |
| Imprisonment | 9 | 0 (0) | 5 | 0 (0) | 14 | 0 (0) | |
| p value | | 0.204 | | 0.634 | | 0.181 | |
| 2022 | | | | | | | |
| No imprisonment | 119 | 15 (13) | 80 | 13 (16) | 203 | 28 (14) | |
| Imprisonment | 26 | 3 (12) | 6 | 0 (0) | 33 | 3 (9) | |
| p value | | 1.000 | | 0.585 | | 0.586 | |

Table 6.4.9 HCV RNA prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| South Australia | | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Aboriginal and Torres Strait Islander origin | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | |
| 2018* | | | | | | | |
| Non Indigenous | 72 | 15 (21) | 47 | 8 (17) | 119 | 23 (19) | |
| Indigenous | 14 | 5 (36) | 7 | 0 (0) | 21 | 5 (24) | |
| p value | | 0.204 | | 0.205 | | 0.554 | |
| 2019* | | | | | | | |
| Non Indigenous | 89 | 14 (16) | 67 | 4 (6) | 157 | 18 (11) | |
| Indigenous | 26 | 4 (15) | 15 | 1 (7) | 42 | 5 (12) | |
| p value | | 0.956 | | 0.981 | | 0.962 | |
| 2020 | | | | | | | |
| Non Indigenous | 7 | 1 (14) | 5 | 0 (0) | 12 | 1 (8) | |
| Indigenous | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) | |
| p value | | 0.571 | | -- | | 0.672 | |
| 2021 | | | | | | | |
| Non Indigenous | 55 | 9 (16) | 34 | 4 (12) | 92 | 13 (14) | |
| Indigenous | 25 | 5 (20) | 17 | 0 (0) | 44 | 6 (14) | |
| p value | | 0.692 | | 0.141 | | 0.938 | |
| 2022 | | | | | | | |
| Non Indigenous | 106 | 15 (14) | 63 | 11 (17) | 174 | 26 (15) | |
| Indigenous | 42 | 4 (10) | 23 | 2 (9) | 66 | 6 (9) | |
| p value | | 0.590 | | 0.499 | | 0.234 | |

Table 6.4.10 HCV RNA prevalence by main language spoken at home by parents, gender and survey year

| South Australia | | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Main language spoken at home by parents | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | |
| 2018* | | | | | | | |
| English speaking | 74 | 18 (24) | 54 | 8 (15) | 129 | 26 (20) | |
| Non-English speaking | 11 | 3 (27) | 2 | 0 (0) | 13 | 3 (23) | |
| p value | | 0.903 | | 0.564 | | 0.824 | |
| 2019* | | | | | | | |
| English speaking | 112 | 18 (16) | 84 | 5 (6) | 197 | 23 (12) | |
| Non-English speaking | 4 | 0 (0) | 1 | 0 (0) | 5 | 0 (0) | |
| p value | | 0.383 | | 0.803 | | 0.416 | |
| 2020 | | | | | | | |
| English speaking | 9 | 1 (11) | 5 | 0 (0) | 14 | 1 (7) | |
| Non-English speaking | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| p value | | -- | | -- | | -- | |
| 2021 | | | | | | | |
| English speaking | 84 | 15 (18) | 51 | 4 (8) | 140 | 20 (14) | |
| Non-English speaking | 2 | 1 (50) | 3 | 1 (33) | 5 | 2 (40) | |
| p value | | 0.248 | | 0.139 | | 0.115 | |
| 2022 | | | | | | | |
| English speaking | 138 | 16 (12) | 81 | 13 (16) | 226 | 29 (13) | |
| Non-English speaking | 9 | 3 (33) | 4 | 0 (0) | 13 | 3 (23) | |
| p value | | 0.093 | | 1.000 | | 0.391 | |

Table 6.4.11 HCV RNA prevalence by region/country of birth, gender and survey year

| South Australia | | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | |
| 2018* | | | | | | | |
| Australia | 73 | 18 (25) | 52 | 8 (15) | 125 | 26 (21) | |
| Other Oceania | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) | |
| Asia | 3 | 1 (33) | 0 | 0 (0) | 3 | 1 (33) | |
| UK & Ireland | 5 | 0 (0) | 2 | 0 (0) | 7 | 0 (0) | |
| Other | 3 | 2 (67) | 1 | 0 (0) | 4 | 2 (50) | |
| p value | | 0.415 | | 0.831 | | 0.332 | |
| 2019* | | | | | | | |
| Australia | 104 | 16 (15) | 74 | 4 (5) | 179 | 20 (11) | |
| Other Oceania | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) | |
| Asia | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) | |
| UK & Ireland | 8 | 2 (25) | 7 | 1 (14) | 15 | 3 (20) | |
| Other | 2 | 0 (0) | 2 | 0 (0) | 4 | 0 (0) | |
| p value | | 0.868 | | 0.846 | | 0.688 | |
| 2020 | | | | | | | |
| Australia | 8 | 1 (13) | 4 | 0 (0) | 12 | 1 (8) | |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| UK & Ireland | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) | |
| Other | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| p value | | 0.708 | | -- | | 0.672 | |
| 2021 | | | | | | | |
| Australia | 76 | 13 (17) | 45 | 2 (4) | 126 | 16 (13) | |
| Other Oceania | 0 | 0 (0) | 2 | 1 (50) | 2 | 1 (50) | |
| Asia | 1 | 1 (100) | 1 | 1 (100) | 2 | 2 (100) | |
| UK & Ireland | 5 | 2 (40) | 3 | 1 (33) | 8 | 3 (38) | |
| Other | 1 | 0 (0) | 2 | 0 (0) | 3 | 0 (0) | |
| p value | | 0.110 | | 0.002 | | 0.002 | |
| 2022 | | | | | | | |
| Australia | 127 | 14 (11) | 72 | 11 (15) | 206 | 25 (12) | |
| Other Oceania | 2 | 0 (0) | 2 | 1 (50) | 4 | 1 (25) | |
| Asia | 5 | 1 (20) | 2 | 0 (0) | 7 | 1 (14) | |
| UK & Ireland | 5 | 1 (20) | 8 | 1 (13) | 13 | 2 (15) | |
| Other | 3 | 1 (33) | 2 | 0 (0) | 5 | 1 (20) | |
| p value | | 0.288 | | 0.680 | | 0.560 | |

Tasmania

Table 7.1.1 Number (percentage) of respondents by demographic characteristics and survey year

| Tasmania | 2018 | 2019 | 2020* | 2021* | 2022 |
|---|---------|----------|----------|---------|---------|
| Number of sites | 4 | 4 | 4 | 4 | 4 |
| N° surveyed | N=62 | N=40 | N=35 | N=95 | N=91 |
| Response rate (%) | 23% | 17% | 17% | 62% | 38% |
| Gender (%) | | | | | |
| Male | 40 (65) | 21 (53) | 21 (60) | 51 (54) | 60 (66) |
| Female | 21 (34) | 19 (48) | 14 (40) | 44 (46) | 29 (32) |
| Other | 1 (2) | 0 (0) | 0 (0) | 0 (0) | 2 (2) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Sexual identity (%) | | | | | |
| Heterosexual | 52 (84) | 35 (88) | 32 (91) | 81 (85) | 77 (85) |
| Bisexual | 5 (8) | 4 (10) | 1 (3) | 9 (9) | 9 (10) |
| Gay/Lesbian | 0 (0) | 0 (0) | 2 (6) | 1 (1) | 3 (3) |
| Other | -- -- | -- -- | 0 (0) | 2 (2) | 2 (2) |
| Not reported | 5 (8) | 1 (3) | 0 (0) | 2 (2) | 0 (0) |
| Aboriginal and Torres Strait Islander origin (%) | | | | | |
| No | 49 (79) | 36 (90) | 31 (89) | 72 (76) | 63 (69) |
| Yes | 11 (18) | 4 (10) | 4 (11) | 20 (21) | 25 (27) |
| Not reported | 2 (3) | 0 (0) | 0 (0) | 3 (3) | 3 (3) |
| Region/country of birth (%) | | | | | |
| Australia | 58 (94) | 39 (98) | 33 (94) | 86 (91) | 87 (96) |
| Other Oceania | 1 (2) | 0 (0) | 0 (0) | 0 (0) | 1 (1) |
| Asia | 0 (0) | 0 (0) | 0 (0) | 1 (1) | 0 (0) |
| UK & Ireland | 2 (3) | 1 (3) | 2 (6) | 0 (0) | 1 (1) |
| Other | 1 (2) | 0 (0) | 0 (0) | 6 (6) | 1 (1) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 2 (2) | 1 (1) |
| Main language spoken at home by parents (%) | | | | | |
| English | 61 (98) | 40 (100) | 35 (100) | 90 (95) | 90 (99) |
| Non-English | 1 (2) | 0 (0) | 0 (0) | 4 (4) | 1 (1) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 1 (1) | 0 (0) |
| Age and time since first injection (years) | | | | | |
| <i>Median age</i> | 41.5 | 42 | 43 | 45 | 44 |
| <i>Age range</i> | 21-59 | 24-59 | 28-66 | 20-67 | 24-62 |
| Age group (%) | | | | | |
| <25 years | 2 (3) | 1 (3) | 0 (0) | 2 (2) | 1 (1) |
| 25+ years | 60 (97) | 39 (98) | 35 (100) | 91 (96) | 90 (99) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 2 (2) | 0 (0) |
| Median age first injection | | | | | |
| <i>Median age first injection</i> | 19 | 20 | 20 | 20 | 19 |
| <i>Age range</i> | 13-48 | 13-38 | 13-36 | 12-55 | 10-56 |
| <i>N° not reported</i> | 2 | 0 | 0 | 2 | 2 |
| Median yrs since first injection | | | | | |
| <i>Median yrs since first injection</i> | 20 | 22 | 23 | 21.5 | 25 |
| <i>Range</i> | 1-40 | 5-45 | 1-46 | 1-48 | 2-46 |
| Years since first injection | | | | | |
| <3 years | 1 (2) | 0 (0) | 1 (3) | 5 (5) | 2 (2) |
| 3+ years | 59 (95) | 40 (100) | 34 (97) | 87 (92) | 87 (96) |
| Not reported | 2 (3) | 0 (0) | 0 (0) | 3 (3) | 2 (2) |
| Imprisonment last year (%) | | | | | |
| No | 56 (90) | 36 (90) | 30 (86) | 82 (86) | 75 (82) |
| Yes | 3 (5) | 4 (10) | 5 (14) | 8 (8) | 14 (15) |
| Not reported | 3 (5) | 0 (0) | 0 (0) | 5 (5) | 2 (2) |
| N° in prison | | | | | |
| <i>N° in prison</i> | N=3 | N=4 | N=5 | N=8 | N=14 |
| <i>Injected in prison</i> | 2 (67) | 0 (0) | 0 (0) | 2 (25) | 4 (29) |

* Implementation of the 2020 and 2021 ANSPS was impacted by the COVID-19 pandemic, see page 1 for more information

Table 7.1.2 Number (percentage) of respondents by last drug injected, frequency of injecting and survey year

| Tasmania | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|---------|---------|---------|---------|---------|
| N° surveyed | N=62 | N=40 | N=35 | N=95 | N=91 |
| Drug last injected (%) | | | | | |
| Cocaine | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Methamphetamine | 22 (35) | 18 (45) | 15 (43) | 49 (52) | 50 (55) |
| Heroin | 2 (3) | 2 (5) | 1 (3) | 7 (7) | 3 (3) |
| Pharm. opioids | 19 (31) | 3 (8) | 9 (26) | 14 (15) | 21 (23) |
| Methadone | 12 (19) | 11 (28) | 7 (20) | 9 (9) | 6 (7) |
| Buprenorphine | 2 (3) | 2 (5) | 0 (0) | 4 (4) | 6 (7) |
| Buprenorphine/naloxone | 2 (3) | 1 (3) | 0 (0) | 4 (4) | 1 (1) |
| PIEDs | 0 (0) | 1 (3) | 1 (3) | 2 (2) | 0 (0) |
| More than one | 1 (2) | 0 (0) | 1 (3) | 3 (3) | 3 (3) |
| Other | 2 (3) | 2 (5) | 0 (0) | 2 (2) | 1 (1) |
| Not reported | 0 (0) | 0 (0) | 1 (3) | 1 (1) | 0 (0) |
| Ever injected opioids | | | | | |
| No | 3 (5) | 3 (8) | 2 (6) | 32 (34) | 16 (18) |
| Yes | 59 (95) | 37 (93) | 33 (94) | 63 (66) | 75 (82) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Frequency of injection last month (%) | | | | | |
| Not last month | 2 (3) | 3 (8) | 5 (14) | 10 (11) | 2 (2) |
| Less than weekly | 7 (11) | 8 (20) | 7 (20) | 32 (34) | 14 (15) |
| Weekly not daily | 27 (44) | 17 (43) | 12 (34) | 20 (21) | 26 (29) |
| Daily or more | 26 (42) | 11 (28) | 11 (31) | 31 (33) | 49 (54) |
| Not reported | 0 (0) | 1 (3) | 0 (0) | 2 (2) | 0 (0) |
| Present when someone injected for the first time in the last 12 months (%) | | | | | |
| No | 57 (92) | 38 (95) | 34 (97) | 78 (82) | 75 (82) |
| Yes | 4 (6) | 2 (5) | 1 (3) | 14 (15) | 16 (18) |
| Not reported | 1 (2) | 0 (0) | 0 (0) | 3 (3) | 0 (0) |
| Overdosed in last 12 months | | | | | |
| No | 55 (89) | 31 (78) | 30 (86) | 85 (89) | 83 (91) |
| Yes | 4 (6) | 9 (23) | 5 (14) | 5 (5) | 8 (9) |
| Not reported | 3 (5) | 0 (0) | 0 (0) | 5 (5) | 0 (0) |

Table 7.1.3 Number (percentage) of respondents by injecting behaviour in the last month and survey year

| Tasmania | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|-------------|-------------|-------------|-------------|-------------|
| N° injected last month | N=60 | N=36 | N=30 | N=83 | N=89 |
| Places injected last month (%) | | | | | |
| All private | 35 (58) | 24 (67) | 24 (80) | 62 (75) | 49 (55) |
| Any public | 25 (42) | 12 (33) | 6 (20) | 21 (25) | 40 (45) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Use of new and sterile needles and syringes last month (%) | | | | | |
| All injections | 47 (78) | 30 (83) | 26 (87) | 71 (86) | 75 (84) |
| Most of the time | 9 (15) | 3 (8) | 4 (13) | 8 (10) | 12 (13) |
| Half of the time | 1 (2) | 1 (3) | 0 (0) | 0 (0) | 1 (1) |
| Some of the time | 1 (2) | 0 (0) | 0 (0) | 0 (0) | 1 (1) |
| Not last month | 2 (3) | 2 (6) | 0 (0) | 2 (2) | 0 (0) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 2 (2) | 0 (0) |
| Re-used someone else's used needle & syringe last month (%) | | | | | |
| None | 56 (93) | 34 (94) | 30 (100) | 76 (92) | 76 (85) |
| Once | 0 (0) | 1 (3) | 0 (0) | 2 (2) | 3 (3) |
| Twice | 1 (2) | 0 (0) | 0 (0) | 4 (5) | 6 (7) |
| 3-5 times | 1 (2) | 1 (3) | 0 (0) | 0 (0) | 1 (1) |
| >5 times | 2 (3) | 0 (0) | 0 (0) | 1 (1) | 3 (3) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Number of people needle & syringe was re-used after last month (%) | | | | | |
| None | 56 (93) | 34 (94) | 30 (100) | 76 (92) | 76 (85) |
| One | 1 (2) | 0 (0) | 0 (0) | 2 (2) | 3 (3) |
| Two | 1 (2) | 0 (0) | 0 (0) | 1 (1) | 1 (1) |
| Three to five | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 2 (2) |
| More than five | 1 (2) | 0 (0) | 0 (0) | 0 (0) | 1 (1) |
| Don't know | 0 (0) | 0 (0) | 0 (0) | 3 (4) | 4 (4) |
| Not reported | 1 (2) | 2 (6) | 0 (0) | 1 (1) | 2 (2) |
| Relationship to people needle & syringe was used after last month (%) [more than one could be selected] | | | | | |
| Regular sex partner | 1 (2) | 0 (0) | 0 (0) | 2 (2) | 2 (2) |
| Casual sex partner | 1 (2) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Close friend | 2 (3) | 0 (0) | 0 (0) | 2 (2) | 3 (3) |
| Acquaintance | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 2 (2) |
| Other | 0 (0) | 0 (0) | 0 (0) | 1 (1) | 3 (3) |
| Equipment used after someone else last month (%) [more than one could be selected] | | | | | |
| Spoon | 5 (8) | 3 (8) | 0 (0) | 11 (13) | 20 (22) |
| Water | 5 (8) | 3 (8) | 1 (3) | 19 (23) | 13 (15) |
| Filter | 2 (3) | 4 (11) | 0 (0) | 9 (11) | 5 (6) |
| Drug mix | 4 (7) | 1 (3) | 0 (0) | 5 (6) | 4 (4) |
| None | 52 (87) | 26 (72) | 28 (93) | 58 (70) | 54 (61) |
| Injected by someone after that person injected themselves or others last month (%) | | | | | |
| No | 53 (88) | 35 (97) | 28 (93) | 73 (88) | 71 (80) |
| Yes | 7 (12) | 1 (3) | 2 (7) | 9 (11) | 18 (20) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 1 (1) | 0 (0) |
| Source of needle acquisition | | | | | |
| Needle Syringe Program | 57 (95) | 36 (100) | 29 (97) | 72 (87) | 86 (97) |
| Chemist/Pharmacy | 8 (13) | 6 (17) | 8 (27) | 19 (23) | 27 (30) |
| Personal sources | 2 (3) | 2 (6) | 0 (0) | 5 (6) | 12 (13) |
| Dispensing/Vending Machine | 9 (15) | 8 (22) | 7 (23) | 7 (8) | 19 (21) |
| Other sources | 2 (3) | 0 (0) | 0 (0) | 1 (1) | 0 (0) |

Table 7.1.4 Number (percentage) of respondents by treatment for drug use and survey year

| Tasmania | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|---------|---------|---------|---------|---------|
| N° surveyed | N=62 | N=40 | N=35 | N=95 | N=91 |
| Ever any treatment/therapy for drug use (%) | | | | | |
| No | 12 (19) | 10 (25) | 11 (31) | 33 (35) | 23 (25) |
| Yes | 50 (81) | 30 (75) | 24 (69) | 62 (65) | 68 (75) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| History of methadone maintenance treatment (%) | | | | | |
| Currently | 16 (26) | 10 (25) | 7 (20) | 10 (11) | 12 (13) |
| Previously | 24 (39) | 11 (28) | 5 (14) | 18 (19) | 24 (26) |
| Never | 20 (32) | 19 (48) | 23 (66) | 62 (65) | 55 (60) |
| Not reported | 2 (3) | 0 (0) | 0 (0) | 5 (5) | 0 (0) |
| History of buprenorphine (Subutex®) treatment (%) | | | | | |
| Currently | 6 (10) | 5 (13) | 2 (6) | 12 (13) | 14 (15) |
| Previously | 19 (31) | 6 (15) | 9 (26) | 16 (17) | 16 (18) |
| Never | 35 (56) | 29 (73) | 24 (69) | 64 (67) | 59 (65) |
| Not reported | 2 (3) | 0 (0) | 0 (0) | 3 (3) | 2 (2) |
| History of buprenorphine-naloxone (Suboxone®) treatment (%) | | | | | |
| Currently | 4 (6) | 2 (5) | 1 (3) | 6 (6) | 9 (10) |
| Previously | 19 (31) | 6 (15) | 9 (26) | 16 (17) | 16 (18) |
| Never | 37 (60) | 32 (80) | 24 (69) | 71 (75) | 66 (73) |
| Not reported | 2 (3) | 0 (0) | 1 (3) | 2 (2) | 0 (0) |

Table 7.1.5 Number (percentage) of respondents by testing for HIV and HCV, HCV treatment by survey year

| Tasmania | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|---------|---------|---------|---------|---------|
| N° surveyed | N=62 | N=40 | N=35 | N=95 | N=91 |
| Previous HIV test (%) | | | | | |
| Yes, ever | 49 (79) | 33 (83) | 29 (83) | 71 (75) | 77 (85) |
| Yes, last year | 21 (34) | 14 (35) | 15 (43) | 40 (42) | 50 (55) |
| >1 year ago | 28 (45) | 19 (48) | 14 (40) | 31 (33) | 27 (30) |
| Never tested | 9 (15) | 6 (15) | 6 (17) | 19 (20) | 12 (13) |
| Not reported | 4 (6) | 1 (3) | 0 (0) | 5 (5) | 2 (2) |
| Previous HCV test (%) | | | | | |
| Yes, ever | 53 (85) | 36 (90) | 29 (83) | 79 (83) | 75 (82) |
| Yes, last year | 35 (56) | 22 (55) | 18 (51) | 52 (55) | 52 (57) |
| >1 year ago | 18 (29) | 14 (35) | 11 (31) | 27 (28) | 23 (25) |
| Never tested | 5 (8) | 1 (3) | 3 (9) | 7 (7) | 6 (7) |
| Not reported | 4 (6) | 3 (8) | 3 (9) | 9 (9) | 10 (11) |
| Lifetime treatment for HCV (%)# | | | | | |
| N° self-reported HCV diagnosis | N=21 | N=11 | N=4 | N=19 | N=17 |
| Antiviral treatment | 15 (71) | 6 (55) | 3 (75) | 11 (42) | 15 (88) |
| No antiviral treatment | 6 (29) | 5 (45) | 1 (25) | 8 (58) | 2 (12) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| Treatment for HCV in past 12 months (%)#* | | | | | |
| N° self-reported HCV diagnosis | N=17 | N=8 | N=3 | N=9 | N=7 |
| Antiviral treatment | 9 (53) | 2 (25) | 1 (33) | 1 (11) | 4 (57) |
| No antiviral treatment | 8 (47) | 6 (75) | 2 (67) | 8 (89) | 3 (43) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |

among people who tested HCV antibody positive and did not report spontaneous clearance

* excludes people who reported treatment induced clearance more than 12 months previously

Table 7.1.6 Number (percentage) of respondents by sexual behaviour in the month prior to survey and survey year

| Tasmania | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|---------|---------|---------|---------|---------|
| N° surveyed | N=62 | N=40 | N=35 | N=95 | N=91 |
| Sex with a regular partner last month (%) | | | | | |
| No | 32 (52) | 15 (38) | 10 (29) | 34 (36) | 43 (47) |
| Yes | 26 (42) | 11 (28) | 11 (31) | 30 (32) | 36 (40) |
| Not reported | 4 (6) | 14 (35) | 14 (40) | 31 (33) | 12 (13) |
| Condom used with regular partner last month (%) | | | | | |
| N° surveyed | N=26 | N=11 | N=11 | N=30 | N=36 |
| Never | 21 (81) | 9 (82) | 9 (82) | 23 (77) | 30 (83) |
| Sometimes | 4 (15) | 1 (9) | 1 (9) | 3 (10) | 2 (6) |
| Every time | 0 (0) | 1 (9) | 1 (9) | 1 (3) | 4 (11) |
| Not reported | 1 (4) | 0 (0) | 0 (0) | 3 (10) | 0 (0) |
| Sex with other partner(s) last month (%) | | | | | |
| No | 51 (82) | 24 (60) | 21 (60) | 57 (60) | 69 (76) |
| Yes | 6 (10) | 2 (5) | 1 (3) | 5 (5) | 9 (10) |
| Not reported | 5 (8) | 14 (35) | 13 (37) | 33 (35) | 13 (14) |
| Condom used with other partner(s) last month (%) | | | | | |
| N° surveyed | N=6 | N=2 | N=1 | N=5 | N=9 |
| Never | 4 (67) | 2 (100) | 1 (100) | 3 (60) | 4 (44) |
| Sometimes | 2 (33) | 0 (0) | 0 (0) | 1 (20) | 3 (33) |
| Every time | 0 (0) | 0 (0) | 0 (0) | 0 (0) | 2 (22) |
| Not reported | 0 (0) | 0 (0) | 0 (0) | 1 (20) | 0 (0) |
| Sex work last month (%) | | | | | |
| No | 55 (89) | 26 (65) | 22 (63) | 57 (60) | 72 (79) |
| Yes | 1 (2) | 0 (0) | 0 (0) | 6 (6) | 6 (7) |
| Not reported | 6 (10) | 14 (35) | 13 (37) | 32 (34) | 13 (14) |
| Condom used at last sex work (%) | | | | | |
| Yes | 0 (0) | 0 (0) | 0 (0) | 3 (50) | 4 (67) |

Table 7.1.7 Number (percentage) of respondents by COVID-19 related impacts, 2020-2022

| Tasmania | 2020 | 2021 | 2022 |
|---|---------|---------|---------|
| N° surveyed | N=35 | N=95 | N=91 |
| COVID-19 test | | | |
| Yes | 5 (14) | 22 (23) | 83 (91) |
| No | 25 (71) | 54 (57) | 6 (7) |
| Don't know | 2 (6) | 2 (2) | -- -- |
| Not reported | 3 (9) | 17 (18) | 2 (2) |
| Self-reported positive COVID-19 test result | | | |
| Yes | 0 (0) | 0 (0) | 44 (53) |
| Yes, last 12 months | -- -- | -- -- | 39 (47) |
| Yes, more than a year ago | -- -- | -- -- | 5 (6) |
| COVID-19 vaccine uptake | | | |
| ≥1 dose | -- -- | 47 (49) | 76 (84) |
| COVID-19 vaccine doses | | | |
| One dose | -- -- | 16 (17) | 7 (8) |
| Two doses | -- -- | 31 (33) | 32 (35) |
| Three or more doses | -- -- | -- -- | 37 (41) |
| No doses | -- -- | 34 (36) | 13 (14) |
| Not reported | -- -- | 14 (15) | 2 (2) |
| Vaccine type * | | | N=76 |
| Astra Zeneca | -- -- | 5 (5) | 16 (21) |
| Pfizer | -- -- | 22 (23) | 44 (58) |
| Don't know | -- -- | 5 (5) | 9 (12) |
| Other | -- -- | -- -- | 13 (17) |
| Not reported | -- -- | 63 (66) | -- -- |
| * 2021 only one response permitted, 2022 multiple responses permitted | | | |
| Last vaccine location | | | N=91 |
| GP or medical centre | -- -- | -- -- | 22 (24) |
| Pharmacy/ chemist | -- -- | -- -- | 26 (29) |
| Vaccine hub | -- -- | -- -- | 13 (14) |
| Opioid treatment/ methadone clinic | -- -- | -- -- | 3 (3) |
| NSP | -- -- | -- -- | 0 (0) |
| Prison | -- -- | -- -- | 8 (9) |
| Other | -- -- | -- -- | 2 (2) |
| Multiple response | -- -- | -- -- | 3 (3) |
| Not reported | -- -- | -- -- | 14 (15) |
| Vaccine hesitancy - if not had three or more doses | | | |
| Haven't got around to it | -- -- | -- -- | 21 (23) |
| Difficulty booking appointment | -- -- | -- -- | 3 (3) |
| Concerns about side effects | -- -- | -- -- | 8 (9) |
| Concerns about vaccine safety | -- -- | -- -- | 6 (7) |
| Not at risk of serious illness | -- -- | -- -- | 4 (4) |
| Other | -- -- | -- -- | 5 (5) |
| Multiple response | -- -- | -- -- | 2 (2) |
| Not reported | -- -- | -- -- | 42 (46) |

HIV antibody prevalence

Table 7.2.1 HIV antibody prevalence by gender and survey year

| Tasmania Survey year | Male | | Female | | Total | |
|-------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | 40 | 0 (0.0) | 21 | 0 (0.0) | 62 | 0 (0.0) |
| 2019 | 20 | 1 (5.0) | 18 | 0 (0.0) | 38 | 1 (2.6) |
| 2020 | 20 | 0 (0.0) | 14 | 0 (0.0) | 34 | 0 (0.0) |
| 2021 | 49 | 0 (0.0) | 42 | 0 (0.0) | 91 | 0 (0.0) |
| 2022 | 59 | 0 (0.0) | 29 | 0 (0.0) | 90 | 0 (0.0) |
| X ² p trend | | 0.375 | | -- | | 0.366 |

HCV antibody prevalence

Table 7.3.1 HCV antibody prevalence by gender and survey year

| Tasmania Survey year | Male | | Female | | Total | |
|-------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | 40 | 21 (53) | 21 | 12 (57) | 62 | 34 (55) |
| 2019 | 19 | 7 (37) | 18 | 6 (33) | 37 | 13 (35) |
| 2020 | 20 | 6 (30) | 14 | 3 (21) | 34 | 9 (26) |
| 2021 | 49 | 18 (37) | 42 | 9 (21) | 91 | 27 (30) |
| 2022 | 59 | 21 (36) | 29 | 6 (21) | 90 | 28 (31) |
| X ² p trend | | 0.128 | | 0.004 | | 0.004 |

Table 7.3.2 HCV antibody prevalence by sexual identity, gender and survey year

| Tasmania Sexual identity | Male | | Female | | Total | |
|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| Heterosexual | 34 | 18 (53) | 18 | 9 (50) | 52 | 27 (52) |
| Bisexual | 2 | 0 (0) | 2 | 2 (100) | 5 | 3 (60) |
| Homosexual | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | 0.486 | | 0.479 | | 1.000 |
| 2019 | | | | | | |
| Heterosexual | 18 | 7 (39) | 14 | 6 (43) | 32 | 13 (41) |
| Bisexual | 1 | 0 (0) | 3 | 0 (0) | 4 | 0 (0) |
| Homosexual | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | 1.000 | | 0.515 | | 0.274 |
| 2020 | | | | | | |
| Heterosexual | 19 | 6 (32) | 12 | 3 (25) | 31 | 9 (29) |
| Bisexual | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| Homosexual | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2021 | | | | | | |
| Heterosexual | 44 | 16 (36) | 33 | 6 (18) | 77 | 22 (29) |
| Bisexual | 3 | 0 (0) | 6 | 2 (33) | 9 | 2 (22) |
| Homosexual | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| p value | | 0.219 | | 0.260 | | 0.147 |
| 2022 | | | | | | |
| Heterosexual | 56 | 20 (36) | 19 | 2 (11) | 76 | 23 (30) |
| Bisexual | 2 | 1 (50) | 7 | 2 (29) | 9 | 3 (33) |
| Homosexual | 1 | 0 (0) | 2 | 1 (50) | 3 | 1 (33) |
| p value | | 1.000 | | 0.085 | | 0.733 |

Table 7.3.3 HCV antibody prevalence by age group, gender and survey year

| Tasmania Age group | Male | | Female | | Total | |
|-----------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <25 years | 1 | 1 (100) | 0 | 0 (0) | 2 | 2 (100) |
| 25-34 years | 6 | 3 (50) | 3 | 2 (67) | 9 | 5 (56) |
| 35-44 years | 19 | 8 (42) | 12 | 8 (67) | 31 | 16 (52) |
| 45+ years | 14 | 9 (64) | 6 | 2 (33) | 20 | 11 (55) |
| p value | | 0.490 | | 0.475 | | 0.798 |
| 2019 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 1 | 0 (0) | 4 | 2 (50) | 5 | 2 (40) |
| 35-44 years | 12 | 3 (25) | 7 | 1 (14) | 19 | 4 (21) |
| 45+ years | 6 | 4 (67) | 7 | 3 (43) | 13 | 7 (54) |
| p value | | 0.129 | | 0.541 | | 0.178 |
| 2020 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 4 | 0 (0) | 5 | 1 (20) | 9 | 1 (11) |
| 35-44 years | 5 | 2 (40) | 3 | 1 (33) | 8 | 3 (38) |
| 45+ years | 11 | 4 (36) | 6 | 1 (17) | 17 | 5 (29) |
| p value | | 0.532 | | 1.000 | | 0.475 |
| 2021 | | | | | | |
| <25 years | 0 | 0 (0) | 2 | 0 (0) | 2 | 0 (0) |
| 25-34 years | 7 | 2 (29) | 7 | 1 (14) | 14 | 3 (21) |
| 35-44 years | 15 | 5 (33) | 12 | 1 (8) | 27 | 6 (22) |
| 45+ years | 27 | 11 (41) | 19 | 6 (32) | 46 | 17 (37) |
| p value | | 0.847 | | 0.489 | | 0.437 |
| 2022 | | | | | | |
| <25 years | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| 25-34 years | 11 | 2 (18) | 6 | 0 (0) | 17 | 2 (12) |
| 35-44 years | 17 | 7 (41) | 10 | 2 (20) | 28 | 10 (36) |
| 45+ years | 31 | 12 (39) | 12 | 4 (33) | 44 | 16 (36) |
| p value | | 0.418 | | 0.494 | | 0.200 |

Table 7.3.4 HCV antibody prevalence by years since first injection, gender and survey year

| Tasmania Years of injection | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <3 years | 1 | 1 (100) | 0 | 0 (0) | 1 | 1 (100) |
| 3 to 10 years | 5 | 2 (40) | 3 | 1 (33) | 9 | 4 (44) |
| 11+ years | 33 | 17 (52) | 17 | 11 (65) | 50 | 28 (56) |
| p value | | 1.000 | | 0.537 | | 0.845 |
| 2019 | | | | | | |
| <3 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 3 to 10 years | 2 | 0 (0) | 4 | 1 (25) | 6 | 1 (17) |
| 11+ years | 17 | 7 (41) | 14 | 5 (36) | 31 | 12 (39) |
| p value | | 0.509 | | 1.000 | | 0.394 |
| 2020 | | | | | | |
| <3 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 3 to 10 years | 2 | 0 (0) | 4 | 1 (25) | 6 | 1 (17) |
| 11+ years | 17 | 6 (35) | 10 | 2 (20) | 27 | 8 (30) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2021 | | | | | | |
| <3 years | 2 | 0 (0) | 3 | 0 (0) | 5 | 0 (0) |
| 3 to 10 years | 2 | 0 (0) | 5 | 0 (0) | 7 | 0 (0) |
| 11+ years | 45 | 18 (40) | 31 | 8 (26) | 76 | 26 (34) |
| p value | | 0.394 | | 0.607 | | 0.067 |
| 2022 | | | | | | |
| <3 years | 0 | 0 (0) | 2 | 0 (0) | 2 | 0 (0) |
| 3 to 10 years | 3 | 0 (0) | 7 | 0 (0) | 10 | 0 (0) |
| 11+ years | 55 | 20 (36) | 19 | 6 (32) | 76 | 27 (36) |
| p value | | 0.544 | | 0.226 | | 0.029 |

Table 7.3.5 HCV antibody prevalence by re-use of someone else's used needle and syringe last month, years since first injection and survey year

| Tasmania Receptively shared syringe last month | <3 years | | 3+ years | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| No receptive sharing | 1 | 1 (100) | 53 | 30 (57) | 56 | 32 (57) |
| Receptive sharing | 0 | 0 (0) | 4 | 1 (25) | 4 | 1 (25) |
| p value | | -- | | 0.322 | | 0.318 |
| 2019 | | | | | | |
| No receptive sharing | 0 | 0 (0) | 32 | 13 (41) | 32 | 13 (41) |
| Receptive sharing | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | 1.000 | | 1.000 |
| 2020 | | | | | | |
| No receptive sharing | 1 | 0 (0) | 28 | 9 (32) | 29 | 9 (31) |
| Receptive sharing | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No receptive sharing | 5 | 0 (0) | 68 | 22 (32) | 75 | 22 (29) |
| Receptive sharing | 0 | 0 (0) | 5 | 2 (40) | 5 | 2 (40) |
| p value | | -- | | 1.000 | | 0.633 |
| 2022 | | | | | | |
| No receptive sharing | 2 | 0 (0) | 71 | 21 (30) | 75 | 22 (29) |
| Receptive sharing | 0 | 0 (0) | 13 | 5 (38) | 13 | 5 (38) |
| p value | | -- | | 0.524 | | 0.510 |

Table 7.3.6 HCV antibody prevalence by last drug injected, years since first injection and survey year

| Tasmania Last drug injected | <3 years | | 3+ years | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Methamphetamine | 0 | 0 (0) | 21 | 10 (48) | 22 | 10 (45) |
| Pharmaceutical opioids | 1 | 1 (100) | 17 | 10 (59) | 19 | 12 (63) |
| OAT* | 0 | 0 (0) | 16 | 9 (56) | 16 | 9 (56) |
| Other drugs | 0 | 0 (0) | 5 | 3 (60) | 5 | 3 (60) |
| p value | | -- | | 0.895 | | 0.710 |
| 2019 | | | | | | |
| Methamphetamine | 0 | 0 (0) | 16 | 6 (38) | 16 | 6 (38) |
| Pharmaceutical opioids | 0 | 0 (0) | 3 | 0 (0) | 3 | 0 (0) |
| OAT* | 0 | 0 (0) | 14 | 6 (43) | 14 | 6 (43) |
| Other drugs | 0 | 0 (0) | 3 | 1 (33) | 3 | 1 (33) |
| p value | | -- | | 0.573 | | 0.573 |
| 2020 | | | | | | |
| Methamphetamine | 1 | 0 (0) | 13 | 2 (15) | 14 | 2 (14) |
| Pharmaceutical opioids | 0 | 0 (0) | 9 | 3 (33) | 9 | 3 (33) |
| OAT* | 0 | 0 (0) | 7 | 3 (43) | 7 | 3 (43) |
| Other drugs | 0 | 0 (0) | 4 | 1 (25) | 4 | 1 (25) |
| p value | | -- | | 0.581 | | 0.521 |
| 2021 | | | | | | |
| Methamphetamine | 4 | 0 (0) | 40 | 8 (20) | 44 | 8 (18) |
| Pharmaceutical opioids | 1 | 0 (0) | 13 | 5 (38) | 14 | 5 (36) |
| OAT* | 0 | 0 (0) | 15 | 6 (40) | 16 | 6 (38) |
| Other drugs | 0 | 0 (0) | 13 | 6 (46) | 14 | 6 (43) |
| p value | | -- | | 0.206 | | 0.194 |
| 2022 | | | | | | |
| Methamphetamine | 0 | 0 (0) | 48 | 14 (29) | 49 | 15 (31) |
| Pharmaceutical opioids | 1 | 0 (0) | 20 | 5 (25) | 21 | 5 (24) |
| OAT* | 1 | 0 (0) | 11 | 6 (55) | 13 | 6 (46) |
| Other drugs | 0 | 0 (0) | 7 | 2 (29) | 7 | 2 (29) |
| p value | | -- | | 0.354 | | 0.589 |

*OAT = methadone, buprenorphine or buprenorphine/naloxone

Table 7.3.7 HCV antibody prevalence by frequency of drug injection last month, years since first injection and survey year

| Tasmania | <3 years | | 3+ years | | Total | |
|-----------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Frequency of injection last month | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Less than daily | 0 | 0 (0) | 33 | 19 (58) | 34 | 19 (56) |
| Daily or more | 1 | 1 (100) | 24 | 12 (50) | 26 | 14 (54) |
| Not last month | 0 | 0 (0) | 2 | 1 (50) | 2 | 1 (50) |
| p value | | -- | | 0.798 | | 1.000 |
| 2019 | | | | | | |
| Less than daily | 0 | 0 (0) | 24 | 7 (29) | 24 | 7 (29) |
| Daily or more | 0 | 0 (0) | 9 | 6 (67) | 9 | 6 (67) |
| Not last month | 0 | 0 (0) | 3 | 0 (0) | 3 | 0 (0) |
| p value | | -- | | 0.081 | | 0.081 |
| 2020 | | | | | | |
| Less than daily | 1 | 0 (0) | 18 | 6 (33) | 19 | 6 (32) |
| Daily or more | 0 | 0 (0) | 10 | 3 (30) | 10 | 3 (30) |
| Not last month | 0 | 0 (0) | 5 | 0 (0) | 5 | 0 (0) |
| p value | | -- | | 0.480 | | 0.485 |
| 2021 | | | | | | |
| Less than daily | 3 | 0 (0) | 47 | 11 (23) | 51 | 11 (22) |
| Daily or more | 2 | 0 (0) | 26 | 13 (50) | 29 | 13 (45) |
| Not last month | 0 | 0 (0) | 9 | 1 (11) | 9 | 1 (11) |
| p value | | -- | | 0.032 | | 0.048 |
| 2022 | | | | | | |
| Less than daily | 0 | 0 (0) | 39 | 11 (28) | 40 | 11 (28) |
| Daily or more | 2 | 0 (0) | 45 | 15 (33) | 48 | 16 (33) |
| Not last month | 0 | 0 (0) | 2 | 1 (50) | 2 | 1 (50) |
| p value | | -- | | 0.676 | | 0.608 |

Table 7.3.8 HCV antibody prevalence by imprisonment last year, years since first injection and survey year

| Tasmania | <3 years | | 3+ years | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Imprisonment last year | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| No imprisonment | 0 | 0 (0) | 55 | 29 (53) | 56 | 29 (52) |
| Imprisonment | 0 | 0 (0) | 3 | 3 (100) | 3 | 3 (100) |
| p value | | -- | | 0.245 | | 0.243 |
| 2019 | | | | | | |
| No imprisonment | 0 | 0 (0) | 33 | 10 (30) | 33 | 10 (30) |
| Imprisonment | 0 | 0 (0) | 4 | 3 (75) | 4 | 3 (75) |
| p value | | -- | | 0.115 | | 0.115 |
| 2020 | | | | | | |
| No imprisonment | 1 | 0 (0) | 28 | 8 (29) | 29 | 8 (28) |
| Imprisonment | 0 | 0 (0) | 5 | 1 (20) | 5 | 1 (20) |
| p value | | -- | | 0.582 | | 1.000 |
| 2021 | | | | | | |
| No imprisonment | 3 | 0 (0) | 73 | 23 (32) | 79 | 24 (30) |
| Imprisonment | 1 | 0 (0) | 6 | 3 (50) | 7 | 3 (43) |
| p value | | -- | | 0.389 | | 0.673 |
| 2022 | | | | | | |
| No imprisonment | 2 | 0 (0) | 70 | 19 (27) | 74 | 20 (27) |
| Imprisonment | 0 | 0 (0) | 14 | 7 (50) | 14 | 7 (50) |
| p value | | -- | | 0.091 | | 0.087 |

Table 7.3.9 HCV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Tasmania Aboriginal and Torres Strait Islander origin | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Non Indigenous | 32 | 14 (44) | 17 | 10 (59) | 49 | 24 (49) |
| Indigenous | 7 | 6 (86) | 3 | 1 (33) | 11 | 8 (73) |
| p value | | 0.091 | | 0.566 | | 0.193 |
| 2019 | | | | | | |
| Non Indigenous | 16 | 5 (31) | 17 | 6 (35) | 33 | 11 (33) |
| Indigenous | 3 | 2 (67) | 1 | 0 (0) | 4 | 2 (50) |
| p value | | 0.523 | | 1.000 | | 0.602 |
| 2020 | | | | | | |
| Non Indigenous | 16 | 5 (31) | 14 | 3 (21) | 30 | 8 (27) |
| Indigenous | 4 | 1 (25) | 0 | 0 (0) | 4 | 1 (25) |
| p value | | 1.000 | | -- | | 1.000 |
| 2021 | | | | | | |
| Non Indigenous | 39 | 13 (33) | 31 | 7 (23) | 70 | 20 (29) |
| Indigenous | 10 | 5 (50) | 8 | 2 (25) | 18 | 7 (39) |
| p value | | 0.329 | | 1.000 | | 0.397 |
| 2022 | | | | | | |
| Non Indigenous | 41 | 12 (29) | 20 | 3 (15) | 62 | 15 (24) |
| Indigenous | 17 | 9 (53) | 8 | 2 (25) | 25 | 11 (44) |
| p value | | 0.088 | | 0.606 | | 0.068 |

Table 7.3.10 HCV antibody prevalence by main language spoken at home by parents, gender and survey year

| Tasmania Main language spoken at home by parents | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| English speaking | 40 | 21 (53) | 21 | 12 (57) | 61 | 33 (54) |
| Non-English speaking | 0 | 0 (0) | 0 | 0 (0) | 1 | 1 (100) |
| p value | | -- | | -- | | 1.000 |
| 2019 | | | | | | |
| English speaking | 19 | 7 (37) | 18 | 6 (33) | 37 | 13 (35) |
| Non-English speaking | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2020 | | | | | | |
| English speaking | 20 | 6 (30) | 14 | 3 (21) | 34 | 9 (26) |
| Non-English speaking | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| English speaking | 45 | 16 (36) | 41 | 9 (22) | 86 | 25 (29) |
| Non-English speaking | 4 | 2 (50) | 0 | 0 (0) | 4 | 2 (50) |
| p value | | 0.618 | | -- | | 0.580 |
| 2022 | | | | | | |
| English speaking | 58 | 20 (34) | 29 | 6 (21) | 89 | 27 (30) |
| Non-English speaking | 1 | 1 (100) | 0 | 0 (0) | 1 | 1 (100) |
| p value | | 0.356 | | -- | | 0.311 |

Table 7.3.11 HCV antibody prevalence by region/country of birth, gender and survey year

| Tasmania | | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | |
| 2018 | | | | | | | |
| Australia | 38 | 21 (55) | 20 | 12 (60) | 58 | 33 (57) | |
| Other Oceania | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) | |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| UK & Ireland | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) | |
| Other | 0 | 0 (0) | 0 | 0 (0) | 1 | 1 (100) | |
| p value | | 0.219 | | 0.429 | | 0.137 | |
| 2019 | | | | | | | |
| Australia | 19 | 7 (37) | 17 | 6 (35) | 36 | 13 (36) | |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| UK & Ireland | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) | |
| Other | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| p value | | -- | | 1.000 | | 1.000 | |
| 2020 | | | | | | | |
| Australia | 19 | 6 (32) | 13 | 3 (23) | 32 | 9 (28) | |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| UK & Ireland | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) | |
| Other | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| p value | | 1.000 | | 1.000 | | 1.000 | |
| 2021 | | | | | | | |
| Australia | 42 | 15 (36) | 40 | 8 (20) | 82 | 23 (28) | |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| Asia | 0 | 0 (0) | 1 | 1 (100) | 1 | 1 (100) | |
| UK & Ireland | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| Other | 5 | 2 (40) | 1 | 0 (0) | 6 | 2 (33) | |
| p value | | 1.000 | | 0.387 | | 0.415 | |
| 2022 | | | | | | | |
| Australia | 56 | 19 (34) | 28 | 6 (21) | 86 | 26 (30) | |
| Other Oceania | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) | |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) | |
| UK & Ireland | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) | |
| Other | 1 | 1 (100) | 0 | 0 (0) | 1 | 1 (100) | |
| p value | | 0.575 | | 1.000 | | 0.667 | |

HCV RNA prevalence

Table 7.4.1 HCV RNA prevalence by gender and survey year

| Tasmania | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Survey year | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | 32 | 6 (19) | 17 | 5 (29) | 49 | 11 (22) |
| 2019* | 16 | 3 (19) | 16 | 3 (19) | 32 | 5 (16) |
| 2020 | 19 | 2 (11) | 12 | 2 (17) | 31 | 4 (13) |
| 2021 | 42 | 6 (14) | 35 | 3 (9) | 77 | 9 (12) |
| 2022 | 55 | 5 (9) | 29 | 0 (0) | 86 | 6 (7) |
| X ² p trend | | 0.189 | | 0.002 | | 0.009 |

* 2018, 2019: Weighted for gender and HCV antibody status

Totals include respondents where gender was reported as other or not reported

Table 7.4.2 HCV RNA prevalence by sexual identity, gender and survey year

| Tasmania | Male | | Female | | Total | |
|-----------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Sexual identity | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heterosexual | 28 | 4 (14) | 13 | 4 (31) | 41 | 8 (20) |
| Bisexual | 0 | 0 (0) | 2 | 1 (50) | 3 | 1 (33) |
| Homosexual | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | 0.544 | | 0.445 |
| 2019* | | | | | | |
| Heterosexual | 15 | 3 (20) | 12 | 2 (17) | 27 | 4 (15) |
| Bisexual | 1 | 0 (0) | 3 | 1 (33) | 4 | 1 (25) |
| Homosexual | 0 | 3 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | 0.637 | | 0.457 | | 0.789 |
| 2020 | | | | | | |
| Heterosexual | 18 | 2 (11) | 10 | 2 (20) | 28 | 4 (14) |
| Bisexual | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| Homosexual | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) |
| p value | | 0.725 | | 0.787 | | 0.782 |
| 2021 | | | | | | |
| Heterosexual | 38 | 5 (13) | 27 | 2 (7) | 65 | 7 (11) |
| Bisexual | 2 | 0 (0) | 6 | 0 (0) | 8 | 0 (0) |
| Homosexual | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| p value | | 0.583 | | 0.759 | | 0.586 |
| 2022 | | | | | | |
| Heterosexual | 52 | 4 (8) | 19 | 0 (0) | 72 | 5 (7) |
| Bisexual | 2 | 1 (50) | 7 | 0 (0) | 9 | 1 (11) |
| Homosexual | 1 | 0 (0) | 2 | 0 (0) | 3 | 0 (0) |
| p value | | 0.253 | | -- | | 0.668 |

Table 7.4.3 HCV RNA prevalence by age group, gender and survey year

| Tasmania Age group | Male | | Female | | Total | |
|-----------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 25-34 years | 5 | 1 (20) | 2 | 1 (50) | 7 | 2 (29) |
| 35-44 years | 15 | 1 (7) | 10 | 2 (20) | 25 | 3 (12) |
| 45+ years | 12 | 4 (33) | 4 | 1 (25) | 16 | 5 (31) |
| p value | | 0.227 | | 0.634 | | 0.430 |
| 2019* | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 0 | 0 (0) | 3 | 1 (33) | 3 | 1 (33) |
| 35-44 years | 10 | 1 (10) | 6 | 1 (17) | 16 | 2 (13) |
| 45+ years | 6 | 2 (33) | 6 | 1 (17) | 12 | 3 (25) |
| p value | | 0.340 | | 0.886 | | 0.669 |
| 2020 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 4 | 0 (0) | 5 | 1 (20) | 9 | 1 (11) |
| 35-44 years | 5 | 1 (20) | 2 | 0 (0) | 7 | 1 (14) |
| 45+ years | 10 | 1 (10) | 5 | 1 (20) | 15 | 2 (13) |
| p value | | 0.622 | | 0.787 | | 0.980 |
| 2021 | | | | | | |
| <25 years | 0 | 0 (0) | 2 | 0 (0) | 2 | 0 (0) |
| 25-34 years | 6 | 0 (0) | 6 | 0 (0) | 12 | 0 (0) |
| 35-44 years | 13 | 0 (0) | 9 | 0 (0) | 22 | 0 (0) |
| 45+ years | 23 | 6 (26) | 16 | 2 (13) | 39 | 8 (21) |
| p value | | 0.056 | | 0.520 | | 0.041 |
| 2022 | | | | | | |
| <25 years | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| 25-34 years | 10 | 1 (10) | 6 | 0 (0) | 16 | 1 (6) |
| 35-44 years | 16 | 0 (0) | 10 | 0 (0) | 27 | 1 (4) |
| 45+ years | 29 | 4 (14) | 12 | 0 (0) | 42 | 4 (10) |
| p value | | 0.373 | | -- | | 0.863 |

Table 7.4.4 HCV RNA prevalence by years since first injection, gender and survey year

| Tasmania Years of injection | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <3 years | 1 | 1 (100) | 0 | 0 (0) | 1 | 1 (100) |
| 3 to 10 years | 4 | 1 (25) | 3 | 1 (33) | 8 | 2 (25) |
| 11+ years | 26 | 3 (12) | 13 | 4 (31) | 39 | 7 (18) |
| p value | | 0.064 | | 0.710 | | 0.127 |
| 2019* | | | | | | |
| <3 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 3 to 10 years | 0 | 0 (0) | 3 | 2 (67) | 3 | 2 (67) |
| 11+ years | 16 | 3 (19) | 12 | 1 (8) | 29 | 3 (10) |
| p value | | -- | | 0.065 | | 0.066 |
| 2020 | | | | | | |
| <3 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 3 to 10 years | 2 | 0 (0) | 4 | 1 (25) | 6 | 1 (17) |
| 11+ years | 16 | 2 (13) | 8 | 1 (13) | 24 | 3 (13) |
| p value | | 0.811 | | 0.584 | | 0.893 |
| 2021 | | | | | | |
| <3 years | 1 | 0 (0) | 2 | 0 (0) | 3 | 0 (0) |
| 3 to 10 years | 2 | 0 (0) | 3 | 0 (0) | 5 | 0 (0) |
| 11+ years | 39 | 6 (15) | 27 | 2 (7) | 66 | 8 (12) |
| p value | | 0.764 | | 0.821 | | 0.581 |
| 2022 | | | | | | |
| <3 years | 0 | 0 (0) | 2 | 0 (0) | 2 | 0 (0) |
| 3 to 10 years | 3 | 0 (0) | 7 | 0 (0) | 10 | 0 (0) |
| 11+ years | 51 | 5 (10) | 19 | 0 (0) | 72 | 6 (8) |
| p value | | 1.000 | | -- | | 1.000 |

Table 7.4.5 HCV RNA prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| Tasmania Receptively shared syringe last month | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| No receptive sharing | 28 | 5 (18) | 16 | 5 (31) | 44 | 10 (23) |
| Receptive sharing | 2 | 0 (0) | 0 | 0 (0) | 3 | 0 (0) |
| p value | | 0.525 | | -- | | 0.375 |
| 2019* | | | | | | |
| No receptive sharing | 16 | 3 (19) | 13 | 3 (23) | 29 | 5 (17) |
| Receptive sharing | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2020 | | | | | | |
| No receptive sharing | 18 | 2 (11) | 8 | 2 (25) | 26 | 4 (15) |
| Receptive sharing | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No receptive sharing | 37 | 5 (14) | 25 | 1 (4) | 62 | 6 (10) |
| Receptive sharing | 2 | 0 (0) | 3 | 1 (33) | 5 | 1 (20) |
| p value | | 0.578 | | 0.062 | | 0.468 |
| 2022 | | | | | | |
| No receptive sharing | 45 | 5 (11) | 26 | 0 (0) | 72 | 5 (7) |
| Receptive sharing | 8 | 0 (0) | 3 | 0 (0) | 12 | 1 (8) |
| p value | | 1.000 | | -- | | 1.000 |

Table 7.4.6 HCV RNA prevalence by last drug injected, gender and survey year

| Tasmania Last drug injected | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Methamphetamine | 7 | 0 (0) | 7 | 2 (29) | 14 | 2 (14) |
| Pharm. Opioids | 14 | 3 (21) | 3 | 1 (33) | 17 | 4 (24) |
| OAT | 9 | 3 (33) | 5 | 1 (20) | 14 | 4 (29) |
| Other drugs | 2 | 0 (0) | 1 | 0 (0) | 4 | 0 (0) |
| p value | | 0.344 | | 0.902 | | 0.650 |
| 2019* | | | | | | |
| Methamphetamine | 8 | 2 (25) | 6 | 2 (33) | 14 | 3 (21) |
| Pharm. Opioids | 1 | 0 (0) | 2 | 0 (0) | 3 | 0 (0) |
| OAT | 6 | 0 (0) | 8 | 1 (13) | 14 | 1 (7) |
| Other drugs | 1 | 1 (100) | 0 | 0 (0) | 1 | 1 (100) |
| p value | | 0.161 | | 0.534 | | 0.067 |
| 2020 | | | | | | |
| Methamphetamine | 8 | 1 (13) | 6 | 1 (17) | 14 | 2 (14) |
| Pharm. Opioids | 6 | 1 (17) | 1 | 0 (0) | 7 | 1 (14) |
| OAT | 3 | 0 (0) | 3 | 1 (33) | 6 | 1 (17) |
| Other drugs | 2 | 0 (0) | 2 | 0 (0) | 4 | 0 (0) |
| p value | | 0.835 | | 0.753 | | 0.872 |
| 2021 | | | | | | |
| Methamphetamine | 12 | 0 (0) | 21 | 2 (10) | 33 | 2 (6) |
| Pharm. Opioids | 11 | 3 (27) | 3 | 0 (0) | 14 | 3 (21) |
| OAT | 11 | 1 (9) | 3 | 0 (0) | 14 | 1 (7) |
| Other drugs | 7 | 2 (29) | 6 | 0 (0) | 13 | 2 (15) |
| p value | | 0.185 | | 0.749 | | 0.409 |
| 2022 | | | | | | |
| Methamphetamine | 28 | 4 (14) | 17 | 0 (0) | 46 | 4 (9) |
| Pharm. Opioids | 16 | 1 (6) | 3 | 0 (0) | 20 | 2 (10) |
| OAT | 6 | 0 (0) | 7 | 0 (0) | 13 | 0 (0) |
| Other drugs | 5 | 0 (0) | 2 | 0 (0) | 7 | 0 (0) |
| p value | | 0.538 | | -- | | 0.574 |

OAT: methadone, buprenorphine or buprenorphine/naloxone

Table 7.4.7 HCV RNA prevalence by frequency of drug injection last month, gender and survey year

| Tasmania Frequency of injection last month | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| Less than daily | 16 | 1 (6) | 12 | 4 (33) | 28 | 5 (18) |
| Daily or more | 15 | 4 (27) | 4 | 1 (25) | 19 | 5 (26) |
| Not last month | 1 | 1 (100) | 1 | 0 (0) | 2 | 1 (50) |
| p value | | 0.045 | | 0.837 | | 0.433 |
| 2019* | | | | | | |
| Less than daily | 12 | 2 (17) | 9 | 2 (22) | 21 | 3 (14) |
| Daily or more | 4 | 1 (25) | 4 | 1 (25) | 8 | 2 (25) |
| Not last month | 0 | 0 (0) | 3 | 0 (0) | 3 | 0 (0) |
| p value | | 0.735 | | 0.705 | | 0.714 |
| 2020 | | | | | | |
| Less than daily | 12 | 1 (8) | 6 | 2 (33) | 18 | 3 (17) |
| Daily or more | 6 | 1 (17) | 2 | 0 (0) | 8 | 1 (13) |
| Not last month | 1 | 0 (0) | 4 | 0 (0) | 5 | 0 (0) |
| p value | | 0.811 | | 0.301 | | 0.616 |
| 2021 | | | | | | |
| Less than daily | 20 | 1 (5) | 21 | 2 (10) | 41 | 3 (7) |
| Daily or more | 19 | 4 (21) | 7 | 0 (0) | 26 | 4 (15) |
| Not last month | 3 | 1 (33) | 5 | 0 (0) | 8 | 1 (13) |
| p value | | 0.222 | | 0.544 | | 0.572 |
| 2022 | | | | | | |
| Less than daily | 25 | 2 (8) | 14 | 0 (0) | 40 | 2 (5) |
| Daily or more | 28 | 3 (11) | 15 | 0 (0) | 44 | 4 (9) |
| Not last month | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| p value | | 1.000 | | -- | | 0.722 |

Table 7.4.8 HCV RNA prevalence by imprisonment last year, gender and survey year

| Tasmania Imprisonment last year | Male | | Female | | Total | |
|------------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| No imprisonment | 28 | 4 (14) | 17 | 5 (29) | 45 | 9 (20) |
| Imprisonment | 1 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| p value | | 0.691 | | -- | | 0.500 |
| 2019* | | | | | | |
| No imprisonment | 14 | 1 (7) | 15 | 3 (20) | 29 | 3 (10) |
| Imprisonment | 3 | 2 (67) | 1 | 0 (0) | 3 | 2 (67) |
| p value | | 0.026 | | 0.654 | | 0.066 |
| 2020 | | | | | | |
| No imprisonment | 15 | 1 (7) | 11 | 2 (18) | 26 | 3 (12) |
| Imprisonment | 4 | 1 (25) | 1 | 0 (0) | 5 | 1 (20) |
| p value | | 0.288 | | 0.640 | | 0.605 |
| 2021 | | | | | | |
| No imprisonment | 36 | 6 (17) | 32 | 3 (9) | 68 | 9 (13) |
| Imprisonment | 4 | 0 (0) | 2 | 0 (0) | 6 | 0 (0) |
| p value | | 0.376 | | 0.650 | | 0.342 |
| 2022 | | | | | | |
| No imprisonment | 44 | 4 (9) | 24 | 0 (0) | 70 | 5 (7) |
| Imprisonment | 10 | 1 (10) | 4 | 0 (0) | 14 | 1 (7) |
| p value | | 1.000 | | -- | | 1.000 |

Table 7.4.9 HCV RNA prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Tasmania | Male | | Female | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| Aboriginal and Torres Strait Islander origin | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| Non Indigenous | 25 | 4 (16) | 12 | 5 (42) | 37 | 9 (24) |
| Indigenous | 6 | 1 (17) | 3 | 0 (0) | 10 | 1 (10) |
| p value | | 0.962 | | 0.221 | | 0.376 |
| 2019* | | | | | | |
| Non Indigenous | 15 | 2 (13) | 15 | 3 (20) | 29 | 4 (14) |
| Indigenous | 2 | 1 (50) | 1 | 0 (0) | 3 | 1 (33) |
| p value | | 0.203 | | 0.654 | | 0.417 |
| 2020 | | | | | | |
| Non Indigenous | 15 | 2 (13) | 12 | 2 (17) | 27 | 4 (15) |
| Indigenous | 4 | 0 (0) | 0 | 0 (0) | 4 | 0 (0) |
| p value | | 0.440 | | -- | | 0.409 |
| 2021 | | | | | | |
| Non Indigenous | 35 | 6 (17) | 24 | 3 (13) | 59 | 9 (15) |
| Indigenous | 7 | 0 (0) | 8 | 0 (0) | 15 | 0 (0) |
| p value | | 0.237 | | 0.294 | | 0.107 |
| 2022 | | | | | | |
| Non Indigenous | 38 | 3 (8) | 20 | 0 (0) | 59 | 3 (5) |
| Indigenous | 16 | 2 (13) | 8 | 0 (0) | 24 | 2 (8) |
| p value | | 0.627 | | -- | | 0.624 |

Table 7.4.10 HCV RNA prevalence by main language spoken at home by parents, gender and survey year

| Tasmania | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| Main language spoken at home by parents | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| English speaking | 31 | 6 (19) | 17 | 5 (29) | 48 | 11 (23) |
| Non-English speaking | 0 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | -- | | 0.605 |
| 2019* | | | | | | |
| English speaking | 16 | 3 (19) | 16 | 3 (19) | 32 | 5 (16) |
| Non-English speaking | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2020 | | | | | | |
| English speaking | 19 | 2 (11) | 12 | 2 (17) | 31 | 4 (13) |
| Non-English speaking | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| English speaking | 39 | 6 (15) | 35 | 3 (9) | 74 | 9 (12) |
| Non-English speaking | 3 | 0 (0) | 0 | 0 (0) | 3 | 0 (0) |
| p value | | 0.463 | | -- | | 0.520 |
| 2022 | | | | | | |
| English speaking | 54 | 5 (9) | 29 | 0 (0) | 85 | 6 (7) |
| Non-English speaking | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | 1.000 | | -- | | 1.000 |

Table 7.4.11 HCV RNA prevalence by region/country of birth, gender and survey year

| Tasmania | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Australia | 29 | 6 (21) | 16 | 5 (31) | 45 | 11 (24) |
| Other Oceania | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) |
| Other | 0 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | 0.786 | | 0.540 | | 0.756 |
| 2019* | | | | | | |
| Australia | 14 | 3 (21) | 15 | 3 (20) | 31 | 5 (16) |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| Other | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | -- | | 0.654 | | 0.664 |
| 2020 | | | | | | |
| Australia | 18 | 2 (11) | 11 | 2 (18) | 29 | 4 (14) |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 1 | 0 (0) | 1 | 0 (0) | 2 | 0 (0) |
| Other | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| p value | | 0.725 | | 0.640 | | 0.574 |
| 2021 | | | | | | |
| Australia | 35 | 6 (17) | 34 | 3 (9) | 69 | 9 (13) |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| Asia | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| UK & Ireland | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| Other | 5 | 0 (0) | 0 | 0 (0) | 5 | 0 (0) |
| p value | | 0.315 | | 0.756 | | 0.641 |
| 2022 | | | | | | |
| Australia | 52 | 5 (10) | 28 | 0 (0) | 82 | 6 (7) |
| Other Oceania | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| Other | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | 1.000 | | -- | | 1.000 |

Victoria

Table 8.1.1 Number (percentage) of respondents by demographic characteristics and survey year

| Victoria | 2018 | 2019 | 2020* | 2021* | 2022 |
|---|----------|----------|-------|----------|----------|
| Number of sites | 6 | 6 | -- | 3 | 5 |
| N° surveyed | N=437 | N=375 | -- | N=40 | N=122 |
| Response rate (%) | 51% | 52% | -- | 44% | 43% |
| Gender (%) | | | | | |
| Male | 290 (66) | 240 (64) | -- -- | 28 (70) | 78 (64) |
| Female | 145 (33) | 130 (35) | -- -- | 11 (28) | 43 (35) |
| Other | 2 (<1) | 3 (1) | -- -- | 1 (3) | 1 (1) |
| Not reported | 0 (0) | 2 (1) | -- -- | 0 (0) | 0 (0) |
| Sexual identity (%) | | | | | |
| Heterosexual | 342 (78) | 277 (74) | -- -- | 30 (75) | 92 (75) |
| Bisexual | 33 (8) | 39 (10) | -- -- | 7 (18) | 14 (11) |
| Gay/Lesbian | 18 (4) | 17 (5) | -- -- | 1 (3) | 9 (7) |
| Other | -- -- | -- -- | -- -- | 2 (5) | 7 (6) |
| Not reported | 44 (10) | 42 (11) | -- -- | 0 (0) | 0 (0) |
| Aboriginal and Torres Strait Islander origin (%) | | | | | |
| No | 359 (82) | 296 (79) | -- -- | 31 (78) | 101 (83) |
| Yes | 71 (16) | 66 (18) | -- -- | 9 (23) | 19 (16) |
| Not reported | 7 (2) | 13 (3) | -- -- | 0 (0) | 2 (2) |
| Region/country of birth (%) | | | | | |
| Australia | 373 (85) | 333 (89) | -- -- | 33 (83) | 96 (79) |
| Other Oceania | 12 (3) | 7 (2) | -- -- | 2 (5) | 3 (2) |
| Asia | 6 (1) | 3 (1) | -- -- | 0 (0) | 0 (0) |
| UK & Ireland | 20 (5) | 7 (2) | -- -- | 2 (5) | 6 (5) |
| Other | 24 (5) | 20 (5) | -- -- | 3 (8) | 12 (10) |
| Not reported | 2 (<1) | 5 (1) | -- -- | 0 (0) | 5 (4) |
| Main language spoken at home by parents (%) | | | | | |
| English | 393 (90) | 345 (92) | -- -- | 36 (90) | 103 (84) |
| Non-English | 43 (10) | 27 (7) | -- -- | 4 (10) | 16 (13) |
| Not reported | 1 (<1) | 3 (1) | -- -- | 0 (0) | 3 (2) |
| Age and time since first injection (years) | | | | | |
| <i>Median age</i> | 42 | 42 | -- -- | 46 | 46 |
| <i>Age range</i> | 17-69 | 20-70 | -- -- | 29-62 | 25-66 |
| <i>Age group (%)</i> | | | | | |
| <25 years | 8 (2) | 6 (2) | -- -- | 0 (0) | 0 (0) |
| 25+ years | 424 (97) | 363 (97) | -- -- | 40 (100) | 121 (99) |
| Not reported | 5 (1) | 6 (2) | -- -- | 0 (0) | 1 (1) |
| <i>Median age first injection</i> | 18 | 18 | -- -- | 18 | 18 |
| <i>Age range</i> | 11-52 | 10-51 | -- -- | 13-50 | 11-50 |
| <i>N° not reported</i> | 23 | 27 | -- -- | 0 | 5 |
| <i>Median yrs since first injection</i> | 22 | 22 | -- -- | 27.5 | 27 |
| <i>Range</i> | <1-46 | <1-52 | -- -- | <1-41 | <1-52 |
| <i>Years since first injection</i> | | | | | |
| <3 years | 18 (4) | 13 (3) | -- -- | 2 (5) | 2 (2) |
| 3+ years | 396 (91) | 333 (89) | -- -- | 38 (95) | 115 (94) |
| Not reported | 23 (5) | 29 (8) | -- -- | 0 (0) | 5 (4) |
| Imprisonment last year (%) | | | | | |
| No | 335 (77) | 268 (71) | -- -- | 34 (85) | 93 (76) |
| Yes | 75 (17) | 83 (22) | -- -- | 2 (5) | 15 (12) |
| Not reported | 27 (6) | 24 (6) | -- -- | 4 (10) | 14 (11) |
| N° in prison | N=75 | N=83 | -- -- | N=2 | N=15 |
| Injected in prison | 19 (25) | 20 (24) | -- -- | 0 (0) | 5 (33) |

* Implementation of the 2020 and 2021 ANSPS was impacted by the COVID-19 pandemic, see page 1 for more information

Table 8.1.2 Number (percentage) of respondents by last drug injected, frequency of injecting and survey year

| Victoria | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|----------|----------|-------|---------|----------|
| N° surveyed | N=437 | N=375 | -- | N=40 | N=122 |
| Drug last injected (%) | | | | | |
| Cocaine | 3 (1) | 4 (1) | -- -- | 0 (0) | 1 (1) |
| Methamphetamine | 158 (36) | 126 (34) | -- -- | 18 (45) | 35 (29) |
| Heroin | 223 (51) | 198 (53) | -- -- | 16 (40) | 70 (57) |
| Pharm. opioids | 4 (1) | 8 (2) | -- -- | 1 (3) | 0 (0) |
| Methadone | 3 (1) | 3 (1) | -- -- | 0 (0) | 1 (1) |
| Buprenorphine | 7 (2) | 1 (<1) | -- -- | 1 (3) | 3 (2) |
| Buprenorphine/naloxone | 8 (2) | 2 (1) | -- -- | 0 (0) | 0 (0) |
| PIEDs | 2 (<1) | 0 (0) | -- -- | 0 (0) | 0 (0) |
| More than one | 22 (5) | 25 (7) | -- -- | 3 (8) | 9 (7) |
| Other | 5 (1) | 5 (1) | -- -- | 1 (3) | 2 (2) |
| Not reported | 2 (<1) | 3 (1) | -- -- | 0 (0) | 1 (1) |
| Ever injected opioids | | | | | |
| No | 40 (9) | 31 (8) | -- -- | 6 (15) | 5 (4) |
| Yes | 393 (90) | 343 (91) | -- -- | 34 (85) | 117 (96) |
| Not reported | 4 (1) | 1 (<1) | -- -- | 0 (0) | 0 (0) |
| Frequency of injection last month (%) | | | | | |
| Not last month | 36 (8) | 23 (6) | -- -- | 2 (5) | 10 (8) |
| Less than weekly | 66 (15) | 47 (13) | -- -- | 10 (25) | 16 (13) |
| Weekly not daily | 104 (24) | 62 (17) | -- -- | 6 (15) | 20 (16) |
| Daily or more | 224 (51) | 240 (64) | -- -- | 22 (55) | 73 (60) |
| Not reported | 7 (2) | 3 (1) | -- -- | 0 (0) | 3 (2) |
| Present when someone injected for the first time in the last 12 months (%) | | | | | |
| No | 356 (81) | 299 (80) | -- -- | 28 (70) | 90 (74) |
| Yes | 67 (15) | 65 (17) | -- -- | 7 (18) | 25 (20) |
| Not reported | 14 (3) | 11 (3) | -- -- | 5 (13) | 7 (6) |
| Overdosed in last 12 months | | | | | |
| No | 297 (68) | 255 (68) | -- -- | 30 (75) | 85 (70) |
| Yes | 125 (29) | 103 (27) | -- -- | 4 (10) | 28 (23) |
| Not reported | 15 (3) | 17 (5) | -- -- | 6 (15) | 9 (7) |

Table 8.1.3 Number (percentage) of respondents by injecting behaviour in the last month and survey year

| Victoria | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|----------|----------|-------|---------|---------|
| N° injected last month | N=394 | N=349 | -- | N=38 | N=109 |
| Places injected last month (%) | | | | | |
| All private | 152 (39) | 125 (36) | -- -- | 18 (47) | 42 (39) |
| Any public | 239 (61) | 222 (64) | -- -- | 20 (53) | 65 (60) |
| Not reported | 3 (1) | 2 (1) | -- -- | 0 (0) | 2 (2) |
| Use of new and sterile needles and syringes last month (%) | | | | | |
| All injections | 301 (76) | 252 (72) | -- -- | 34 (89) | 82 (75) |
| Most of the time | 69 (18) | 78 (22) | -- -- | 2 (5) | 21 (19) |
| Half of the time | 6 (2) | 3 (1) | -- -- | 0 (0) | 3 (3) |
| Some of the time | 6 (2) | 3 (1) | -- -- | 0 (0) | 1 (1) |
| Not last month | 5 (1) | 2 (1) | -- -- | 2 (5) | 1 (1) |
| Not reported | 7 (2) | 11 (3) | -- -- | 0 (0) | 1 (1) |
| Re-used someone else's used needle & syringe last month (%) | | | | | |
| None | 321 (81) | 291 (83) | -- -- | 35 (92) | 84 (77) |
| Once | 22 (6) | 20 (6) | -- -- | 0 (0) | 7 (6) |
| Twice | 14 (4) | 13 (4) | -- -- | 2 (5) | 8 (7) |
| 3-5 times | 16 (4) | 15 (4) | -- -- | 0 (0) | 3 (3) |
| >5 times | 18 (5) | 6 (2) | -- -- | 1 (3) | 6 (6) |
| Not reported | 3 (1) | 4 (1) | -- -- | 0 (0) | 1 (1) |
| Number of people needle & syringe was re-used after last month (%) | | | | | |
| None | 321 (81) | 291 (83) | -- -- | 34 (89) | 84 (77) |
| One | 26 (7) | 27 (8) | -- -- | 1 (3) | 4 (4) |
| Two | 5 (1) | 4 (1) | -- -- | 0 (0) | 3 (3) |
| Three to five | 6 (2) | 3 (1) | -- -- | 0 (0) | 4 (4) |
| More than five | 6 (2) | 0 (0) | -- -- | 0 (0) | 2 (2) |
| Don't know | 17 (4) | 15 (4) | -- -- | 2 (5) | 4 (4) |
| Not reported | 13 (3) | 9 (3) | -- -- | 1 (3) | 8 (7) |
| Relationship to people needle & syringe was used after last month (%) [more than one could be selected] | | | | | |
| Regular sex partner | 26 (7) | 21 (6) | -- -- | 0 (0) | 4 (4) |
| Casual sex partner | 6 (2) | 3 (1) | -- -- | 0 (0) | 1 (1) |
| Close friend | 18 (5) | 10 (3) | -- -- | 1 (3) | 4 (4) |
| Acquaintance | 4 (1) | 11 (3) | -- -- | 0 (0) | 4 (4) |
| Other | 6 (2) | 11 (3) | -- -- | 0 (0) | 2 (2) |
| Equipment used after someone else last month (%) [more than one could be selected] | | | | | |
| Spoon | 132 (34) | 96 (28) | -- -- | 9 (24) | 30 (28) |
| Water | 77 (20) | 73 (21) | -- -- | 4 (11) | 22 (20) |
| Filter | 42 (11) | 51 (15) | -- -- | 5 (13) | 12 (11) |
| Drug mix | 46 (12) | 37 (11) | -- -- | 3 (8) | 9 (8) |
| None | 233 (59) | 222 (64) | -- -- | 28 (74) | 71 (65) |
| Injected by someone after that person injected themselves or others last month (%) | | | | | |
| No | 303 (77) | 291 (83) | -- -- | 35 (92) | 85 (78) |
| Yes | 84 (21) | 54 (15) | -- -- | 3 (8) | 19 (17) |
| Not reported | 7 (2) | 4 (1) | -- -- | 0 (0) | 5 (5) |
| Source of needle acquisition | | | | | |
| Needle Syringe Program | 358 (91) | 314 (90) | -- -- | 31 (82) | 92 (84) |
| Chemist/Pharmacy | 61 (15) | 63 (18) | -- -- | 4 (11) | 10 (9) |
| Personal sources | 50 (13) | 53 (15) | -- -- | 5 (13) | 10 (9) |
| Dispensing/Vending Machine | 90 (23) | 68 (19) | -- -- | 16 (42) | 28 (26) |
| Other sources | 21 (5) | 13 (4) | -- -- | 0 (0) | 5 (5) |

Table 8.1.4 Number (percentage) of respondents by treatment for drug use and survey year

| Victoria | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|----------|----------|-------|---------|----------|
| N° surveyed | N=437 | N=375 | -- | N=40 | N=122 |
| Ever any treatment/therapy for drug use (%) | | | | | |
| No | 51 (12) | 53 (14) | -- -- | 6 (15) | 13 (11) |
| Yes | 385 (88) | 320 (85) | -- -- | 34 (85) | 108 (89) |
| Not reported | 1 (<1) | 2 (1) | -- -- | 0 (0) | 1 (1) |
| History of methadone maintenance treatment (%) | | | | | |
| Currently | 139 (32) | 143 (38) | -- -- | 16 (40) | 37 (30) |
| Previously | 155 (35) | 114 (30) | -- -- | 12 (30) | 51 (42) |
| Never | 134 (31) | 109 (29) | -- -- | 11 (28) | 30 (25) |
| Not reported | 9 (2) | 9 (2) | -- -- | 1 (3) | 4 (3) |
| History of buprenorphine (Subutex®) treatment (%) | | | | | |
| Currently | 33 (8) | 25 (7) | -- -- | 4 (10) | 16 (13) |
| Previously | 170 (39) | 146 (39) | -- -- | 14 (35) | 53 (43) |
| Never | 229 (52) | 194 (52) | -- -- | 21 (53) | 49 (40) |
| Not reported | 5 (1) | 10 (3) | -- -- | 1 (3) | 4 (3) |
| History of buprenorphine-naloxone (Suboxone®) treatment (%) | | | | | |
| Currently | 57 (13) | 27 (7) | -- -- | 3 (8) | 8 (7) |
| Previously | 159 (36) | 149 (40) | -- -- | 14 (35) | 56 (46) |
| Never | 216 (49) | 193 (51) | -- -- | 23 (58) | 56 (46) |
| Not reported | 5 (1) | 6 (2) | -- -- | 0 (0) | 2 (2) |

Table 8.1.5 Number (percentage) of respondents by testing for HIV and HCV, HCV treatment by survey year

| Victoria | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|----------|----------|-------|---------|----------|
| N° surveyed | N=437 | N=375 | -- | N=40 | N=122 |
| Previous HIV test (%) | | | | | |
| Yes, ever | 364 (83) | 306 (82) | -- -- | 35 (88) | 100 (82) |
| Yes, last year | 226 (52) | 206 (55) | -- -- | 12 (30) | 58 (48) |
| >1 year ago | 138 (32) | 100 (27) | -- -- | 23 (58) | 42 (34) |
| Never tested | 57 (13) | 54 (14) | -- -- | 1 (3) | 13 (11) |
| Not reported | 16 (4) | 15 (4) | -- -- | 4 (10) | 9 (7) |
| Previous HCV test (%) | | | | | |
| Yes, ever | 363 (83) | 309 (82) | -- -- | 33 (83) | 104 (85) |
| Yes, last year | 243 (56) | 218 (58) | -- -- | 17 (43) | 66 (54) |
| >1 year ago | 120 (27) | 91 (24) | -- -- | 17 (43) | 38 (31) |
| Never tested | 30 (7) | 23 (6) | -- -- | 1 (3) | 8 (7) |
| Not reported | 44 (10) | 43 (11) | -- -- | 5 (13) | 10 (8) |
| Lifetime treatment for HCV (%)# | | | | | |
| N° self-reported HCV diagnosis | N=150 | N=131 | -- | N=11 | N=21 |
| Antiviral treatment | 86 (57) | 87 (66) | -- -- | 5 (45) | 15 (71) |
| No antiviral treatment | 63 (42) | 44 (34) | -- -- | 6 (55) | 6 (29) |
| Not reported | 1 (1) | 0 (0) | -- -- | 0 (0) | 0 (0) |
| Treatment for HCV in past 12 months (%)#* | | | | | |
| N° self-reported HCV diagnosis | N=121 | N=100 | -- | N=10 | N=14 |
| Antiviral treatment | 53 (44) | 48 (48) | -- -- | 2 (20) | 5 (36) |
| No antiviral treatment | 67 (55) | 52 (52) | -- -- | 8 (80) | 9 (64) |
| Not reported | 1 (1) | 0 (0) | -- -- | 0 (0) | 0 (0) |

among people who tested HCV antibody positive and did not report spontaneous clearance

* excludes people who reported treatment induced clearance more than 12 months previously

Table 8.1.6 Number (percentage) of respondents by sexual behaviour in the month prior to survey and survey year

| Victoria | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|----------|----------|-------|---------|----------|
| N° surveyed | N=437 | N=375 | -- | N=40 | N=122 |
| Sex with a regular partner last month (%) | | | | | |
| No | 253 (58) | 212 (57) | -- -- | 31 (78) | 74 (61) |
| Yes | 173 (40) | 147 (39) | -- -- | 6 (15) | 39 (32) |
| Not reported | 11 (3) | 16 (4) | -- -- | 3 (8) | 9 (7) |
| Condom used with regular partner last month (%) | | | | | |
| N° surveyed | N=173 | N=147 | -- | N=6 | N=39 |
| Never | 126 (73) | 102 (69) | -- -- | 5 (83) | 29 (74) |
| Sometimes | 23 (13) | 20 (14) | -- -- | 0 (0) | 4 (10) |
| Every time | 10 (6) | 13 (9) | -- -- | 1 (17) | 4 (10) |
| Not reported | 14 (8) | 12 (8) | -- -- | 0 (0) | 2 (5) |
| Sex with other partner(s) last month (%) | | | | | |
| No | 355 (81) | 297 (79) | -- -- | 25 (63) | 94 (77) |
| Yes | 58 (13) | 54 (14) | -- -- | 8 (20) | 19 (16) |
| Not reported | 24 (5) | 24 (6) | -- -- | 7 (18) | 9 (7) |
| Condom used with other partner(s) last month (%) | | | | | |
| N° surveyed | N=58 | N=54 | -- | N=8 | N=19 |
| Never | 22 (38) | 25 (46) | -- -- | 2 (25) | 9 (47) |
| Sometimes | 18 (31) | 14 (26) | -- -- | 1 (13) | 6 (32) |
| Every time | 15 (26) | 11 (20) | -- -- | 4 (50) | 4 (21) |
| Not reported | 3 (5) | 4 (7) | -- -- | 1 (13) | 0 (0) |
| Sex work last month (%) | | | | | |
| No | 378 (87) | 325 (87) | -- -- | 30 (75) | 102 (84) |
| Yes | 29 (7) | 30 (8) | -- -- | 5 (13) | 8 (7) |
| Not reported | 30 (7) | 20 (5) | -- -- | 5 (13) | 12 (10) |
| Condom used at last sex work (%) | | | | | |
| Yes | 22 (76) | 17 (57) | -- -- | 4 (80) | 6 (75) |

Table 8.1.7 Number (percentage) of respondents by COVID-19 related impacts, 2020-2022

| Victoria | 2020 | 2021 | 2022 |
|---|-------|---------|----------|
| N° surveyed | -- | N=40 | N=122 |
| COVID-19 test | | | |
| Yes | -- -- | 27 (68) | 109 (89) |
| No | -- -- | 9 (23) | 10 (8) |
| Don't know | -- -- | 0 (0) | -- -- |
| Not reported | -- -- | 4 (10) | 3 (2) |
| Self-reported positive COVID-19 test result | | | |
| Yes | -- -- | 2 (5) | 38 (35) |
| Yes, last 12 months | -- -- | -- -- | 28 (26) |
| Yes, more than a year ago | -- -- | -- -- | 10 (9) |
| COVID-19 vaccine uptake | | | |
| ≥1 dose | -- -- | 31 (78) | 102 (84) |
| COVID-19 vaccine doses | | | |
| One dose | -- -- | 9 (23) | 16 (13) |
| Two doses | -- -- | 22 (55) | 27 (22) |
| Three or more doses | -- -- | -- -- | 59 (48) |
| No doses | -- -- | 7 (18) | 15 (12) |
| Not reported | -- -- | 2 (5) | 5 (4) |
| Vaccine type * | | | N=102 |
| Astra Zeneca | -- -- | 6 (15) | 25 (25) |
| Pfizer | -- -- | 19 (48) | 73 (72) |
| Don't know | -- -- | 4 (10) | 13 (13) |
| Other | -- -- | -- -- | 5 (5) |
| Not reported | -- -- | 11 (28) | -- -- |
| * 2021 only one response permitted, 2022 multiple responses permitted | | | |
| Last vaccine location | | | N=122 |
| GP or medical centre | -- -- | -- -- | 43 (35) |
| Pharmacy/ chemist | -- -- | -- -- | 16 (13) |
| Vaccine hub | -- -- | -- -- | 21 (17) |
| Opioid treatment/ methadone clinic | -- -- | -- -- | 3 (2) |
| NSP | -- -- | -- -- | 9 (7) |
| Prison | -- -- | -- -- | 4 (3) |
| Other | -- -- | -- -- | 3 (2) |
| Multiple response | -- -- | -- -- | 9 (7) |
| Not reported | -- -- | -- -- | 14 (11) |
| Vaccine hesitancy - if not had three or more doses | | | |
| Haven't got around to it | -- -- | -- -- | 20 (16) |
| Difficulty booking appointment | -- -- | -- -- | 4 (3) |
| Concerns about side effects | -- -- | -- -- | 5 (4) |
| Concerns about vaccine safety | -- -- | -- -- | 5 (4) |
| Not at risk of serious illness | -- -- | -- -- | 6 (5) |
| Other | -- -- | -- -- | 11 (9) |
| Multiple response | -- -- | -- -- | 10 (8) |
| Not reported | -- -- | -- -- | 61 (50) |

HIV antibody prevalence

Table 8.2.1 HIV antibody prevalence by gender and survey year

| Victoria Survey year | Male | | Female | | Total | |
|-------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | 289 | 8 (2.8) | 145 | 1 (0.7) | 436 | 9 (2.1) |
| 2019 | 234 | 8 (3.4) | 128 | 1 (0.8) | 367 | 9 (2.5) |
| 2020 | -- | -- -- | -- | -- -- | -- | -- -- |
| 2021 | 25 | 1 (4.0) | 11 | 0 (0.0) | 37 | 2 (5.4) |
| 2022 | 68 | 1 (1.5) | 41 | 1 (2.4) | 109 | 2 (1.8) |
| X ² p trend | | 0.710 | | 0.376 | | 0.761 |

Table 8.2.2 HIV antibody prevalence by sexual identity, gender and survey year

| Victoria Sexual identity | Male | | Female | | Total | |
|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) | N ^o tested | N ^o with HIV (%) |
| 2018 | | | | | | |
| Heterosexual | 240 | 5 (2.1) | 101 | 1 (1.0) | 341 | 6 (1.8) |
| Bisexual | 11 | 1 (9.1) | 22 | 0 (0.0) | 33 | 1 (3.0) |
| Homosexual | 11 | 2 (18.2) | 6 | 0 (0.0) | 18 | 2 (11.1) |
| p value | | 0.022 | | 1.000 | | 0.042 |
| 2019 | | | | | | |
| Heterosexual | 194 | 7 (3.6) | 78 | 1 (1.3) | 272 | 8 (2.9) |
| Bisexual | 11 | 0 (0.0) | 22 | 0 (0.0) | 37 | 0 (0.0) |
| Homosexual | 6 | 1 (16.7) | 10 | 0 (0.0) | 16 | 1 (6.3) |
| p value | | 0.258 | | 1.000 | | 0.299 |
| 2020 | | | | | | |
| Heterosexual | -- | -- -- | -- | -- -- | -- | -- -- |
| Bisexual | -- | -- -- | -- | -- -- | -- | -- -- |
| Homosexual | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Heterosexual | 22 | 0 (0.0) | 4 | 0 (0.0) | 27 | 1 (3.7) |
| Bisexual | 1 | 0 (0.0) | 6 | 0 (0.0) | 7 | 0 (0.0) |
| Homosexual | 1 | 1 (100.0) | 1 | 0 (0.0) | 1 | 1 (100.0) |
| p value | | 0.120 | | -- | | 0.108 |
| 2022 | | | | | | |
| Heterosexual | 59 | 0 (0.0) | 24 | 1 (4.2) | 83 | 1 (1.2) |
| Bisexual | 3 | 0 (0.0) | 10 | 0 (0.0) | 13 | 0 (0.0) |
| Homosexual | 4 | 1 (25.0) | 3 | 0 (0.0) | 7 | 1 (14.3) |
| p value | | 0.132 | | 1.000 | | 0.239 |

Table 8.2.3 HIV antibody prevalence by age group, gender and survey year

| Victoria Age group | Male | | Female | | Total | |
|-----------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| <25 years | 2 | 0 (0.0) | 6 | 0 (0.0) | 8 | 0 (0.0) |
| 25-34 years | 43 | 4 (9.3) | 30 | 0 (0.0) | 73 | 4 (5.5) |
| 35-44 years | 117 | 0 (0.0) | 57 | 0 (0.0) | 175 | 0 (0.0) |
| 45+ years | 123 | 4 (3.3) | 51 | 1 (2.0) | 175 | 5 (2.9) |
| p value | | 0.013 | | 0.604 | | 0.024 |
| 2019 | | | | | | |
| <25 years | 3 | 0 (0.0) | 3 | 0 (0.0) | 6 | 0 (0.0) |
| 25-34 years | 32 | 3 (9.4) | 25 | 0 (0.0) | 59 | 3 (5.1) |
| 35-44 years | 97 | 3 (3.1) | 54 | 0 (0.0) | 152 | 3 (2.0) |
| 45+ years | 99 | 1 (1.0) | 45 | 1 (2.2) | 145 | 2 (1.4) |
| p value | | 0.139 | | 0.575 | | 0.387 |
| 2020 | | | | | | |
| <25 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 25-34 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 35-44 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 45+ years | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| <25 years | 0 | 0 (0.0) | 0 | 0 (0.0) | 0 | 0 (0.0) |
| 25-34 years | 0 | 0 (0.0) | 1 | 0 (0.0) | 2 | 1 (50.0) |
| 35-44 years | 6 | 1 (16.7) | 4 | 0 (0.0) | 10 | 1 (10.0) |
| 45+ years | 19 | 0 (0.0) | 6 | 0 (0.0) | 25 | 0 (0.0) |
| p value | | 0.240 | | -- | | 0.032 |
| 2022 | | | | | | |
| <25 years | 0 | 0 (0.0) | 0 | 0 (0.0) | 0 | 0 (0.0) |
| 25-34 years | 5 | 0 (0.0) | 6 | 0 (0.0) | 11 | 0 (0.0) |
| 35-44 years | 17 | 0 (0.0) | 18 | 1 (5.6) | 35 | 1 (2.9) |
| 45+ years | 46 | 1 (2.2) | 16 | 0 (0.0) | 62 | 1 (1.6) |
| p value | | 1.000 | | 1.000 | | 1.000 |

Table 8.2.4 HIV antibody prevalence by last drug injected, gender and survey year

| Victoria Last drug injected | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Heroin | 145 | 5 (3.5) | 77 | 1 (1.3) | 222 | 6 (2.7) |
| Methamphetamine | 109 | 3 (2.8) | 48 | 0 (0.0) | 158 | 3 (1.9) |
| Other opioids | 12 | 0 (0.0) | 9 | 0 (0.0) | 22 | 0 (0.0) |
| Other drugs | 23 | 0 (0.0) | 9 | 0 (0.0) | 32 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 0.922 |
| 2019 | | | | | | |
| Heroin | 120 | 5 (4.2) | 72 | 1 (1.4) | 194 | 6 (3.1) |
| Methamphetamine | 78 | 2 (2.6) | 44 | 0 (0.0) | 124 | 2 (1.6) |
| Other opioids | 11 | 1 (9.1) | 2 | 0 (0.0) | 13 | 1 (7.7) |
| Other drugs | 22 | 0 (0.0) | 10 | 0 (0.0) | 33 | 0 (0.0) |
| p value | | 0.555 | | 1.000 | | 0.318 |
| 2020 | | | | | | |
| Heroin | -- | -- -- | -- | -- -- | -- | -- -- |
| Methamphetamine | -- | -- -- | -- | -- -- | -- | -- -- |
| Other opioids | -- | -- -- | -- | -- -- | -- | -- -- |
| Other drugs | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Heroin | 10 | 0 (0.0) | 4 | 0 (0.0) | 15 | 1 (6.7) |
| Methamphetamine | 10 | 1 (10.0) | 6 | 0 (0.0) | 16 | 1 (6.3) |
| Other opioids | 2 | 0 (0.0) | 0 | 0 (0.0) | 2 | 0 (0.0) |
| Other drugs | 3 | 0 (0.0) | 1 | 0 (0.0) | 4 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2022 | | | | | | |
| Heroin | 43 | 0 (0.0) | 23 | 1 (4.4) | 66 | 1 (1.5) |
| Methamphetamine | 16 | 1 (6.3) | 13 | 0 (0.0) | 29 | 1 (3.5) |
| Other opioids | 2 | 0 (0.0) | 1 | 0 (0.0) | 3 | 0 (0.0) |
| Other drugs | 7 | 0 (0.0) | 4 | 0 (0.0) | 11 | 0 (0.0) |
| p value | | 0.368 | | 1.000 | | 0.636 |

Table 8.2.5 HIV antibody prevalence by years of injection, gender and survey year

| Victoria Years of injection | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| <3 years | 12 | 0 (0.0) | 6 | 0 (0.0) | 18 | 0 (0.0) |
| 3 to 10 years | 24 | 2 (8.3) | 24 | 0 (0.0) | 48 | 2 (4.2) |
| 11+ years | 253 | 6 (2.4) | 115 | 1 (0.9) | 370 | 7 (1.9) |
| p value | | 0.171 | | 1.000 | | 0.507 |
| 2019 | | | | | | |
| <3 years | 6 | 0 (0.0) | 7 | 0 (0.0) | 13 | 0 (0.0) |
| 3 to 10 years | 23 | 2 (8.7) | 20 | 0 (0.0) | 45 | 2 (4.4) |
| 11+ years | 188 | 5 (2.7) | 93 | 1 (1.1) | 282 | 6 (2.1) |
| p value | | 0.320 | | 1.000 | | 0.492 |
| 2020 | | | | | | |
| <3 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 3 to 10 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 11+ years | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| <3 years | 1 | 0 (0.0) | 1 | 0 (0.0) | 2 | 0 (0.0) |
| 3 to 10 years | 4 | 1 (25.0) | 1 | 0 (0.0) | 6 | 2 (33.3) |
| 11+ years | 20 | 0 (0.0) | 9 | 0 (0.0) | 29 | 0 (0.0) |
| p value | | 0.200 | | -- | | 0.042 |
| 2022 | | | | | | |
| <3 years | 2 | 0 (0.0) | 0 | 0 (0.0) | 2 | 0 (0.0) |
| 3 to 10 years | 3 | 0 (0.0) | 3 | 0 (0.0) | 6 | 0 (0.0) |
| 11+ years | 60 | 1 (1.7) | 36 | 1 (2.8) | 96 | 2 (2.1) |
| p value | | 1.000 | | 1.000 | | 1.000 |

Table 8.2.6 HIV antibody prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| Victoria Receptively shared syringe last month | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| No receptive sharing | 210 | 7 (3.3) | 110 | 1 (0.9) | 320 | 8 (2.5) |
| Receptive sharing | 48 | 1 (2.1) | 21 | 0 (0.0) | 70 | 1 (1.4) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2019 | | | | | | |
| No receptive sharing | 185 | 6 (3.2) | 98 | 1 (1.0) | 286 | 7 (2.5) |
| Receptive sharing | 29 | 1 (3.5) | 21 | 0 (0.0) | 52 | 1 (1.9) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2020 | | | | | | |
| No receptive sharing | -- | -- -- | -- | -- -- | -- | -- -- |
| Receptive sharing | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No receptive sharing | 22 | 1 (4.6) | 9 | 0 (0.0) | 32 | 2 (6.3) |
| Receptive sharing | 2 | 0 (0.0) | 1 | 0 (0.0) | 3 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2022 | | | | | | |
| No receptive sharing | 45 | 1 (2.2) | 30 | 1 (3.3) | 75 | 2 (2.7) |
| Receptive sharing | 15 | 0 (0.0) | 8 | 0 (0.0) | 23 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 1.000 |

Table 8.2.7 HIV antibody prevalence by frequency of drug injection last month, gender and survey year

| Victoria | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Frequency of injection | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Less than daily | 117 | 3 (2.6) | 51 | 0 (0.0) | 169 | 3 (1.8) |
| Daily or more | 143 | 5 (3.5) | 81 | 1 (1.2) | 224 | 6 (2.7) |
| Not last month | 24 | 0 (0.0) | 11 | 0 (0.0) | 36 | 0 (0.0) |
| p value | | 0.870 | | 1.000 | | 0.882 |
| 2019 | | | | | | |
| Less than daily | 70 | 1 (1.4) | 36 | 0 (0.0) | 107 | 1 (0.9) |
| Daily or more | 145 | 6 (4.1) | 86 | 1 (1.2) | 235 | 7 (3.0) |
| Not last month | 17 | 1 (5.9) | 5 | 0 (0.0) | 22 | 1 (4.6) |
| p value | | 0.324 | | 1.000 | | 0.328 |
| 2020 | | | | | | |
| Less than daily | -- | -- -- | -- | -- -- | -- | -- -- |
| Daily or more | -- | -- -- | -- | -- -- | -- | -- -- |
| Not last month | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Less than daily | 12 | 1 (8.3) | 3 | 0 (0.0) | 15 | 1 (6.7) |
| Daily or more | 12 | 0 (0.0) | 7 | 0 (0.0) | 20 | 1 (5.0) |
| Not last month | 1 | 0 (0.0) | 1 | 0 (0.0) | 2 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2022 | | | | | | |
| Less than daily | 15 | 1 (6.7) | 15 | 0 (0.0) | 30 | 1 (3.3) |
| Daily or more | 45 | 0 (0.0) | 24 | 1 (4.2) | 69 | 1 (1.5) |
| Not last month | 5 | 0 (0.0) | 2 | 0 (0.0) | 7 | 0 (0.0) |
| p value | | 0.308 | | 1.000 | | 0.578 |

Table 8.2.8 HIV antibody prevalence by imprisonment last year, gender and survey year

| Victoria | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Imprisonment last year | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| No imprisonment | 218 | 7 (3.2) | 115 | 1 (0.9) | 334 | 8 (2.4) |
| Imprisonment | 52 | 1 (1.9) | 23 | 0 (0.0) | 75 | 1 (1.3) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2019 | | | | | | |
| No imprisonment | 162 | 5 (3.1) | 95 | 1 (1.1) | 261 | 6 (2.3) |
| Imprisonment | 55 | 2 (3.6) | 26 | 0 (0.0) | 82 | 2 (2.4) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2020 | | | | | | |
| No imprisonment | -- | -- -- | -- | -- -- | -- | -- -- |
| Imprisonment | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No imprisonment | 24 | 1 (4.2) | 7 | 0 (0.0) | 32 | 2 (6.3) |
| Imprisonment | 1 | 0 (0.0) | 1 | 0 (0.0) | 2 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2022 | | | | | | |
| No imprisonment | 54 | 1 (1.9) | 30 | 0 (0.0) | 84 | 1 (1.2) |
| Imprisonment | 8 | 0 (0.0) | 6 | 0 (0.0) | 14 | 0 (0.0) |
| p value | | 1.000 | | | | 1.000 |

Table 8.2.9 HIV antibody prevalence by condom use at last sex with regular partner, gender and survey year

| Victoria | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Condom use at last sex | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| No condom use | 73 | 2 (2.7) | 53 | 0 (0.0) | 126 | 2 (1.6) |
| Condom use | 25 | 1 (4.0) | 8 | 0 (0.0) | 33 | 1 (3.0) |
| p value | | 1.000 | | -- | | 0.505 |
| 2019 | | | | | | |
| No condom use | 57 | 1 (1.8) | 42 | 0 (0.0) | 100 | 1 (1.0) |
| Condom use | 18 | 0 (0.0) | 12 | 0 (0.0) | 32 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2020 | | | | | | |
| No condom use | -- | -- -- | -- | -- -- | -- | -- -- |
| Condom use | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No condom use | 2 | 0 (0.0) | 3 | 0 (0.0) | 5 | 0 (0.0) |
| Condom use | 0 | 0 (0.0) | 1 | 0 (0.0) | 1 | 0 (0.0) |
| p value | | -- | | -- | | -- |
| 2022 | | | | | | |
| No condom use | 14 | 1 (7.1) | 12 | 0 (0.0) | 26 | 1 (3.9) |
| Condom use | 3 | 0 (0.0) | 4 | 1 (25.0) | 7 | 1 (14.3) |
| p value | | 1.000 | | 0.250 | | 0.384 |

Table 8.2.10 HIV antibody prevalence by sex work last month, gender and survey year

| Victoria | Male | | Female | | Total | |
|---------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Sex work last month | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| No sex work | 257 | 8 (3.1) | 118 | 1 (0.9) | 377 | 9 (2.4) |
| Sex work | 13 | 0 (0.0) | 16 | 0 (0.0) | 29 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2019 | | | | | | |
| No sex work | 210 | 8 (3.8) | 107 | 1 (0.9) | 321 | 9 (2.8) |
| Sex work | 14 | 0 (0.0) | 14 | 0 (0.0) | 29 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2020 | | | | | | |
| No sex work | -- | -- -- | -- | -- -- | -- | -- -- |
| Sex work | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No sex work | 23 | 1 (4.4) | 4 | 0 (0.0) | 27 | 1 (3.7) |
| Sex work | 0 | 0 (0.0) | 4 | 0 (0.0) | 5 | 1 (20.0) |
| p value | | -- | | -- | | 0.292 |
| 2022 | | | | | | |
| No sex work | 58 | 1 (1.7) | 32 | 1 (3.1) | 90 | 2 (2.2) |
| Sex work | 4 | 0 (0.0) | 4 | 0 (0.0) | 8 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 1.000 |

Table 8.2.11 HIV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Victoria | Male | | Female | | Total | |
|---|-------------|-----------------|---------------|-----------------|--------------|-----------------|
| Aboriginal and Torres Strait Islander origin | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Non Indigenous | 240 | 4 (1.7) | 117 | 0 (0.0) | 358 | 4 (1.1) |
| Indigenous | 43 | 4 (9.3) | 27 | 1 (3.7) | 71 | 5 (7.0) |
| p value | | 0.021 | | 0.187 | | 0.008 |
| 2019 | | | | | | |
| Non Indigenous | 183 | 4 (2.2) | 102 | 0 (0.0) | 289 | 4 (1.4) |
| Indigenous | 42 | 4 (9.5) | 23 | 1 (4.4) | 65 | 5 (7.7) |
| p value | | 0.042 | | 0.184 | | 0.012 |
| 2020 | | | | | | |
| Non Indigenous | -- | -- -- | -- | -- -- | -- | -- -- |
| Indigenous | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Non Indigenous | 19 | 1 (5.3) | 10 | 0 (0.0) | 30 | 2 (6.7) |
| Indigenous | 6 | 0 (0.0) | 1 | 0 (0.0) | 7 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2022 | | | | | | |
| Non Indigenous | 55 | 1 (1.8) | 34 | 1 (2.9) | 89 | 2 (2.3) |
| Indigenous | 12 | 0 (0.0) | 6 | 0 (0.0) | 18 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 1.000 |

Table 8.2.12 HIV antibody prevalence by main language spoken at home by parents, gender and survey year

| Victoria | Male | | Female | | Total | |
|--|-------------|-----------------|---------------|-----------------|--------------|-----------------|
| Main language spoken at home by parents | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| English speaking | 254 | 8 (3.2) | 136 | 1 (0.7) | 392 | 9 (2.3) |
| Non-English speaking | 34 | 0 (0.0) | 9 | 0 (0.0) | 43 | 0 (0.0) |
| p value | | 0.602 | | 1.000 | | 0.609 |
| 2019 | | | | | | |
| English speaking | 209 | 7 (3.4) | 125 | 1 (0.8) | 338 | 8 (2.4) |
| Non-English speaking | 23 | 1 (4.4) | 2 | 0 (0.0) | 26 | 1 (3.9) |
| p value | | 0.572 | | 1.000 | | 0.491 |
| 2020 | | | | | | |
| English speaking | -- | -- -- | -- | -- -- | -- | -- -- |
| Non-English speaking | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| English speaking | 21 | 1 (4.8) | 11 | 0 (0.0) | 33 | 2 (6.1) |
| Non-English speaking | 4 | 0 (0.0) | 0 | 0 (0.0) | 4 | 0 (0.0) |
| p value | | 1.000 | | -- | | 1.000 |
| 2022 | | | | | | |
| English speaking | 55 | 1 (1.8) | 37 | 1 (2.7) | 92 | 2 (2.2) |
| Non-English speaking | 11 | 0 (0.0) | 4 | 0 (0.0) | 15 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 1.000 |

Table 8.2.13 HIV antibody prevalence by region/country of birth, gender and survey year

| Victoria | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Region/country of birth | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Australia | 243 | 8 (3.3) | 127 | 1 (0.8) | 372 | 9 (2.4) |
| Other Oceania | 9 | 0 (0.0) | 3 | 0 (0.0) | 12 | 0 (0.0) |
| Asia | 6 | 0 (0.0) | 0 | 0 (0.0) | 6 | 0 (0.0) |
| UK & Ireland | 8 | 0 (0.0) | 12 | 0 (0.0) | 20 | 0 (0.0) |
| Other | 21 | 0 (0.0) | 3 | 0 (0.0) | 24 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2019 | | | | | | |
| Australia | 206 | 6 (2.9) | 119 | 1 (0.8) | 328 | 7 (2.1) |
| Other Oceania | 2 | 0 (0.0) | 4 | 0 (0.0) | 7 | 0 (0.0) |
| Asia | 2 | 0 (0.0) | 0 | 0 (0.0) | 2 | 0 (0.0) |
| UK & Ireland | 6 | 0 (0.0) | 1 | 0 (0.0) | 7 | 0 (0.0) |
| Other | 16 | 2 (12.5) | 3 | 0 (0.0) | 19 | 2 (10.5) |
| p value | | 0.283 | | 1.000 | | 0.234 |
| 2020 | | | | | | |
| Australia | -- | -- -- | -- | -- -- | -- | -- -- |
| Other Oceania | -- | -- -- | -- | -- -- | -- | -- -- |
| Asia | -- | -- -- | -- | -- -- | -- | -- -- |
| UK & Ireland | -- | -- -- | -- | -- -- | -- | -- -- |
| Other | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Australia | 20 | 1 (0.0) | 10 | 0 (0.0) | 30 | 1 (3.3) |
| Other Oceania | 0 | 0 (0.0) | 1 | 0 (0.0) | 2 | 1 (50.0) |
| Asia | 0 | 0 (0.0) | 0 | 0 (0.0) | 0 | 0 (0.0) |
| UK & Ireland | 2 | 0 (0.0) | 0 | 0 (0.0) | 2 | 0 (0.0) |
| Other | 3 | 0 (0.0) | 0 | 0 (0.0) | 3 | 0 (0.0) |
| p value | | 1.000 | | -- | | 0.212 |
| 2022 | | | | | | |
| Australia | 51 | 1 (2.0) | 34 | 1 (2.9) | 85 | 2 (2.4) |
| Other Oceania | 1 | 0 (0.0) | 2 | 0 (0.0) | 3 | 0 (0.0) |
| Asia | 0 | 0 (0.0) | 0 | 0 (0.0) | 0 | 0 (0.0) |
| UK & Ireland | 4 | 0 (0.0) | 2 | 0 (0.0) | 6 | 0 (0.0) |
| Other | 9 | 0 (0.0) | 1 | 0 (0.0) | 10 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 1.000 |

HCV antibody prevalence

Table 8.3.1 HCV antibody prevalence by gender and survey year

| Victoria Survey year | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | 284 | 161 (57) | 145 | 81 (56) | 431 | 243 (56) |
| 2019 | 236 | 153 (65) | 126 | 76 (60) | 367 | 232 (63) |
| 2020 | -- | -- -- | -- | -- -- | -- | -- -- |
| 2021 | 25 | 8 (32) | 11 | 8 (73) | 37 | 16 (43) |
| 2022 | 64 | 19 (30) | 39 | 18 (46) | 103 | 37 (36) |
| X ² p trend | | <0.001 | | 0.510 | | <0.001 |

Table 8.3.2 HCV antibody prevalence by sexual identity, gender and survey year

| Victoria Sexual identity | Male | | Female | | Total | |
|-----------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heterosexual | 236 | 136 (58) | 101 | 58 (57) | 337 | 194 (58) |
| Bisexual | 10 | 4 (40) | 22 | 12 (55) | 32 | 16 (50) |
| Homosexual | 11 | 7 (64) | 6 | 4 (67) | 18 | 11 (61) |
| p value | | 0.471 | | 0.889 | | 0.668 |
| 2019 | | | | | | |
| Heterosexual | 196 | 129 (66) | 77 | 46 (60) | 273 | 175 (64) |
| Bisexual | 11 | 6 (55) | 22 | 12 (55) | 37 | 20 (54) |
| Homosexual | 6 | 1 (17) | 10 | 7 (70) | 16 | 8 (50) |
| p value | | 0.041 | | 0.737 | | 0.290 |
| 2020 | | | | | | |
| Heterosexual | -- | -- -- | -- | -- -- | -- | -- -- |
| Bisexual | -- | -- -- | -- | -- -- | -- | -- -- |
| Homosexual | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Heterosexual | 22 | 8 (36.4) | 4 | 4 (100.0) | 27 | 12 (44.4) |
| Bisexual | 1 | 0 (0.0) | 6 | 3 (50.0) | 7 | 3 (42.9) |
| Homosexual | 1 | 0 (0.0) | 0 | 0 (0.0) | 1 | 0 (0.0) |
| p value | | 1.000 | | 0.273 | | 1.000 |
| 2022 | | | | | | |
| Heterosexual | 56 | 16 (29) | 22 | 9 (41) | 78 | 25 (32) |
| Bisexual | 2 | 1 (50) | 10 | 2 (20) | 12 | 3 (25) |
| Homosexual | 4 | 1 (25) | 3 | 3 (100) | 7 | 4 (57) |
| p value | | 0.723 | | 0.018 | | 0.076 |

Table 8.3.3 HCV antibody prevalence by age group, gender and survey year

| Victoria Age group | Male | | Female | | Total | |
|-----------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <25 years | 2 | 0 (0) | 6 | 2 (33) | 8 | 2 (25) |
| 25-34 years | 43 | 22 (51) | 30 | 17 (57) | 73 | 39 (53) |
| 35-44 years | 115 | 69 (60) | 57 | 31 (54) | 173 | 101 (58) |
| 45+ years | 120 | 68 (57) | 51 | 31 (61) | 172 | 99 (58) |
| p value | | 0.300 | | 0.647 | | 0.294 |
| 2019 | | | | | | |
| <25 years | 3 | 2 (67) | 3 | 0 (0) | 6 | 2 (33) |
| 25-34 years | 32 | 13 (41) | 25 | 17 (68) | 59 | 30 (51) |
| 35-44 years | 98 | 56 (57) | 54 | 30 (56) | 153 | 87 (57) |
| 45+ years | 100 | 79 (79) | 43 | 29 (67) | 144 | 109 (76) |
| p value | | <0.001 | | 0.090 | | <0.001 |
| 2020 | | | | | | |
| <25 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 25-34 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 35-44 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 45+ years | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| <25 years | 0 | 0 (0.0) | 0 | 0 (0.0) | 0 | 0 (0.0) |
| 25-34 years | 0 | 0 (0.0) | 1 | 0 (0.0) | 2 | 0 (0.0) |
| 35-44 years | 6 | 2 (33.3) | 4 | 2 (50.0) | 10 | 4 (40.0) |
| 45+ years | 19 | 6 (31.6) | 6 | 6 (100.0) | 25 | 12 (48.0) |
| p value | | 1.000 | | 0.061 | | 0.542 |
| 2022 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 5 | 2 (40) | 6 | 2 (33) | 11 | 4 (36) |
| 35-44 years | 16 | 7 (44) | 17 | 8 (47) | 33 | 15 (45) |
| 45+ years | 43 | 10 (23) | 15 | 8 (53) | 58 | 18 (31) |
| p value | | 0.248 | | 0.760 | | 0.361 |

Table 8.3.4 HCV antibody prevalence by years since first injection, gender and survey year

| Victoria Years of injection | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <3 years | 12 | 2 (17) | 6 | 0 (0) | 18 | 2 (11) |
| 3 to 10 years | 24 | 13 (54) | 24 | 13 (54) | 48 | 26 (54) |
| 11+ years | 233 | 137 (59) | 107 | 64 (60) | 342 | 202 (59) |
| p value | | 0.014 | | 0.013 | | <0.001 |
| 2019 | | | | | | |
| <3 years | 6 | 1 (17) | 7 | 0 (0) | 13 | 1 (8) |
| 3 to 10 years | 23 | 10 (43) | 20 | 8 (40) | 45 | 19 (42) |
| 11+ years | 189 | 129 (68) | 91 | 62 (68) | 281 | 191 (68) |
| p value | | 0.003 | | <0.001 | | <0.001 |
| 2020 | | | | | | |
| <3 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 3 to 10 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 11+ years | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| <3 years | 1 | 0 (0.0) | 1 | 0 (0.0) | 2 | 0 (0.0) |
| 3 to 10 years | 4 | 1 (25.0) | 1 | 1 (100.0) | 6 | 2 (33.3) |
| 11+ years | 20 | 7 (35.0) | 9 | 7 (77.8) | 29 | 14 (48.3) |
| p value | | 1.000 | | 0.491 | | 0.575 |
| 2022 | | | | | | |
| <3 years | 2 | 1 (50) | 0 | 0 (0) | 2 | 1 (50) |
| 3 to 10 years | 3 | 0 (0) | 3 | 0 (0) | 6 | 0 (0) |
| 11+ years | 56 | 18 (32) | 34 | 17 (50) | 90 | 35 (39) |
| p value | | 0.588 | | 0.234 | | 0.095 |

Table 8.3.5 HCV antibody prevalence by re-use of someone else's used needle and syringe last month, years since first injection and survey year

| Victoria Receptively shared syringe last month | <3 years | | 3+ years | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| No receptive sharing | 14 | 1 (7) | 291 | 174 (60) | 315 | 180 (57) |
| Receptive sharing | 3 | 1 (33) | 60 | 39 (65) | 70 | 46 (66) |
| p value | | 0.331 | | 0.452 | | 0.188 |
| 2019 | | | | | | |
| No receptive sharing | 11 | 1 (9) | 259 | 169 (65) | 285 | 184 (65) |
| Receptive sharing | 0 | 0 (0) | 44 | 30 (68) | 52 | 35 (67) |
| p value | | -- | | 0.705 | | 0.703 |
| 2020 | | | | | | |
| No receptive sharing | -- | -- -- | -- | -- -- | -- | -- -- |
| Receptive sharing | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No receptive sharing | 2 | 0 (0.0) | 30 | 14 (46.7) | 32 | 14 (43.8) |
| Receptive sharing | 0 | 0 (0.0) | 3 | 2 (66.7) | 3 | 2 (66.7) |
| p value | | -- | | 0.601 | | 0.582 |
| 2022 | | | | | | |
| No receptive sharing | 2 | 1 (50.0) | 66 | 26 (39.3) | 71 | 27 (38.0) |
| Receptive sharing | 0 | 0 (0.0) | 19 | 3 (15.8) | 21 | 4 (19.1) |
| p value | | -- | | 0.970 | | 0.123 |

Table 8.3.6 HCV antibody prevalence by last drug injected, years since first injection and survey year

| Victoria Last drug injected | <3 years | | 3+ years | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heroin | 6 | 1 (17) | 200 | 134 (67) | 216 | 141 (65) |
| Methamphetamine | 8 | 1 (13) | 142 | 68 (48) | 156 | 72 (46) |
| Other opioids | 0 | 0 (0) | 19 | 8 (42) | 20 | 8 (40) |
| Other drugs | 4 | 0 (0) | 22 | 15 (68) | 30 | 17 (57) |
| p value | | 0.704 | | 0.001 | | 0.001 |
| 2019 | | | | | | |
| Heroin | 4 | 1 (25) | 177 | 126 (71) | 194 | 138 (71) |
| Methamphetamine | 9 | 0 (0) | 104 | 63 (61) | 121 | 69 (57) |
| Other opioids | 0 | 0 (0) | 12 | 6 (50) | 13 | 7 (54) |
| Other drugs | 0 | 0 (0) | 29 | 13 (45) | 33 | 16 (48) |
| p value | | 0.118 | | 0.017 | | 0.013 |
| 2020 | | | | | | |
| Heroin | -- | -- -- | -- | -- -- | -- | -- -- |
| Methamphetamine | -- | -- -- | -- | -- -- | -- | -- -- |
| Other opioids | -- | -- -- | -- | -- -- | -- | -- -- |
| Other drugs | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Heroin | 0 | 0 (0.0) | 15 | 6 (40.0) | 15 | 6 (40.0) |
| Methamphetamine | 2 | 0 (0.0) | 14 | 7 (50.0) | 16 | 7 (43.8) |
| Other opioids | 0 | 0 (0.0) | 2 | 1 (50.0) | 2 | 1 (50.0) |
| Other drugs | 0 | 0 (0.0) | 4 | 2 (50.0) | 4 | 2 (50.0) |
| p value | | -- | | 0.951 | | 0.981 |
| 2022 | | | | | | |
| Heroin | 0 | 0 (0.0) | 57 | 26 (45.6) | 60 | 26 (43.3) |
| Methamphetamine | 1 | 0 (0.0) | 25 | 6 (24.0) | 28 | 7 (25.0) |
| Other opioids | 1 | 1 (100.0) | 2 | 1 (50.0) | 3 | 2 (66.7) |
| Other drugs | 0 | 0 (0.0) | 9 | 2 (22.2) | 9 | 2 (22.2) |
| p value | | 0.157 | | 0.206 | | 0.192 |

Table 8.3.7 HCV antibody prevalence by frequency of drug injection last month, years since first injection and survey year

| Victoria | <3 years | | 3+ years | | Total | |
|-----------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Frequency of injection last month | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Less than daily | 11 | 0 (0) | 153 | 89 (58) | 168 | 91 (54) |
| Daily or more | 6 | 2 (33) | 201 | 125 (62) | 220 | 136 (62) |
| Not last month | 1 | 0 (0) | 31 | 12 (39) | 36 | 12 (33) |
| p value | | 0.209 | | 0.046 | | 0.005 |
| 2019 | | | | | | |
| Less than daily | 3 | 0 (0) | 98 | 58 (59) | 107 | 62 (58) |
| Daily or more | 9 | 1 (11) | 208 | 142 (68) | 234 | 158 (68) |
| Not last month | 1 | 0 (0) | 17 | 9 (53) | 23 | 11 (48) |
| p value | | 1.000 | | 0.174 | | 0.064 |
| 2020 | | | | | | |
| Less than daily | -- | -- -- | -- | -- -- | -- | -- -- |
| Daily or more | -- | -- -- | -- | -- -- | -- | -- -- |
| Not last month | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Less than daily | 0 | 0 (0.0) | 15 | 7 (46.7) | 15 | 7 (46.7) |
| Daily or more | 2 | 0 (0.0) | 18 | 9 (50.0) | 20 | 9 (45.0) |
| Not last month | 0 | 0 (0.0) | 2 | 0 (0.0) | 2 | 0 (0.0) |
| p value | | -- | | 0.641 | | 0.657 |
| 2022 | | | | | | |
| Less than daily | 1 | 0 (0.0) | 24 | 9 (37.5) | 27 | 9 (33.3) |
| Daily or more | 1 | 1 (100.0) | 62 | 21 (33.9) | 66 | 23 (34.9) |
| Not last month | 0 | 0 (0.0) | 7 | 5 (71.4) | 7 | 5 (71.4) |
| p value | | 1.000 | | 0.179 | | 0.183 |

Table 8.3.8 HCV antibody prevalence by imprisonment last year, years since first injection and survey year

| Victoria | <3 years | | 3+ years | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Imprisonment last year | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| No imprisonment | 16 | 2 (13) | 304 | 165 (54) | 332 | 173 (52) |
| Imprisonment | 2 | 0 (0) | 64 | 50 (78) | 72 | 53 (74) |
| p value | | 1.000 | | <0.001 | | 0.001 |
| 2019 | | | | | | |
| No imprisonment | 11 | 1 (9) | 236 | 140 (59) | 263 | 152 (58) |
| Imprisonment | 1 | 0 (0) | 69 | 52 (75) | 81 | 61 (75) |
| p value | | 1.000 | | 0.015 | | 0.005 |
| 2020 | | | | | | |
| No imprisonment | -- | -- -- | -- | -- -- | -- | -- -- |
| Imprisonment | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No imprisonment | 2 | 0 (0.0) | 30 | 12 (40.0) | 32 | 12 (37.5) |
| Imprisonment | 0 | 0 (0.0) | 2 | 1 (50.0) | 2 | 1 (50.0) |
| p value | | -- | | 1.000 | | 1.000 |
| 2022 | | | | | | |
| No imprisonment | 1 | 0 (0.0) | 74 | 26 (35.1) | 79 | 26 (32.9) |
| Imprisonment | 1 | 1 (100.0) | 12 | 3 (25.0) | 14 | 5 (35.7) |
| p value | | 1.000 | | 0.743 | | 1.000 |

Table 8.3.9 HCV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Victoria | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Aboriginal and Torres Strait Islander origin | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Non Indigenous | 237 | 131 (55) | 117 | 64 (55) | 355 | 195 (55) |
| Indigenous | 42 | 27 (64) | 27 | 17 (63) | 70 | 45 (64) |
| p value | | 0.277 | | 0.435 | | 0.149 |
| 2019 | | | | | | |
| Non Indigenous | 185 | 113 (61) | 100 | 62 (62) | 289 | 177 (61) |
| Indigenous | 42 | 34 (81) | 23 | 12 (52) | 65 | 46 (71) |
| p value | | 0.015 | | 0.385 | | 0.151 |
| 2020 | | | | | | |
| Non Indigenous | -- | -- -- | -- | -- -- | -- | -- -- |
| Indigenous | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Non Indigenous | 19 | 7 (36.8) | 10 | 7 (70.0) | 30 | 14 (46.7) |
| Indigenous | 6 | 1 (16.7) | 1 | 1 (100.0) | 7 | 2 (28.6) |
| p value | | 0.624 | | 1.000 | | 0.674 |
| 2022 | | | | | | |
| Non Indigenous | 51 | 13 (25.5) | 33 | 14 (42.4) | 84 | 27 (32.1) |
| Indigenous | 12 | 6 (50.0) | 5 | 3 (60.0) | 17 | 9 (52.9) |
| p value | | 0.096 | | 0.640 | | 0.103 |

Table 8.3.10 HCV antibody prevalence by main language spoken at home by parents, gender and survey year

| Victoria | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Main language spoken at home by parents | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| English speaking | 249 | 137 (55) | 136 | 75 (55) | 387 | 213 (55) |
| Non-English speaking | 34 | 24 (71) | 9 | 6 (67) | 43 | 30 (70) |
| p value | | 0.086 | | 0.731 | | 0.065 |
| 2019 | | | | | | |
| English speaking | 210 | 134 (64) | 123 | 73 (59) | 337 | 209 (62) |
| Non-English speaking | 24 | 19 (79) | 2 | 2 (100) | 27 | 22 (81) |
| p value | | 0.134 | | 0.516 | | 0.043 |
| 2020 | | | | | | |
| English speaking | -- | -- -- | -- | -- -- | -- | -- -- |
| Non-English speaking | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| English speaking | 21 | 6 (28.6) | 11 | 8 (72.7) | 33 | 14 (42.4) |
| Non-English speaking | 4 | 2 (50.0) | 0 | 0 (0.0) | 4 | 2 (50.0) |
| p value | | 0.570 | | -- | | 1.000 |
| 2022 | | | | | | |
| English speaking | 53 | 17 (32.1) | 36 | 17 (42.2) | 89 | 34 (38.2) |
| Non-English speaking | 9 | 2 (22.2) | 3 | 1 (33.3) | 12 | 3 (25.0) |
| p value | | 0.709 | | 1.000 | | 0.373 |

Table 8.3.11 HCV antibody prevalence by region/country of birth, gender and survey year

| Victoria | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Australia | 238 | 132 (55) | 127 | 72 (57) | 367 | 205 (56) |
| Other Oceania | 9 | 4 (44) | 3 | 1 (33) | 12 | 5 (42) |
| Asia | 6 | 5 (83) | 0 | 0 (0) | 6 | 5 (83) |
| UK & Ireland | 8 | 3 (38) | 12 | 6 (50) | 20 | 9 (45) |
| Other | 21 | 16 (76) | 3 | 2 (67) | 24 | 18 (75) |
| p value | | 0.140 | | 0.826 | | 0.121 |
| 2019 | | | | | | |
| Australia | 207 | 133 (64) | 117 | 71 (61) | 327 | 205 (63) |
| Other Oceania | 2 | 2 (100) | 4 | 1 (25) | 7 | 4 (57) |
| Asia | 3 | 2 (67) | 0 | 0 (0) | 3 | 2 (67) |
| UK & Ireland | 6 | 5 (83) | 1 | 1 (100) | 7 | 6 (86) |
| Other | 16 | 11 (69) | 3 | 2 (67) | 19 | 13 (68) |
| p value | | 0.855 | | 0.485 | | 0.802 |
| 2020 | | | | | | |
| Australia | -- | -- -- | -- | -- -- | -- | -- -- |
| Other Oceania | -- | -- -- | -- | -- -- | -- | -- -- |
| Asia | -- | -- -- | -- | -- -- | -- | -- -- |
| UK & Ireland | -- | -- -- | -- | -- -- | -- | -- -- |
| Other | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Australia | 20 | 7 (35.0) | 10 | 8 (80.0) | 30 | 15 (50.0) |
| Other Oceania | 0 | 0 (0.0) | 1 | 0 (0.0) | 2 | 0 (0.0) |
| Asia | 0 | 0 (0.0) | 0 | 0 (0.0) | 0 | 0 (0.0) |
| UK & Ireland | 2 | 0 (0.0) | 0 | 0 (0.0) | 2 | 0 (0.0) |
| Other | 3 | 1 (33.3) | 0 | 0 (0.0) | 3 | 1 (33.3) |
| p value | | 1.000 | | 0.273 | | 0.435 |
| 2022 | | | | | | |
| Australia | 48 | 17 (35.4) | 32 | 15 (46.9) | 80 | 32 (40.0) |
| Other Oceania | 1 | 0 (0.0) | 2 | 1 (50.0) | 3 | 1 (33.3) |
| Asia | 0 | 0 (0.0) | 0 | 0 (0.0) | 0 | 0 (0.0) |
| UK & Ireland | 4 | 1 (25.0) | 2 | 0 (0.0) | 6 | 1 (16.7) |
| Other | 8 | 1 (12.5) | 1 | 0 (0.0) | 9 | 1 (11.1) |
| p value | | 0.716 | | 0.746 | | 0.270 |

HCV RNA prevalence

Table 8.4.1 HCV RNA prevalence by gender and survey year

| Victoria | Male | | Female | | Total | |
|---------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Survey year | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | 198 | 46 (23) | 99 | 20 (20) | 299 | 66 (22) |
| 2019* | 219 | 49 (22) | 117 | 23 (20) | 341 | 73 (21) |
| 2020 | -- | -- -- | -- | -- -- | -- | -- -- |
| 2021 | 19 | 0 (0) | 8 | 4 (50) | 28 | 5 (18) |
| 2022 | 37 | 6 (16) | 29 | 2 (7) | 66 | 8 (12) |
| X ² p trend (4 year) | | 0.081 | | 0.387 | | 0.077 |

* 2018, 2019: Weighted for gender and HCV antibody status

Totals include respondents where gender was reported as other or not reported

Table 8.4.2 HCV RNA prevalence by sexual identity, gender and survey year

| Victoria | Male | | Female | | Total | |
|-----------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Sexual identity | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heterosexual | 163 | 41 (25) | 71 | 16 (23) | 234 | 57 (24) |
| Bisexual | 8 | 0 (0) | 15 | 3 (20) | 23 | 3 (13) |
| Homosexual | 6 | 2 (33) | 4 | 0 (0) | 10 | 2 (20) |
| p value | | 0.222 | | 0.548 | | 0.325 |
| 2019* | | | | | | |
| Heterosexual | 183 | 42 (23) | 71 | 15 (21) | 254 | 57 (22) |
| Bisexual | 9 | 2 (22) | 20 | 2 (10) | 33 | 6 (18) |
| Homosexual | 6 | 0 (0) | 10 | 5 (50) | 16 | 5 (31) |
| p value | | 0.394 | | 0.045 | | 0.604 |
| 2020 | | | | | | |
| Heterosexual | -- | -- -- | -- | -- -- | -- | -- -- |
| Bisexual | -- | -- -- | -- | -- -- | -- | -- -- |
| Homosexual | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Heterosexual | 16 | 0 (0) | 3 | 1 (33) | 20 | 2 (10) |
| Bisexual | 1 | 0 (0) | 4 | 2 (50) | 5 | 2 (40) |
| Homosexual | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | 0.659 | | 0.228 |
| 2022 | | | | | | |
| Heterosexual | 34 | 6 (18) | 18 | 1 (6) | 52 | 7 (13) |
| Bisexual | 1 | 0 (0) | 5 | 0 (0) | 6 | 0 (0) |
| Homosexual | 2 | 0 (0) | 2 | 1 (50) | 4 | 1 (25) |
| p value | | 1.000 | | 0.401 | | 0.729 |

Table 8.4.3 HCV RNA prevalence by age group, gender and survey year

| Victoria Age group | Male | | Female | | Total | |
|-----------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <25 years | 2 | 0 (0) | 5 | 3 (60) | 8 | 3 (38) |
| 25-34 years | 23 | 5 (22) | 23 | 5 (22) | 47 | 10 (21) |
| 35-44 years | 86 | 25 (29) | 34 | 7 (21) | 121 | 32 (26) |
| 45+ years | 84 | 15 (18) | 35 | 4 (11) | 120 | 20 (17) |
| p value | | 0.295 | | 0.077 | | 0.131 |
| 2019* | | | | | | |
| <25 years | 2 | 0 (0) | 3 | 0 (0) | 5 | 0 (0) |
| 25-34 years | 30 | 6 (20) | 22 | 3 (14) | 53 | 9 (17) |
| 35-44 years | 92 | 20 (22) | 51 | 12 (24) | 144 | 31 (22) |
| 45+ years | 92 | 22 (24) | 40 | 8 (20) | 134 | 31 (23) |
| p value | | 0.814 | | 0.656 | | 0.493 |
| 2020 | | | | | | |
| <25 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 25-34 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 35-44 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 45+ years | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 0 | 0 (0) | 1 | 0 (0) | 2 | 1 (50) |
| 35-44 years | 4 | 0 (0) | 2 | 1 (50) | 6 | 1 (17) |
| 45+ years | 15 | 0 (0) | 5 | 3 (60) | 20 | 3 (15) |
| p value | | -- | | 0.549 | | 0.466 |
| 2022 | | | | | | |
| <25 years | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| 25-34 years | 2 | 1 (50) | 3 | 1 (33) | 5 | 2 (40) |
| 35-44 years | 10 | 1 (10) | 11 | 1 (9) | 21 | 2 (10) |
| 45+ years | 25 | 4 (16) | 14 | 0 (0) | 39 | 4 (10) |
| p value | | 0.408 | | 0.095 | | 0.164 |

Table 8.4.4 HCV RNA prevalence by years since first injection, gender and survey year

| Victoria Years of injection | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <3 years | 5 | 0 (0) | 4 | 1 (25) | 9 | 1 (11) |
| 3 to 10 years | 11 | 3 (27) | 13 | 6 (46) | 24 | 9 (38) |
| 11+ years | 170 | 39 (23) | 77 | 12 (16) | 248 | 51 (21) |
| p value | | 0.442 | | 0.092 | | 0.243 |
| 2019* | | | | | | |
| <3 years | 6 | 1 (17) | 7 | 0 (0) | 13 | 1 (8) |
| 3 to 10 years | 20 | 6 (30) | 19 | 3 (16) | 40 | 10 (25) |
| 11+ years | 175 | 37 (21) | 85 | 20 (24) | 261 | 57 (22) |
| p value | | 0.639 | | 0.290 | | 0.398 |
| 2020 | | | | | | |
| <3 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 3 to 10 years | -- | -- -- | -- | -- -- | -- | -- -- |
| 11+ years | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| <3 years | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| 3 to 10 years | 3 | 0 (0) | 1 | 1 (100) | 5 | 2 (40) |
| 11+ years | 15 | 0 (0) | 7 | 3 (43) | 22 | 3 (14) |
| p value | | -- | | 0.285 | | 0.340 |
| 2022 | | | | | | |
| <3 years | 1 | 1 (100) | 0 | 0 (0) | 1 | 1 (100) |
| 3 to 10 years | 2 | 0 (0) | 2 | 0 (0) | 4 | 0 (0) |
| 11+ years | 34 | 5 (15) | 25 | 1 (4) | 59 | 6 (10) |
| p value | | 0.182 | | 1.000 | | 0.161 |

Table 8.4.5 HCV RNA prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| Victoria Receptively shared syringe last month | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| No receptive sharing | 141 | 33 (23) | 71 | 12 (17) | 212 | 45 (21) |
| Receptive sharing | 36 | 9 (25) | 18 | 8 (44) | 55 | 18 (33) |
| p value | | 0.761 | | 0.005 | | 0.065 |
| 2019* | | | | | | |
| No receptive sharing | 174 | 38 (22) | 91 | 16 (18) | 269 | 55 (20) |
| Receptive sharing | 25 | 7 (28) | 19 | 5 (26) | 45 | 13 (29) |
| p value | | 0.549 | | 0.358 | | 0.225 |
| 2020 | | | | | | |
| No receptive sharing | -- | -- -- | -- | -- -- | -- | -- -- |
| Receptive sharing | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No receptive sharing | 18 | 0 (0) | 6 | 3 (50) | 25 | 4 (16) |
| Receptive sharing | 1 | 0 (0) | 1 | 1 (100) | 2 | 1 (50) |
| p value | | -- | | 0.350 | | 0.234 |
| 2022 | | | | | | |
| No receptive sharing | 27 | 5 (19) | 22 | 0 (0) | 49 | 5 (10) |
| Receptive sharing | 7 | 0 (0) | 5 | 1 (20) | 12 | 1 (8) |
| p value | | 0.559 | | 0.185 | | 1.000 |

Table 8.4.6 HCV RNA prevalence by last drug injected, gender and survey year

| Victoria Last drug injected | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heroin | 111 | 25 (23) | 55 | 9 (16) | 167 | 34 (20) |
| Methamphetamine | 65 | 16 (25) | 27 | 8 (30) | 93 | 24 (26) |
| Other opioids | 7 | 1 (14) | 6 | 0 (0) | 14 | 1 (7) |
| Other drugs | 12 | 4 (33) | 7 | 3 (43) | 19 | 7 (37) |
| p value | | 0.800 | | 0.187 | | 0.199 |
| 2019* | | | | | | |
| Heroin | 115 | 19 (17) | 67 | 12 (18) | 183 | 30 (16) |
| Methamphetamine | 69 | 19 (28) | 38 | 9 (24) | 109 | 29 (27) |
| Other opioids | 11 | 11 (100) | 2 | 0 (0) | 13 | 5 (38) |
| Other drugs | 19 | 5 (26) | 10 | 2 (20) | 31 | 7 (23) |
| p value | | 0.088 | | 0.814 | | 0.092 |
| 2020 | | | | | | |
| Heroin | -- | -- -- | -- | -- -- | -- | -- -- |
| Methamphetamine | -- | -- -- | -- | -- -- | -- | -- -- |
| Other opioids | -- | -- -- | -- | -- -- | -- | -- -- |
| Other drugs | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Heroin | 9 | 0 (0) | 3 | 0 (0) | 13 | 1 (8) |
| Methamphetamine | 8 | 0 (0) | 4 | 3 (75) | 12 | 3 (25) |
| Other opioids | 1 | 0 (0) | 0 | 0 (0) | 1 | 0 (0) |
| Other drugs | 1 | 0 (0) | 1 | 1 (100) | 2 | 1 (50) |
| p value | | -- | | 0.082 | | 0.398 |
| 2022 | | | | | | |
| Heroin | 22 | 3 (14) | 15 | 0 (0) | 37 | 3 (8) |
| Methamphetamine | 10 | 1 (10) | 11 | 2 (18) | 21 | 3 (14) |
| Other opioids | 2 | 1 (50) | 1 | 0 (0) | 3 | 1 (33) |
| Other drugs | 2 | 1 (50) | 2 | 0 (0) | 4 | 1 (25) |
| p value | | 0.300 | | 0.319 | | 0.474 |

Table 8.4.7 HCV RNA prevalence by frequency of drug injection last month, gender and survey year

| Victoria Frequency of injection last month | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| Less than daily | 70 | 14 (20) | 33 | 6 (18) | 104 | 20 (19) |
| Daily or more | 107 | 28 (26) | 57 | 14 (25) | 164 | 42 (26) |
| Not last month | 19 | 4 (21) | 8 | 0 (0) | 27 | 4 (15) |
| p value | | 0.538 | | 0.219 | | 0.186 |
| 2019* | | | | | | |
| Less than daily | 65 | 13 (20) | 32 | 6 (19) | 96 | 19 (20) |
| Daily or more | 135 | 33 (24) | 91 | 15 (16) | 220 | 50 (23) |
| Not last month | 17 | 2 (12) | 5 | 2 (40) | 22 | 4 (18) |
| p value | | 0.409 | | 0.490 | | 0.742 |
| 2020 | | | | | | |
| Less than daily | -- | -- -- | -- | -- -- | -- | -- -- |
| Daily or more | -- | -- -- | -- | -- -- | -- | -- -- |
| Not last month | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Less than daily | 9 | 0 (0) | 2 | 2 (100) | 11 | 2 (18) |
| Daily or more | 10 | 0 (0) | 5 | 2 (40) | 16 | 3 (19) |
| Not last month | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | 0.202 | | 0.893 |
| 2022 | | | | | | |
| Less than daily | 6 | 1 (17) | 10 | 0 (0) | 16 | 1 (6) |
| Daily or more | 28 | 4 (14) | 17 | 1 (6) | 45 | 5 (11) |
| Not last month | 2 | 1 (50) | 2 | 1 (50) | 4 | 2 (50) |
| p value | | 0.346 | | 0.135 | | 0.109 |

Table 8.4.8 HCV RNA prevalence by imprisonment last year, gender and survey year

| Victoria Imprisonment last year | Male | | Female | | Total | |
|------------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018* | | | | | | |
| No imprisonment | 148 | 34 (23) | 81 | 15 (19) | 229 | 49 (21) |
| Imprisonment | 33 | 8 (24) | 13 | 4 (31) | 46 | 12 (26) |
| p value | | 0.691 | | 0.333 | | 0.370 |
| 2019* | | | | | | |
| No imprisonment | 156 | 27 (17) | 90 | 18 (20) | 250 | 46 (18) |
| Imprisonment | 48 | 16 (33) | 22 | 4 (18) | 71 | 20 (28) |
| p value | | 0.030 | | 0.865 | | 0.058 |
| 2020 | | | | | | |
| No imprisonment | -- | -- -- | -- | -- -- | -- | -- -- |
| Imprisonment | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| No imprisonment | 19 | 0 (0) | 4 | 3 (75) | 24 | 4 (17) |
| Imprisonment | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| p value | | -- | | 0.171 | | 0.656 |
| 2022 | | | | | | |
| No imprisonment | 28 | 5 (18) | 20 | 1 (5) | 48 | 6 (13) |
| Imprisonment | 6 | 1 (17) | 5 | 1 (20) | 11 | 2 (18) |
| p value | | 1.000 | | 0.367 | | 0.635 |

Table 8.4.9 HCV RNA prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Victoria | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Aboriginal and Torres Strait Islander origin | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Non Indigenous | 159 | 38 (24) | 80 | 17 (21) | 240 | 55 (23) |
| Indigenous | 37 | 7 (19) | 19 | 3 (16) | 57 | 10 (18) |
| p value | | 0.417 | | 0.840 | | 0.411 |
| 2019* | | | | | | |
| Non Indigenous | 175 | 34 (19) | 91 | 18 (20) | 270 | 53 (20) |
| Indigenous | 39 | 14 (36) | 23 | 5 (22) | 61 | 19 (31) |
| p value | | 0.035 | | 0.801 | | 0.066 |
| 2020 | | | | | | |
| Non Indigenous | -- | -- -- | -- | -- -- | -- | -- -- |
| Indigenous | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Non Indigenous | 14 | 0 (0) | 7 | 3 (43) | 22 | 4 (18) |
| Indigenous | 5 | 0 (0) | 1 | 1 (100) | 6 | 1 (17) |
| p value | | -- | | 0.285 | | 0.932 |
| 2022 | | | | | | |
| Non Indigenous | 30 | 4 (13) | 24 | 1 (4) | 54 | 5 (9) |
| Indigenous | 7 | 2 (29) | 4 | 1 (25) | 11 | 3 (27) |
| p value | | 0.315 | | 0.270 | | 0.126 |

Table 8.4.10 HCV RNA prevalence by main language spoken at home by parents, gender and survey year

| Victoria | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Main language spoken at home by parents | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| English speaking | 172 | 36 (21) | 94 | 18 (19) | 267 | 54 (20) |
| Non-English speaking | 26 | 10 (38) | 5 | 2 (40) | 31 | 12 (39) |
| p value | | 0.061 | | 0.097 | | 0.013 |
| 2019* | | | | | | |
| English speaking | 195 | 44 (23) | 114 | 21 (18) | 313 | 66 (21) |
| Non-English speaking | 23 | 5 (22) | 2 | 2 (100) | 26 | 8 (31) |
| p value | | 0.891 | | 0.004 | | 0.274 |
| 2020 | | | | | | |
| English speaking | -- | -- -- | -- | -- -- | -- | -- -- |
| Non-English speaking | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| English speaking | 16 | 0 (0) | 8 | 4 (50) | 25 | 5 (20) |
| Non-English speaking | 3 | 0 (0) | 0 | 0 (0) | 3 | 0 (0) |
| p value | | -- | | -- | | 0.393 |
| 2022 | | | | | | |
| English speaking | 31 | 5 (16) | 26 | 1 (4) | 57 | 6 (11) |
| Non-English speaking | 5 | 1 (20) | 3 | 1 (33) | 8 | 2 (25) |
| p value | | 1.000 | | 0.200 | | 0.254 |

Table 8.4.11 HCV RNA prevalence by region/country of birth, gender and survey year

| Victoria | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Australia | 168 | 34 (20) | 84 | 14 (17) | 254 | 48 (19) |
| Other Oceania | 6 | 2 (33) | 3 | 1 (33) | 10 | 3 (30) |
| Asia | 3 | 3 (100) | 0 | 0 (0) | 3 | 3 (100) |
| UK & Ireland | 4 | 0 (0) | 9 | 3 (33) | 13 | 3 (23) |
| Other | 15 | 7 (47) | 3 | 2 (67) | 18 | 9 (50) |
| p value | | 0.004 | | 0.249 | | 0.001 |
| 2019* | | | | | | |
| Australia | 192 | 42 (22) | 109 | 21 (19) | 304 | 64 (21) |
| Other Oceania | 2 | 1 (50) | 3 | 1 (33) | 6 | 3 (50) |
| Asia | 3 | 0 (0) | 0 | 0 (0) | 3 | 0 (0) |
| UK & Ireland | 6 | 2 (33) | 1 | 1 (100) | 7 | 3 (43) |
| Other | 15 | 4 (27) | 3 | 0 (0) | 18 | 4 (22) |
| p value | | 0.690 | | 0.155 | | 0.248 |
| 2020 | | | | | | |
| Australia | -- | -- -- | -- | -- -- | -- | -- -- |
| Other Oceania | -- | -- -- | -- | -- -- | -- | -- -- |
| Asia | -- | -- -- | -- | -- -- | -- | -- -- |
| UK & Ireland | -- | -- -- | -- | -- -- | -- | -- -- |
| Other | -- | -- -- | -- | -- -- | -- | -- -- |
| p value | | -- | | -- | | -- |
| 2021 | | | | | | |
| Australia | 14 | 0 (0) | 8 | 4 (50) | 22 | 4 (18) |
| Other Oceania | 0 | 0 (0) | 0 | 0 (0) | 1 | 1 (100) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 2 | 0 (0) | 0 | 0 (0) | 2 | 0 (0) |
| Other | 3 | 0 (0) | 0 | 0 (0) | 3 | 0 (0) |
| p value | | -- | | -- | | 0.128 |
| 2022 | | | | | | |
| Australia | 30 | 6 (20) | 24 | 1 (4) | 54 | 7 (13) |
| Other Oceania | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) |
| Asia | 0 | 0 (0) | 0 | 0 (0) | 0 | 0 (0) |
| UK & Ireland | 1 | 0 (0) | 2 | 0 (0) | 3 | 0 (0) |
| Other | 5 | 0 (0) | 0 | 0 (0) | 5 | 0 (0) |
| p value | | 0.634 | | 1.000 | | 1.000 |

Western Australia

Table 9.1.1 Number (percentage) of respondents by demographic characteristics and survey year

| Western Australia | 2018 | 2019 | 2020* | 2021* | 2022 |
|---|----------|----------|----------|----------|----------|
| Number of sites | 4 | 4 | 4 | 4 | 4 |
| N° surveyed | N=521 | N=528 | N=399 | N=485 | N=421 |
| Response rate (%) | 71% | 78% | 57% | 68% | 62% |
| Gender (%) | | | | | |
| Male | 309 (59) | 327 (62) | 246 (62) | 292 (60) | 265 (63) |
| Female | 211 (41) | 199 (38) | 152 (38) | 187 (39) | 151 (36) |
| Other* | 1 (<1) | 1 (<1) | 1 (<1) | 4 (1) | 4 (1) |
| Not reported | 0 (0) | 1 (<1) | 0 (0) | 2 (<1) | 1 (<1) |
| Sexual identity (%) | | | | | |
| Heterosexual | 408 (78) | 415 (79) | 311 (78) | 397 (82) | 344 (82) |
| Bisexual | 48 (9) | 39 (7) | 37 (9) | 49 (10) | 34 (8) |
| Gay/Lesbian | 16 (3) | 12 (2) | 13 (3) | 16 (3) | 12 (3) |
| Other | -- -- | -- -- | 13 (3) | 18 (4) | 27 (6) |
| Not reported | 49 (9) | 62 (12) | 25 (6) | 5 (1) | 4 (1) |
| Aboriginal and Torres Strait Islander origin (%) | | | | | |
| No | 367 (70) | 374 (71) | 258 (65) | 306 (63) | 279 (66) |
| Yes | 139 (27) | 146 (28) | 123 (31) | 150 (31) | 139 (33) |
| Not reported | 15 (3) | 8 (2) | 18 (5) | 29 (6) | 3 (1) |
| Region/country of birth (%) | | | | | |
| Australia | 427 (82) | 457 (87) | 336 (84) | 401 (83) | 365 (87) |
| Other Oceania | 21 (4) | 14 (3) | 14 (4) | 22 (5) | 12 (3) |
| Asia | 11 (2) | 4 (1) | 4 (1) | 4 (1) | 5 (1) |
| UK & Ireland | 33 (6) | 31 (6) | 24 (6) | 27 (6) | 16 (4) |
| Other | 23 (4) | 21 (4) | 14 (4) | 14 (3) | 18 (4) |
| Not reported | 6 (1) | 1 (<1) | 7 (2) | 17 (4) | 5 (1) |
| Main language spoken at home by parents (%) | | | | | |
| English | 489 (94) | 509 (96) | 381 (95) | 462 (95) | 396 (94) |
| Non-English | 26 (5) | 17 (3) | 15 (4) | 17 (4) | 22 (5) |
| Not reported | 6 (1) | 2 (<1) | 3 (1) | 6 (1) | 3 (1) |
| Age and time since first injection (years) | | | | | |
| <i>Median age</i> | 41 | 41 | 43 | 44 | 43 |
| <i>Age range</i> | 18-69 | 15-70 | 18-74 | 23-66 | 18-73 |
| Age group (%) | | | | | |
| <25 years | 30 (6) | 22 (4) | 13 (3) | 7 (1) | 11 (3) |
| 25+ years | 490 (94) | 505 (96) | 385 (96) | 472 (97) | 404 (96) |
| Not reported | 1 (<1) | 1 (<1) | 1 (<1) | 6 (1) | 6 (1) |
| Median age first injection | | | | | |
| <i>Median age first injection</i> | 19 | 19 | 19 | 19 | 19 |
| <i>Age range</i> | 11-59 | 10-66 | 10-57 | 10-85 | 10-72 |
| <i>N° not reported</i> | 20 | 31 | 11 | 17 | 27 |
| Median yrs since first injection | | | | | |
| <i>Median yrs since first injection</i> | 20 | 21 | 22 | 23 | 22 |
| <i>Range</i> | <1-51 | <1-55 | <1-52 | <1-49 | <1-51 |
| Years since first injection | | | | | |
| <3 years | 35 (7) | 28 (5) | 16 (4) | 17 (4) | 16 (4) |
| 3+ years | 465 (89) | 467 (88) | 372 (93) | 448 (92) | 378 (90) |
| Not reported | 21 (4) | 33 (6) | 11 (3) | 20 (4) | 27 (6) |
| Imprisonment last year (%) | | | | | |
| No | 404 (78) | 429 (81) | 329 (82) | 395 (81) | 343 (81) |
| Yes | 80 (15) | 69 (13) | 54 (14) | 61 (13) | 57 (14) |
| Not reported | 37 (7) | 30 (6) | 16 (4) | 29 (6) | 21 (5) |
| N° in prison | | | | | |
| <i>N° in prison</i> | N=80 | N=69 | N=54 | N=61 | N=57 |
| Injected in prison | 25 (31) | 23 (33) | 12 (22) | 14 (23) | 22 (39) |

* Implementation of the 2020 and 2021 ANSPS was impacted by the COVID-19 pandemic, see page 1 for more information

Table 9.1.2 Number (percentage) of respondents by last drug injected, frequency of injecting and survey year

| Western Australia | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|----------|----------|----------|----------|----------|
| N° surveyed | N=521 | N=528 | N=399 | N=485 | N=421 |
| Drug last injected (%) | | | | | |
| Cocaine | 1 (<1) | 0 (0) | 1 (<1) | 1 (<1) | 1 (<1) |
| Methamphetamine | 357 (69) | 352 (67) | 254 (64) | 282 (58) | 283 (67) |
| Heroin | 80 (15) | 104 (20) | 73 (18) | 86 (18) | 64 (15) |
| Pharm. opioids | 18 (3) | 17 (3) | 12 (3) | 12 (2) | 6 (1) |
| Methadone | 8 (2) | 7 (1) | 11 (3) | 21 (4) | 14 (3) |
| Buprenorphine | 9 (2) | 5 (1) | 10 (3) | 21 (4) | 8 (2) |
| Buprenorphine/naloxone | 6 (1) | 8 (2) | 8 (2) | 17 (4) | 9 (2) |
| PIEDs | 4 (1) | 7 (1) | 4 (1) | 9 (2) | 16 (4) |
| More than one | 33 (6) | 21 (4) | 20 (5) | 32 (7) | 16 (4) |
| Other | 3 (1) | 6 (1) | 1 (<1) | 3 (1) | 2 (<1) |
| Not reported | 2 (<1) | 1 (<1) | 5 (1) | 1 (<1) | 2 (<1) |
| Ever injected opioids | | | | | |
| No | 187 (36) | 181 (34) | 137 (34) | 154 (32) | 151 (36) |
| Yes | 330 (63) | 341 (65) | 261 (65) | 327 (67) | 270 (64) |
| Not reported | 4 (1) | 6 (1) | 1 (<1) | 4 (1) | 0 (0) |
| Frequency of injection last month (%) | | | | | |
| Not last month | 28 (5) | 26 (5) | 32 (8) | 40 (8) | 29 (7) |
| Less than weekly | 55 (11) | 70 (13) | 84 (21) | 98 (20) | 69 (16) |
| Weekly not daily | 104 (20) | 124 (23) | 91 (23) | 121 (25) | 85 (20) |
| Daily or more | 321 (62) | 298 (56) | 189 (47) | 215 (44) | 229 (54) |
| Not reported | 13 (3) | 10 (2) | 3 (1) | 11 (2) | 9 (2) |
| Present when someone injected for the first time in the last 12 months (%) | | | | | |
| No | 372 (71) | 401 (76) | 317 (79) | 395 (81) | 334 (79) |
| Yes | 125 (24) | 104 (20) | 77 (19) | 83 (17) | 79 (19) |
| Not reported | 24 (5) | 23 (4) | 5 (1) | 7 (1) | 8 (2) |
| Overdosed in last 12 months | | | | | |
| No | 390 (75) | 409 (77) | 346 (87) | 412 (85) | 352 (84) |
| Yes | 107 (21) | 96 (18) | 47 (12) | 63 (13) | 63 (15) |
| Not reported | 24 (5) | 23 (4) | 6 (2) | 10 (2) | 6 (1) |

Table 9.1.3 Number (percentage) of respondents by injecting behaviour in the last month and survey year

| Western Australia | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|-------------|-------------|-------------|-------------|-------------|
| N° injected last month | N=480 | N=492 | N=364 | N=434 | N=421 |
| Places injected last month (%) | | | | | |
| All private | 193 (40) | 217 (44) | 169 (46) | 183 (42) | 153 (40) |
| Any public | 286 (60) | 274 (56) | 194 (53) | 247 (57) | 228 (60) |
| Not reported | 1 (<1) | 1 (<1) | 1 (<1) | 4 (1) | 2 (1) |
| Use of new and sterile needles and syringes last month (%) | | | | | |
| All injections | 278 (58) | 322 (65) | 244 (67) | 269 (62) | 234 (61) |
| Most of the time | 146 (30) | 124 (25) | 96 (26) | 126 (29) | 113 (30) |
| Half of the time | 31 (6) | 24 (5) | 9 (2) | 14 (3) | 16 (4) |
| Some of the time | 14 (3) | 16 (3) | 10 (3) | 17 (4) | 11 (3) |
| Not last month | 5 (1) | 1 (<1) | 2 (1) | 3 (1) | 5 (1) |
| Not reported | 6 (1) | 5 (1) | 3 (1) | 5 (1) | 4 (1) |
| Re-used someone else's used needle & syringe last month (%) | | | | | |
| None | 330 (69) | 354 (72) | 276 (76) | 321 (74) | 265 (69) |
| Once | 56 (12) | 39 (8) | 30 (8) | 34 (8) | 34 (9) |
| Twice | 27 (6) | 48 (10) | 22 (6) | 27 (6) | 37 (10) |
| 3-5 times | 32 (7) | 22 (4) | 27 (7) | 22 (5) | 27 (7) |
| >5 times | 31 (6) | 21 (4) | 9 (2) | 26 (6) | 20 (5) |
| Not reported | 4 (1) | 8 (2) | 0 (0) | 4 (1) | 0 (0) |
| Number of people needle & syringe was re-used after last month (%) | | | | | |
| None | 330 (69) | 354 (72) | 276 (76) | 321 (74) | 265 (69) |
| One | 62 (13) | 72 (15) | 39 (11) | 49 (11) | 62 (16) |
| Two | 14 (3) | 14 (3) | 8 (2) | 17 (4) | 14 (4) |
| Three to five | 15 (3) | 7 (1) | 6 (2) | 11 (3) | 8 (2) |
| More than five | 8 (2) | 8 (2) | 4 (1) | 7 (2) | 5 (1) |
| Don't know | 18 (4) | 20 (4) | 21 (6) | 18 (4) | 14 (4) |
| Not reported | 33 (7) | 17 (3) | 10 (3) | 11 (3) | 15 (4) |
| Relationship to people needle & syringe was used after last month (%) [more than one could be selected] | | | | | |
| Regular sex partner | 58 (12) | 56 (11) | 39 (11) | 43 (10) | 49 (13) |
| Casual sex partner | 7 (1) | 7 (1) | 6 (2) | 6 (1) | 4 (1) |
| Close friend | 47 (10) | 25 (5) | 16 (4) | 33 (8) | 43 (11) |
| Acquaintance | 17 (4) | 15 (3) | 15 (4) | 26 (6) | 15 (4) |
| Other | 14 (3) | 15 (3) | 11 (3) | 6 (1) | 8 (2) |
| Equipment used after someone else last month (%) [more than one could be selected] | | | | | |
| Spoon | 91 (19) | 94 (19) | 56 (15) | 94 (22) | 60 (16) |
| Water | 132 (28) | 115 (23) | 86 (24) | 130 (30) | 87 (23) |
| Filter | 59 (12) | 32 (7) | 21 (6) | 46 (11) | 33 (9) |
| Drug mix | 63 (13) | 47 (10) | 36 (10) | 56 (13) | 46 (12) |
| None | 284 (59) | 304 (62) | 230 (63) | 255 (59) | 243 (63) |
| Injected by someone after that person injected themselves or others last month (%) | | | | | |
| No | 361 (75) | 379 (77) | 304 (84) | 327 (75) | 301 (79) |
| Yes | 110 (23) | 105 (21) | 60 (16) | 103 (24) | 82 (21) |
| Not reported | 9 (2) | 8 (2) | 0 (0) | 4 (1) | 0 (0) |
| Source of needle acquisition | | | | | |
| Needle Syringe Program | 374 (78) | 398 (81) | 293 (80) | 350 (81) | 295 (77) |
| Chemist/Pharmacy | 185 (39) | 177 (36) | 138 (38) | 153 (35) | 126 (33) |
| Personal sources | 72 (15) | 72 (15) | 49 (13) | 57 (13) | 54 (14) |
| Dispensing/Vending Machine | 11 (2) | 15 (3) | 3 (1) | 4 (1) | 10 (3) |
| Other sources | 21 (4) | 23 (5) | 10 (3) | 8 (2) | 11 (3) |

Table 9.1.4 Number (percentage) of respondents by treatment for drug use and survey year

| Western Australia | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|----------|----------|----------|----------|----------|
| N° surveyed | N=521 | N=528 | N=399 | N=485 | N=421 |
| Ever any treatment/therapy for drug use (%) | | | | | |
| No | 239 (49) | 225 (43) | 177 (44) | 206 (42) | 176 (42) |
| Yes | 277 (53) | 303 (57) | 222 (56) | 279 (58) | 245 (58) |
| Not reported | 5 (1) | 0 (0) | 0 (0) | 0 (0) | 0 (0) |
| History of methadone maintenance treatment (%) | | | | | |
| Currently | 56 (11) | 68 (13) | 42 (11) | 59 (12) | 46 (11) |
| Previously | 72 (14) | 83 (16) | 55 (14) | 70 (14) | 74 (18) |
| Never | 373 (72) | 365 (69) | 295 (74) | 345 (71) | 293 (70) |
| Not reported | 20 (4) | 12 (2) | 7 (2) | 11 (2) | 8 (2) |
| History of buprenorphine (Subutex®) treatment (%) | | | | | |
| Currently | 28 (5) | 34 (6) | 19 (5) | 33 (7) | 21 (5) |
| Previously | 88 (17) | 93 (18) | 51 (13) | 81 (17) | 78 (19) |
| Never | 388 (74) | 399 (76) | 323 (81) | 358 (74) | 315 (75) |
| Not reported | 17 (3) | 2 (<1) | 6 (2) | 13 (3) | 7 (2) |
| History of buprenorphine-naloxone (Suboxone®) treatment (%) | | | | | |
| Currently | 30 (6) | 41 (8) | 17 (4) | 30 (6) | 26 (6) |
| Previously | 80 (15) | 87 (16) | 52 (13) | 76 (16) | 70 (17) |
| Never | 393 (75) | 394 (75) | 327 (82) | 372 (77) | 323 (77) |
| Not reported | 18 (3) | 6 (1) | 3 (1) | 7 (1) | 2 (<1) |

Table 9.1.5 Number (percentage) of respondents by testing for HIV and HCV, HCV treatment by survey year

| Western Australia | 2018 | 2019 | 2020 | 2021 | 2022 |
|--|----------|----------|----------|----------|----------|
| N° surveyed | N=521 | N=528 | N=399 | N=485 | N=421 |
| Previous HIV test (%) | | | | | |
| Yes, ever | 389 (75) | 391 (74) | 303 (76) | 370 (76) | 311 (74) |
| Yes, last year | 252 (48) | 253 (48) | 179 (45) | 216 (45) | 187 (44) |
| >1 year ago | 137 (26) | 138 (26) | 124 (31) | 154 (32) | 124 (29) |
| Never tested | 99 (19) | 106 (20) | 83 (21) | 92 (19) | 98 (23) |
| Not reported | 33 (6) | 31 (6) | 13 (3) | 23 (5) | 12 (3) |
| Previous HCV test (%) | | | | | |
| Yes, ever | 397 (76) | 396 (75) | 315 (79) | 377 (78) | 337 (80) |
| Yes, last year | 273 (52) | 264 (50) | 188 (47) | 258 (53) | 219 (52) |
| >1 year ago | 124 (24) | 132 (25) | 127 (32) | 119 (25) | 118 (28) |
| Never tested | 64 (12) | 73 (14) | 52 (13) | 63 (13) | 44 (10) |
| Not reported | 60 (12) | 59 (11) | 32 (8) | 45 (9) | 40 (10) |
| Lifetime treatment for HCV (%)# | | | | | |
| N° self-reported HCV diagnosis | N=99 | N=103 | N=87 | N=110 | N=83 |
| Antiviral treatment | 42 (42) | 60 (58) | 45 (52) | 58 (53) | 48 (58) |
| No antiviral treatment | 55 (56) | 42 (41) | 41 (47) | 52 (47) | 35 (42) |
| Not reported | 2 (2) | 1 (1) | 1 (1) | 0 (0) | 0 (0) |
| Treatment for HCV in past 12 months (%)** | | | | | |
| N° self-reported HCV diagnosis | N=93 | N=76 | N=64 | N=86 | N=62 |
| Antiviral treatment | 28 (30) | 25 (33) | 12 (19) | 22 (26) | 17 (27) |
| No antiviral treatment | 63 (68) | 50 (66) | 51 (80) | 64 (74) | 45 (73) |
| Not reported | 2 (2) | 1 (1) | 1 (2) | 0 (0) | 0 (0) |

among people who tested HCV antibody positive and did not report spontaneous clearance

* excludes people who reported treatment induced clearance more than 12 months previously

Table 9.1.6 Number (percentage) of respondents by sexual behaviour in the month prior to survey and survey year

| Western Australia | 2018 | 2019 | 2020 | 2021 | 2022 |
|---|-------------|-------------|-------------|-------------|-------------|
| N° surveyed | N=521 | N=528 | N=399 | N=485 | N=421 |
| Sex with a regular partner last month (%) | | | | | |
| No | 235 (45) | 252 (48) | 209 (52) | 226 (47) | 203 (48) |
| Yes | 256 (49) | 249 (47) | 181 (45) | 242 (50) | 200 (48) |
| Not reported | 30 (6) | 27 (5) | 9 (2) | 17 (4) | 18 (4) |
| Condom used with regular partner last month (%) | | | | | |
| N° surveyed | N=256 | N=249 | N=181 | N=242 | N=200 |
| Never | 195 (76) | 174 (70) | 141 (78) | 192 (79) | 154 (77) |
| Sometimes | 37 (14) | 42 (17) | 16 (9) | 16 (7) | 19 (10) |
| Every time | 15 (6) | 23 (9) | 13 (7) | 20 (8) | 16 (8) |
| Not reported | 9 (4) | 10 (4) | 11 (6) | 14 (6) | 11 (6) |
| Sex with other partner(s) last month (%) | | | | | |
| No | 390 (75) | 414 (78) | 329 (82) | 389 (80) | 327 (78) |
| Yes | 99 (19) | 82 (16) | 59 (15) | 72 (15) | 77 (18) |
| Not reported | 32 (6) | 32 (6) | 11 (3) | 24 (5) | 17 (4) |
| Condom used with other partner(s) last month (%) | | | | | |
| N° surveyed | N=99 | N=82 | N=59 | N=72 | N=77 |
| Never | 50 (51) | 44 (54) | 33 (56) | 41 (57) | 42 (55) |
| Sometimes | 25 (25) | 24 (29) | 16 (27) | 14 (19) | 19 (25) |
| Every time | 20 (20) | 9 (11) | 8 (14) | 13 (18) | 12 (16) |
| Not reported | 4 (4) | 5 (6) | 2 (3) | 4 (6) | 4 (5) |
| Sex work last month (%) | | | | | |
| No | 446 (86) | 460 (87) | 364 (91) | 432 (89) | 379 (90) |
| Yes | 35 (7) | 27 (5) | 20 (5) | 26 (5) | 26 (6) |
| Not reported | 40 (8) | 41 (8) | 15 (4) | 27 (6) | 16 (4) |
| Condom used at last sex work (%) | | | | | |
| Yes | 23 (66) | 11 (41) | 5 (25) | 11 (42) | 16 (62) |

Table 9.1.7 Number (percentage) of respondents by COVID-19 related impacts, 2020-2022

| Western Australia | 2020 | 2021 | 2022 |
|---|----------|----------|----------|
| N° surveyed | N=399 | N=485 | N=421 |
| COVID-19 test | | | |
| Yes | 77 (19) | 119 (25) | 368 (87) |
| No | 265 (66) | 267 (55) | 47 (11) |
| Don't know | 26 (7) | 28 (6) | -- -- |
| Not reported | 31 (8) | 71 (15) | 6 (1) |
| Self-reported positive COVID-19 test result | | | |
| Yes | 2 (3) | 3 (1) | 172 (47) |
| Yes, last 12 months | -- -- | -- -- | 157 (43) |
| Yes, more than a year ago | -- -- | -- -- | 15 (4) |
| COVID-19 vaccine uptake | | | |
| ≥1 dose | -- -- | 182 (38) | 346 (82) |
| COVID-19 vaccine doses | | | |
| One dose | -- -- | 86 (18) | 32 (8) |
| Two doses | -- -- | 96 (20) | 121 (29) |
| Three or more doses | -- -- | -- -- | 193 (46) |
| No doses | -- -- | 236 (49) | 61 (14) |
| Not reported | -- -- | 67 (14) | 14 (3) |
| Vaccine type * | | | N=346 |
| Astra Zeneca | -- -- | 33 (7) | 82 (24) |
| Pfizer | -- -- | 106 (22) | 228 (66) |
| Don't know | -- -- | 43 (9) | 73 (21) |
| Other | -- -- | -- -- | 19 (5) |
| Not reported | -- -- | 303 (62) | -- -- |
| * 2021 only one response permitted, 2022 multiple responses permitted | | | |
| Last vaccine location | | | N=421 |
| GP or medical centre | -- -- | -- -- | 145 (34) |
| Pharmacy/ chemist | -- -- | -- -- | 80 (19) |
| Vaccine hub | -- -- | -- -- | 49 (12) |
| Opioid treatment/ methadone clinic | -- -- | -- -- | 2 (0) |
| NSP | -- -- | -- -- | 12 (3) |
| Prison | -- -- | -- -- | 23 (5) |
| Other | -- -- | -- -- | 10 (2) |
| Multiple response | -- -- | -- -- | 37 (9) |
| Not reported | -- -- | -- -- | 63 (15) |
| Vaccine hesitancy - if not had three or more doses | | | |
| Haven't got around to it | -- -- | -- -- | 97 (23) |
| Difficulty booking appointment | -- -- | -- -- | 13 (3) |
| Concerns about side effects | -- -- | -- -- | 13 (3) |
| Concerns about vaccine safety | -- -- | -- -- | 12 (3) |
| Not at risk of serious illness | -- -- | -- -- | 39 (9) |
| Other | -- -- | -- -- | 40 (10) |
| Multiple response | -- -- | -- -- | 16 (4) |
| Not reported | -- -- | -- -- | 191 (45) |

HIV antibody prevalence

Table 9.2.1 HIV antibody prevalence by gender and survey year

| Western Australia | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Survey year | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | 299 | 6 (2.0) | 207 | 4 (1.9) | 507 | 10 (2.0) |
| 2019 | 320 | 4 (1.3) | 198 | 3 (1.5) | 520 | 7 (1.4) |
| 2020 | 239 | 3 (1.3) | 152 | 2 (1.3) | 392 | 5 (1.3) |
| 2021 | 289 | 5 (1.7) | 187 | 1 (0.5) | 482 | 6 (1.2) |
| 2022 | 265 | 3 (1.1) | 151 | 1 (0.7) | 421 | 4 (1.0) |
| X ² p trend | | 0.586 | | 0.165 | | 0.206 |

Table 9.2.2 HIV antibody prevalence by sexual identity, gender and survey year

| Western Australia | Male | | Female | | Total | |
|-------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Sexual identity | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Heterosexual | 249 | 4 (1.6) | 148 | 2 (1.4) | 397 | 6 (1.5) |
| Bisexual | 16 | 1 (6.3) | 31 | 2 (6.5) | 47 | 3 (6.4) |
| Homosexual | 10 | 1 (10.0) | 5 | 0 (0.0) | 16 | 1 (6.3) |
| p value | | 0.102 | | 0.230 | | 0.037 |
| 2019 | | | | | | |
| Heterosexual | 264 | 2 (0.8) | 145 | 2 (1.4) | 409 | 4 (1.0) |
| Bisexual | 15 | 2 (13.3) | 23 | 0 (0.0) | 39 | 2 (5.1) |
| Homosexual | 9 | 0 (0.0) | 3 | 0 (0.0) | 12 | 0 (0.0) |
| p value | | 0.019 | | 1.000 | | 0.135 |
| 2020 | | | | | | |
| Heterosexual | 195 | 1 (0.5) | 111 | 2 (1.8) | 306 | 3 (1.0) |
| Bisexual | 16 | 1 (6.3) | 18 | 0 (0.0) | 35 | 1 (2.9) |
| Homosexual | 10 | 1 (10.0) | 3 | 0 (0.0) | 13 | 1 (7.7) |
| p value | | 0.058 | | 1.000 | | 0.142 |
| 2021 | | | | | | |
| Heterosexual | 258 | 3 (1.2) | 136 | 1 (0.7) | 395 | 4 (1.0) |
| Bisexual | 12 | 0 (0.0) | 36 | 0 (0.0) | 49 | 0 (0.0) |
| Homosexual | 9 | 2 (22.2) | 5 | 0 (0.0) | 15 | 2 (13.3) |
| p value | | 0.038 | | 1.000 | | 0.067 |
| 2022 | | | | | | |
| Heterosexual | 233 | 1 (0.4) | 110 | 1 (0.9) | 344 | 2 (0.6) |
| Bisexual | 12 | 0 (0.0) | 21 | 0 (0.0) | 34 | 0 (0.0) |
| Homosexual | 7 | 1 (14.3) | 5 | 0 (0.0) | 12 | 1 (8.3) |
| p value | | 0.018 | | 1.000 | | 0.058 |

Table 9.2.3 HIV antibody prevalence by age group, gender and survey year

| Western Australia Age group | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| <25 years | 15 | 0 (0.0) | 13 | 0 (0.0) | 28 | 0 (0.0) |
| 25-34 years | 54 | 1 (1.9) | 58 | 0 (0.0) | 112 | 1 (0.9) |
| 35-44 years | 99 | 1 (1.0) | 78 | 0 (0.0) | 178 | 1 (0.6) |
| 45+ years | 131 | 4 (3.1) | 57 | 4 (7.0) | 188 | 8 (4.3) |
| p value | | 0.822 | | 0.031 | | 0.081 |
| 2019 | | | | | | |
| <25 years | 10 | 0 (0.0) | 11 | 0 (0.0) | 22 | 0 (0.0) |
| 25-34 years | 54 | 1 (1.9) | 49 | 0 (0.0) | 103 | 1 (1.0) |
| 35-44 years | 124 | 1 (0.8) | 79 | 1 (1.3) | 203 | 2 (1.0) |
| 45+ years | 131 | 2 (1.5) | 59 | 2 (3.4) | 191 | 4 (2.1) |
| p value | | 0.847 | | 0.560 | | 0.718 |
| 2020 | | | | | | |
| <25 years | 3 | 0 (0.0) | 9 | 0 (0.0) | 12 | 0 (0.0) |
| 25-34 years | 37 | 0 (0.0) | 32 | 1 (3.1) | 70 | 1 (1.4) |
| 35-44 years | 80 | 1 (1.3) | 60 | 0 (0.0) | 140 | 1 (0.7) |
| 45+ years | 118 | 2 (1.7) | 51 | 1 (2.0) | 169 | 3 (1.8) |
| p value | | 1.000 | | 0.412 | | 0.870 |
| 2021 | | | | | | |
| <25 years | 4 | 1 (25.0) | 3 | 0 (0.0) | 7 | 1 (14.3) |
| 25-34 years | 31 | 1 (3.2) | 28 | 0 (0.0) | 61 | 1 (1.6) |
| 35-44 years | 103 | 0 (0.0) | 83 | 0 (0.0) | 189 | 0 (0.0) |
| 45+ years | 147 | 3 (2.0) | 71 | 1 (1.4) | 219 | 4 (1.8) |
| p value | | 0.018 | | 0.551 | | 0.016 |
| 2022 | | | | | | |
| <25 years | 8 | 0 (0.0) | 3 | 0 (0.0) | 11 | 0 (0.0) |
| 25-34 years | 42 | 1 (2.4) | 19 | 0 (0.0) | 62 | 1 (1.6) |
| 35-44 years | 98 | 1 (1.0) | 66 | 0 (0.0) | 166 | 1 (0.6) |
| 45+ years | 114 | 1 (0.9) | 60 | 1 (1.7) | 176 | 2 (1.1) |
| p value | | 0.604 | | 0.554 | | 0.827 |

Table 9.2.4 HIV antibody prevalence by last drug injected, gender and survey year

| Western Australia Last drug injected | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Heroin | 48 | 0 (0.0) | 31 | 1 (3.2) | 79 | 1 (1.3) |
| Methamphetamine | 204 | 5 (2.5) | 144 | 3 (2.1) | 348 | 8 (2.3) |
| Other opioids | 23 | 0 (0.0) | 17 | 0 (0.0) | 40 | 0 (0.0) |
| Other drugs | 24 | 1 (4.2) | 15 | 0 (0.0) | 40 | 1 (2.5) |
| p value | | 0.508 | | 0.769 | | 1.000 |
| 2019 | | | | | | |
| Heroin | 66 | 1 (1.5) | 38 | 1 (2.6) | 104 | 2 (1.9) |
| Methamphetamine | 201 | 2 (1.0) | 142 | 2 (1.4) | 344 | 4 (1.2) |
| Other opioids | 29 | 0 (0.0) | 8 | 0 (0.0) | 37 | 0 (0.0) |
| Other drugs | 24 | 1 (4.2) | 10 | 0 (0.0) | 34 | 1 (2.9) |
| p value | | 0.286 | | 0.633 | | 0.511 |
| 2020 | | | | | | |
| Heroin | 44 | 1 (2.3) | 27 | 1 (3.7) | 71 | 2 (2.8) |
| Methamphetamine | 150 | 1 (0.7) | 100 | 1 (1.0) | 250 | 2 (0.8) |
| Other opioids | 26 | 0 (0.0) | 14 | 0 (0.0) | 41 | 0 (0.0) |
| Other drugs | 17 | 1 (5.9) | 10 | 0 (0.0) | 27 | 1 (3.7) |
| p value | | 0.161 | | 0.563 | | 0.180 |
| 2021 | | | | | | |
| Heroin | 48 | 2 (4.2) | 36 | 1 (2.8) | 86 | 3 (3.5) |
| Methamphetamine | 168 | 3 (1.8) | 112 | 0 (0.0) | 281 | 3 (1.1) |
| Other opioids | 44 | 0 (0.0) | 24 | 0 (0.0) | 69 | 0 (0.0) |
| Other drugs | 29 | 0 (0.0) | 15 | 0 (0.0) | 45 | 0 (0.0) |
| p value | | 0.468 | | 0.401 | | 0.259 |
| 2022 | | | | | | |
| Heroin | 41 | 0 (0.0) | 22 | 1 (4.6) | 64 | 1 (1.6) |
| Methamphetamine | 167 | 2 (1.2) | 114 | 0 (0.0) | 283 | 2 (0.7) |
| Other opioids | 27 | 0 (0.0) | 8 | 0 (0.0) | 37 | 0 (0.0) |
| Other drugs | 30 | 1 (3.3) | 5 | 0 (0.0) | 35 | 1 (2.9) |
| p value | | 0.566 | | 0.235 | | 0.317 |

Table 9.2.5 HIV antibody prevalence by years of injection, gender and survey year

| Western Australia Years of injection | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| <3 years | 15 | 1 (6.7) | 17 | 0 (0.0) | 32 | 1 (3.1) |
| 3 to 10 years | 46 | 0 (0.0) | 46 | 0 (0.0) | 92 | 0 (0.0) |
| 11 + years | 238 | 5 (2.1) | 144 | 4 (2.8) | 383 | 9 (2.4) |
| p value | | 0.309 | | 0.698 | | 0.259 |
| 2019 | | | | | | |
| <3 years | 18 | 0 (0.0) | 10 | 0 (0.0) | 28 | 0 (0.0) |
| 3 to 10 years | 36 | 0 (0.0) | 50 | 1 (2.0) | 87 | 1 (1.2) |
| 11 + years | 248 | 4 (1.6) | 124 | 2 (1.6) | 372 | 6 (1.6) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2020 | | | | | | |
| <3 years | 7 | 0 (0.0) | 9 | 0 (0.0) | 16 | 0 (0.0) |
| 3 to 10 years | 39 | 0 (0.0) | 30 | 1 (3.3) | 70 | 1 (1.4) |
| 11 + years | 187 | 3 (1.6) | 108 | 1 (0.9) | 295 | 4 (1.4) |
| p value | | 1.000 | | 0.462 | | 1.000 |
| 2021 | | | | | | |
| <3 years | 13 | 0 (0.0) | 3 | 0 (0.0) | 17 | 0 (0.0) |
| 3 to 10 years | 38 | 3 (7.9) | 35 | 0 (0.0) | 73 | 3 (4.1) |
| 11 + years | 224 | 2 (0.9) | 144 | 1 (0.7) | 372 | 3 (0.8) |
| p value | | 0.039 | | 1.000 | | 0.099 |
| 2022 | | | | | | |
| <3 years | 11 | 0 (0.0) | 5 | 0 (0.0) | 16 | 0 (0.0) |
| 3 to 10 years | 37 | 1 (2.7) | 17 | 0 (0.0) | 55 | 1 (1.8) |
| 11 + years | 199 | 2 (1.0) | 120 | 1 (0.8) | 323 | 3 (0.9) |
| p value | | 0.479 | | 1.000 | | 0.550 |

Table 9.2.6 HIV antibody prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| Western Australia Receptively shared syringe last month | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| No receptive sharing | 193 | 4 (2.1) | 130 | 3 (2.3) | 323 | 7 (2.2) |
| Receptive sharing | 85 | 0 (0.0) | 58 | 0 (0.0) | 143 | 0 (0.0) |
| p value | | 0.317 | | 0.554 | | 0.106 |
| 2019 | | | | | | |
| No receptive sharing | 208 | 3 (1.4) | 139 | 2 (1.4) | 348 | 5 (1.4) |
| Receptive sharing | 85 | 1 (1.2) | 43 | 0 (0.0) | 128 | 1 (0.8) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2020 | | | | | | |
| No receptive sharing | 175 | 2 (1.1) | 94 | 2 (2.1) | 270 | 4 (1.5) |
| Receptive sharing | 46 | 1 (2.2) | 42 | 0 (0.0) | 88 | 1 (1.1) |
| p value | | 0.505 | | 1.000 | | 1.000 |
| 2021 | | | | | | |
| No receptive sharing | 184 | 5 (2.7) | 133 | 1 (0.8) | 318 | 6 (1.9) |
| Receptive sharing | 68 | 0 (0.0) | 38 | 0 (0.0) | 109 | 0 (0.0) |
| p value | | 0.328 | | 1.000 | | 0.345 |
| 2022 | | | | | | |
| No receptive sharing | 165 | 3 (1.8) | 98 | 0 (0.0) | 265 | 3 (1.1) |
| Receptive sharing | 74 | 0 (0.0) | 41 | 0 (0.0) | 118 | 0 (0.0) |
| p value | | 0.554 | | -- | | 0.556 |

Table 9.2.7 HIV antibody prevalence by frequency of drug injection last month, gender and survey year

| Western Australia | | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Frequency of injection | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | |
| 2018 | | | | | | | |
| Less than daily | 101 | 2 (2.0) | 56 | 2 (3.6) | 157 | 4 (2.6) | |
| Daily or more | 179 | 2 (1.1) | 134 | 2 (1.5) | 313 | 4 (1.3) | |
| Not last month | 12 | 2 (16.7) | 15 | 0 (0.0) | 27 | 2 (7.4) | |
| p value | | 0.016 | | 0.693 | | 0.093 | |
| 2019 | | | | | | | |
| Less than daily | 130 | 2 (1.5) | 61 | 2 (3.3) | 192 | 4 (2.1) | |
| Daily or more | 169 | 2 (1.2) | 123 | 1 (0.8) | 292 | 3 (1.0) | |
| Not last month | 15 | 0 (0.0) | 11 | 0 (0.0) | 26 | 0 (0.0) | |
| p value | | 1.000 | | 0.375 | | 0.615 | |
| 2020 | | | | | | | |
| Less than daily | 111 | 1 (0.9) | 61 | 1 (1.6) | 173 | 2 (1.2) | |
| Daily or more | 110 | 2 (1.8) | 75 | 1 (1.3) | 185 | 3 (1.6) | |
| Not last month | 17 | 0 (0.0) | 14 | 0 (0.0) | 31 | 0 (0.0) | |
| p value | | 0.697 | | 1.000 | | 1.000 | |
| 2021 | | | | | | | |
| Less than daily | 130 | 4 (3.1) | 84 | 0 (0.0) | 216 | 4 (1.9) | |
| Daily or more | 124 | 1 (0.8) | 89 | 1 (1.1) | 215 | 2 (0.9) | |
| Not last month | 28 | 0 (0.0) | 12 | 0 (0.0) | 40 | 0 (0.0) | |
| p value | | 0.504 | | 1.000 | | 0.816 | |
| 2022 | | | | | | | |
| Less than daily | 91 | 2 (2.2) | 60 | 0 (0.0) | 154 | 2 (1.3) | |
| Daily or more | 148 | 1 (0.7) | 79 | 0 (0.0) | 229 | 1 (0.4) | |
| Not last month | 21 | 0 (0.0) | 8 | 1 (12.5) | 29 | 1 (3.5) | |
| p value | | 0.658 | | 0.054 | | 0.259 | |

Table 9.2.8 HIV antibody prevalence by imprisonment last year, gender and survey year

| Western Australia | | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Imprisonment last year | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | |
| 2018 | | | | | | | |
| No imprisonment | 227 | 3 (1.3) | 168 | 3 (1.8) | 396 | 6 (1.5) | |
| Imprisonment | 51 | 3 (5.9) | 28 | 0 (0.0) | 79 | 3 (3.8) | |
| p value | | 0.077 | | 1.000 | | 0.176 | |
| 2019 | | | | | | | |
| No imprisonment | 252 | 1 (0.4) | 168 | 3 (1.8) | 421 | 4 (1.0) | |
| Imprisonment | 50 | 1 (2.0) | 19 | 0 (0.0) | 69 | 1 (1.5) | |
| p value | | 0.304 | | 1.000 | | 0.533 | |
| 2020 | | | | | | | |
| No imprisonment | 191 | 2 (1.1) | 131 | 2 (1.5) | 323 | 4 (1.2) | |
| Imprisonment | 40 | 1 (2.5) | 13 | 0 (0.0) | 53 | 1 (1.9) | |
| p value | | 0.436 | | 1.000 | | 0.534 | |
| 2021 | | | | | | | |
| No imprisonment | 222 | 3 (1.4) | 166 | 1 (0.6) | 392 | 4 (1.0) | |
| Imprisonment | 47 | 1 (2.1) | 13 | 0 (0.0) | 61 | 1 (1.6) | |
| p value | | 0.538 | | 1.000 | | 0.516 | |
| 2022 | | | | | | | |
| No imprisonment | 203 | 3 (1.5) | 135 | 1 (0.7) | 343 | 4 (1.2) | |
| Imprisonment | 47 | 0 (0.0) | 10 | 0 (0.0) | 57 | 0 (0.0) | |
| p value | | 1.000 | | 1.000 | | 1.000 | |

Table 9.2.9 HIV antibody prevalence by condom use at last sex with regular partner, gender and survey year

| Western Australia | | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Condom use at last sex | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | |
| 2018 | | | | | | | |
| No condom use | 103 | 2 (1.9) | 91 | 1 (1.1) | 194 | 3 (1.6) | |
| Condom use | 28 | 0 (0.0) | 21 | 0 (0.0) | 49 | 0 (0.0) | |
| p value | | 1.000 | | 1.000 | | 1.000 | |
| 2019 | | | | | | | |
| No condom use | 86 | 1 (1.2) | 86 | 0 (0.0) | 172 | 1 (0.6) | |
| Condom use | 40 | 0 (0.0) | 24 | 1 (4.2) | 64 | 1 (1.6) | |
| p value | | 1.000 | | 0.218 | | 0.470 | |
| 2020 | | | | | | | |
| No condom use | 75 | 1 (1.3) | 63 | 1 (1.6) | 138 | 2 (1.5) | |
| Condom use | 20 | 1 (5.0) | 8 | 0 (0.0) | 28 | 1 (3.6) | |
| p value | | 0.378 | | 1.000 | | 0.428 | |
| 2021 | | | | | | | |
| No condom use | 99 | 3 (3.0) | 90 | 0 (0.0) | 192 | 3 (1.6) | |
| Condom use | 19 | 0 (0.0) | 16 | 0 (0.0) | 36 | 0 (0.0) | |
| p value | | 1.000 | | -- | | 1.000 | |
| 2022 | | | | | | | |
| No condom use | 88 | 1 (1.1) | 64 | 0 (0.0) | 154 | 1 (0.7) | |
| Condom use | 22 | 0 (0.0) | 42 | 0 (0.0) | 35 | 0 (0.0) | |
| p value | | 1.000 | | -- | | 1.000 | |

Table 9.2.10 HIV antibody prevalence by sex work last month, gender and survey year

| Western Australia | | Male | | Female | | Total | |
|---------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Sex work last month | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | |
| 2018 | | | | | | | |
| No sex work | 262 | 5 (1.9) | 174 | 2 (1.2) | 436 | 7 (1.6) | |
| Sex work | 15 | 0 (0.0) | 18 | 0 (0.0) | 33 | 0 (0.0) | |
| p value | | 1.000 | | 1.000 | | 1.000 | |
| 2019 | | | | | | | |
| No sex work | 279 | 4 (1.4) | 174 | 3 (1.7) | 454 | 7 (1.5) | |
| Sex work | 14 | 0 (0.0) | 12 | 0 (0.0) | 26 | 0 (0.0) | |
| p value | | 1.000 | | 1.000 | | 1.000 | |
| 2020 | | | | | | | |
| No sex work | 223 | 3 (1.4) | 133 | 1 (0.8) | 357 | 4 (1.1) | |
| Sex work | 10 | 0 (0.0) | 10 | 1 (10.0) | 20 | 1 (5.0) | |
| p value | | 1.000 | | 0.135 | | 0.240 | |
| 2021 | | | | | | | |
| No sex work | 263 | 3 (1.1) | 162 | 1 (0.6) | 430 | 4 (0.9) | |
| Sex work | 11 | 1 (9.1) | 15 | 0 (0.0) | 26 | 1 (3.9) | |
| p value | | 0.152 | | 1.000 | | 0.255 | |
| 2022 | | | | | | | |
| No sex work | 243 | 3 (1.2) | 132 | 1 (0.8) | 379 | 4 (1.1) | |
| Sex work | 12 | 0 (0.0) | 13 | 0 (0.0) | 26 | 0 (0.0) | |
| p value | | 1.000 | | 1.000 | | 1.000 | |

Table 9.2.11 HIV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Western Australia Aboriginal and Torres Strait Islander origin | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| Non Indigenous | 225 | 3 (1.3) | 137 | 1 (0.7) | 362 | 4 (1.1) |
| Indigenous | 67 | 2 (3.0) | 65 | 2 (3.1) | 132 | 4 (3.0) |
| p value | | 0.324 | | 0.243 | | 0.219 |
| 2019 | | | | | | |
| Non Indigenous | 238 | 2 (0.8) | 131 | 1 (0.8) | 369 | 3 (0.8) |
| Indigenous | 76 | 2 (2.6) | 66 | 2 (3.0) | 144 | 4 (2.8) |
| p value | | 0.248 | | 0.260 | | 0.101 |
| 2020 | | | | | | |
| Non Indigenous | 165 | 1 (0.6) | 87 | 1 (1.2) | 252 | 2 (0.8) |
| Indigenous | 65 | 2 (3.1) | 57 | 1 (1.8) | 122 | 3 (2.5) |
| p value | | 0.193 | | 1.000 | | 0.335 |
| 2021 | | | | | | |
| Non Indigenous | 185 | 4 (2.2) | 115 | 1 (0.9) | 303 | 5 (1.7) |
| Indigenous | 80 | 1 (1.3) | 68 | 0 (0.0) | 150 | 1 (0.7) |
| p value | | 1.000 | | 1.000 | | 0.669 |
| 2022 | | | | | | |
| Non Indigenous | 190 | 2 (1.1) | 87 | 1 (1.2) | 279 | 3 (1.1) |
| Indigenous | 73 | 1 (1.4) | 63 | 0 (0.0) | 139 | 1 (0.7) |
| p value | | 1.000 | | 1.000 | | 1.000 |

Table 9.2.12 HIV antibody prevalence by main language spoken at home by parents, gender and survey year

| Western Australia Main language spoken at home by parents | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) |
| 2018 | | | | | | |
| English speaking | 282 | 6 (2.1) | 196 | 3 (1.5) | 478 | 9 (1.9) |
| Non-English speaking | 15 | 0 (0.0) | 9 | 0 (0.0) | 25 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2019 | | | | | | |
| English speaking | 306 | 4 (1.3) | 194 | 3 (1.6) | 502 | 7 (1.4) |
| Non-English speaking | 13 | 0 (0.0) | 3 | 0 (0.0) | 16 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2020 | | | | | | |
| English speaking | 227 | 3 (1.3) | 147 | 2 (1.4) | 375 | 5 (1.3) |
| Non-English speaking | 11 | 0 (0.0) | 4 | 0 (0.0) | 15 | 0 (0.0) |
| p value | | 1.000 | | 1.000 | | 1.000 |
| 2021 | | | | | | |
| English speaking | 274 | 4 (1.5) | 179 | 1 (0.6) | 459 | 5 (1.1) |
| Non-English speaking | 11 | 1 (9.1) | 6 | 0 (0.0) | 17 | 1 (5.9) |
| p value | | 0.180 | | 1.000 | | 0.197 |
| 2022 | | | | | | |
| English speaking | 248 | 2 (0.8) | 143 | 1 (0.7) | 396 | 3 (0.8) |
| Non-English speaking | 16 | 1 (6.3) | 6 | 0 (0.0) | 22 | 1 (4.6) |
| p value | | 0.172 | | 1.000 | | 0.195 |

Table 9.2.13 HIV antibody prevalence by region/country of birth, gender and survey year

| Western Australia | | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Region/country of birth | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | N° tested | N° with HIV (%) | |
| 2018 | | | | | | | |
| Australia | 238 | 6 (2.5) | 180 | 3 (1.7) | 418 | 9 (2.2) | |
| Other Oceania | 12 | 0 (0.0) | 8 | 0 (0.0) | 20 | 0 (0.0) | |
| Asia | 7 | 0 (0.0) | 4 | 0 (0.0) | 11 | 0 (0.0) | |
| UK & Ireland | 26 | 0 (0.0) | 6 | 0 (0.0) | 32 | 0 (0.0) | |
| Other | 14 | 0 (0.0) | 7 | 0 (0.0) | 22 | 0 (0.0) | |
| p value | | 1.000 | | 1.000 | | 1.000 | |
| 2019 | | | | | | | |
| Australia | 273 | 4 (1.5) | 175 | 3 (1.7) | 450 | 7 (1.6) | |
| Other Oceania | 7 | 0 (0.0) | 7 | 0 (0.0) | 14 | 0 (0.0) | |
| Asia | 4 | 0 (0.0) | 0 | 0 (0.0) | 4 | 0 (0.0) | |
| UK & Ireland | 19 | 0 (0.0) | 11 | 0 (0.0) | 30 | 0 (0.0) | |
| Other | 17 | 0 (0.0) | 4 | 0 (0.0) | 21 | 0 (0.0) | |
| p value | | 1.000 | | 1.000 | | 1.000 | |
| 2020 | | | | | | | |
| Australia | 190 | 3 (1.6) | 139 | 2 (1.4) | 330 | 5 (1.5) | |
| Other Oceania | 10 | 0 (0.0) | 4 | 0 (0.0) | 14 | 0 (0.0) | |
| Asia | 4 | 0 (0.0) | 0 | 0 (0.0) | 4 | 0 (0.0) | |
| UK & Ireland | 18 | 0 (0.0) | 6 | 0 (0.0) | 24 | 0 (0.0) | |
| Other | 12 | 0 (0.0) | 2 | 0 (0.0) | 14 | 0 (0.0) | |
| p value | | 1.000 | | 1.000 | | 1.000 | |
| 2021 | | | | | | | |
| Australia | 235 | 3 (1.3) | 158 | 1 (0.6) | 399 | 4 (1.0) | |
| Other Oceania | 11 | 0 (0.0) | 11 | 0 (0.0) | 22 | 0 (0.0) | |
| Asia | 2 | 0 (0.0) | 2 | 0 (0.0) | 4 | 0 (0.0) | |
| UK & Ireland | 19 | 0 (0.0) | 7 | 0 (0.0) | 26 | 0 (0.0) | |
| Other | 9 | 2 (22.2) | 5 | 0 (0.0) | 14 | 2 (14.3) | |
| p value | | 0.037 | | 1.000 | | 0.058 | |
| 2022 | | | | | | | |
| Australia | 228 | 2 (0.9) | 133 | 1 (0.8) | 365 | 3 (0.8) | |
| Other Oceania | 9 | 0 (0.0) | 3 | 0 (0.0) | 12 | 0 (0.0) | |
| Asia | 1 | 0 (0.0) | 4 | 0 (0.0) | 5 | 0 (0.0) | |
| UK & Ireland | 8 | 0 (0.0) | 7 | 0 (0.0) | 16 | 0 (0.0) | |
| Other | 15 | 1 (6.7) | 3 | 0 (0.0) | 18 | 1 (5.6) | |
| p value | | 0.334 | | 1.000 | | 0.409 | |

HCV antibody prevalence

Table 9.3.1 HCV antibody prevalence by gender and survey year

| Western Australia | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Survey year | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | 297 | 115 (39) | 204 | 72 (35) | 502 | 188 (37) |
| 2019 | 320 | 149 (47) | 197 | 76 (39) | 519 | 225 (43) |
| 2020 | 239 | 114 (48) | 151 | 63 (42) | 391 | 177 (45) |
| 2021 | 289 | 120 (42) | 187 | 68 (36) | 482 | 192 (40) |
| 2022 | 265 | 105 (40) | 151 | 44 (29) | 421 | 151 (36) |
| X ² p trend | | 0.751 | | 0.282 | | 0.424 |

Table 9.3.2 HCV antibody prevalence by sexual identity, gender and survey year

| Western Australia | Male | | Female | | Total | |
|-------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Sexual identity | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heterosexual | 249 | 94 (38) | 147 | 57 (39) | 396 | 151 (38) |
| Bisexual | 16 | 8 (50) | 30 | 5 (17) | 46 | 13 (28) |
| Homosexual | 10 | 3 (30) | 5 | 3 (60) | 16 | 7 (44) |
| p value | | 0.552 | | 0.024 | | 0.366 |
| 2019 | | | | | | |
| Heterosexual | 264 | 120 (45) | 144 | 55 (38) | 408 | 175 (43) |
| Bisexual | 15 | 6 (40) | 23 | 6 (26) | 39 | 12 (31) |
| Homosexual | 9 | 1 (11) | 3 | 1 (33) | 12 | 2 (17) |
| p value | | 0.113 | | 0.665 | | 0.078 |
| 2020 | | | | | | |
| Heterosexual | 196 | 95 (48) | 110 | 42 (38) | 306 | 137 (45) |
| Bisexual | 16 | 8 (50) | 18 | 11 (61) | 35 | 19 (54) |
| Homosexual | 9 | 3 (33) | 3 | 2 (67) | 12 | 5 (42) |
| p value | | 0.193 | | 0.227 | | 0.283 |
| 2021 | | | | | | |
| Heterosexual | 257 | 113 (44) | 136 | 43 (32) | 394 | 157 (40) |
| Bisexual | 12 | 4 (33) | 36 | 19 (53) | 49 | 23 (47) |
| Homosexual | 10 | 1 (10) | 5 | 2 (40) | 16 | 3 (19) |
| p value | | 0.089 | | 0.050 | | 0.378 |
| 2022 | | | | | | |
| Heterosexual | 233 | 94 (40) | 110 | 36 (33) | 344 | 131 (38) |
| Bisexual | 12 | 4 (33) | 21 | 2 (10) | 34 | 6 (18) |
| Homosexual | 7 | 2 (29) | 5 | 0 (0) | 12 | 2 (17) |
| p value | | 0.987 | | 0.079 | | 0.164 |

Table 9.3.3 HCV antibody prevalence by age group, gender and survey year

| Western Australia Age group | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| <25 years | 14 | 5 (36) | 13 | 4 (31) | 27 | 9 (33) |
| 25-34 years | 53 | 19 (36) | 57 | 16 (28) | 110 | 35 (32) |
| 35-44 years | 99 | 30 (30) | 77 | 26 (34) | 177 | 57 (32) |
| 45+ years | 131 | 61 (47) | 56 | 25 (45) | 187 | 86 (46) |
| p value | | 0.086 | | 0.311 | | 0.022 |
| 2019 | | | | | | |
| <25 years | 10 | 2 (20) | 11 | 2 (18) | 22 | 4 (18) |
| 25-34 years | 54 | 13 (24) | 49 | 18 (37) | 103 | 31 (30) |
| 35-44 years | 124 | 55 (44) | 78 | 28 (36) | 202 | 83 (41) |
| 45+ years | 131 | 79 (60) | 59 | 28 (47) | 191 | 107 (56) |
| p value | | <0.001 | | 0.258 | | <0.001 |
| 2020 | | | | | | |
| <25 years | 3 | 0 (0) | 9 | 4 (44) | 12 | 4 (33) |
| 25-34 years | 37 | 15 (41) | 32 | 9 (28) | 70 | 24 (34) |
| 35-44 years | 80 | 43 (54) | 59 | 29 (49) | 139 | 72 (52) |
| 45+ years | 118 | 55 (47) | 51 | 21 (41) | 169 | 76 (45) |
| p value | | 0.220 | | 0.281 | | 0.089 |
| 2021 | | | | | | |
| <25 years | 4 | 1 (25) | 3 | 1 (33) | 7 | 2 (29) |
| 25-34 years | 31 | 7 (23) | 28 | 9 (32) | 61 | 17 (28) |
| 35-44 years | 103 | 41 (40) | 83 | 25 (30) | 189 | 68 (36) |
| 45+ years | 147 | 71 (48) | 71 | 32 (45) | 219 | 104 (47) |
| p value | | 0.039 | | 0.236 | | 0.013 |
| 2022 | | | | | | |
| <25 years | 8 | 4 (50) | 3 | 0 (0) | 11 | 4 (36) |
| 25-34 years | 42 | 13 (31) | 19 | 4 (21) | 62 | 17 (27) |
| 35-44 years | 98 | 34 (35) | 66 | 23 (35) | 166 | 58 (35) |
| 45+ years | 114 | 53 (46) | 60 | 17 (28) | 176 | 71 (40) |
| p value | | 0.173 | | 0.533 | | 0.328 |

Table 9.3.4 HCV antibody prevalence by years since first injection, gender and survey year

| Western Australia Years of injection | Male | | Female | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| <3 years | 15 | 2 (13) | 17 | 1 (6) | 32 | 3 (9) |
| 3 to 10 years | 45 | 11 (24) | 45 | 15 (33) | 90 | 26 (29) |
| 11+ years | 227 | 96 (42) | 135 | 52 (39) | 363 | 149 (41) |
| p value | | 0.009 | | 0.019 | | <0.001 |
| 2019 | | | | | | |
| <3 years | 18 | 2 (11) | 10 | 0 (0) | 28 | 2 (7) |
| 3 to 10 years | 36 | 8 (22) | 50 | 17 (34) | 87 | 25 (29) |
| 11+ years | 248 | 132 (53) | 123 | 51 (41) | 371 | 183 (49) |
| p value | | <0.001 | | 0.021 | | <0.001 |
| 2020 | | | | | | |
| <3 years | 7 | 0 (0) | 9 | 0 (0) | 16 | 0 (0) |
| 3 to 10 years | 39 | 11 (28) | 30 | 11 (37) | 70 | 22 (31) |
| 11+ years | 187 | 97 (52) | 107 | 49 (46) | 294 | 146 (50) |
| p value | | 0.001 | | 0.014 | | <0.001 |
| 2021 | | | | | | |
| <3 years | 13 | 0 (0) | 3 | 1 (33) | 17 | 1 (6) |
| 3 to 10 years | 38 | 9 (24) | 35 | 10 (29) | 73 | 19 (26) |
| 11+ years | 224 | 107 (48) | 144 | 54 (38) | 372 | 164 (44) |
| p value | | <0.001 | | 0.746 | | <0.001 |
| 2022 | | | | | | |
| <3 years | 11 | 2 (18) | 5 | 0 (0) | 16 | 2 (13) |
| 3 to 10 years | 37 | 7 (19) | 17 | 3 (18) | 55 | 10 (18) |
| 11+ years | 199 | 88 (44) | 120 | 38 (32) | 323 | 128 (40) |
| p value | | 0.004 | | 0.265 | | 0.001 |

Table 9.3.5 HCV antibody prevalence by re-use of someone else's used needle and syringe last month, years since first injection and survey year

| Western Australia Receptively shared syringe last month | <3 years | | 3+ years | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|
| | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) |
| 2018 | | | | | | |
| No receptive sharing | 17 | 2 (12) | 296 | 100 (34) | 320 | 105 (33) |
| Receptive sharing | 9 | 1 (11) | 125 | 63 (50) | 141 | 69 (49) |
| p value | | 1.000 | | 0.001 | | 0.001 |
| 2019 | | | | | | |
| No receptive sharing | 19 | 1 (5) | 309 | 140 (45) | 348 | 151 (43) |
| Receptive sharing | 8 | 0 (0) | 111 | 51 (46) | 128 | 54 (42) |
| p value | | 1.000 | | 0.908 | | 0.814 |
| 2020 | | | | | | |
| No receptive sharing | 14 | 0 (0) | 247 | 113 (46) | 268 | 118 (44) |
| Receptive sharing | 1 | 0 (0) | 85 | 43 (51) | 88 | 45 (51) |
| p value | | -- | | 0.441 | | 0.246 |
| 2021 | | | | | | |
| No receptive sharing | 9 | 0 (0) | 298 | 116 (39) | 319 | 119 (37) |
| Receptive sharing | 2 | 1 (50) | 102 | 54 (53) | 108 | 58 (54) |
| p value | | 0.182 | | 0.013 | | 0.003 |
| 2022 | | | | | | |
| No receptive sharing | 10 | 0 (0) | 235 | 78 (33) | 265 | 85 (32) |
| Receptive sharing | 3 | 2 (67) | 110 | 51 (46) | 118 | 56 (47) |
| p value | | 0.038 | | 0.018 | | 0.004 |

Table 9.3.6 HCV antibody prevalence by last drug injected, years since first injection and survey year

| Western Australia Last drug injected | <3 years | | 3+ years | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Heroin | 0 | 0 (0) | 76 | 40 (53) | 78 | 41 (53) |
| Methamphetamine | 29 | 3 (10) | 294 | 93 (32) | 337 | 105 (31) |
| Other opioids | 0 | 0 (0) | 40 | 20 (50) | 40 | 20 (50) |
| Other drugs | 3 | 0 (0) | 33 | 17 (52) | 37 | 17 (46) |
| p value | | 0.558 | | 0.001 | | 0.001 |
| 2019 | | | | | | |
| Heroin | 1 | 1 (100) | 95 | 67 (71) | 100 | 71 (71) |
| Methamphetamine | 23 | 1 (4) | 293 | 99 (34) | 339 | 109 (32) |
| Other opioids | 1 | 0 (0) | 34 | 20 (59) | 37 | 21 (57) |
| Other drugs | 3 | 0 (0) | 29 | 19 (66) | 34 | 20 (59) |
| p value | | 0.004 | | <0.001 | | <0.001 |
| 2020 | | | | | | |
| Heroin | 1 | 0 (0) | 69 | 51 (74) | 71 | 52 (73) |
| Methamphetamine | 15 | 0 (0) | 226 | 76 (34) | 247 | 80 (32) |
| Other opioids | 0 | 0 (0) | 40 | 27 (68) | 41 | 28 (68) |
| Other drugs | 0 | 0 (0) | 24 | 12 (50) | 26 | 14 (54) |
| p value | | -- | | <0.001 | | <0.001 |
| 2021 | | | | | | |
| Heroin | 2 | 1 (50) | 82 | 45 (55) | 85 | 47 (55) |
| Methamphetamine | 8 | 0 (0) | 257 | 82 (32) | 274 | 85 (31) |
| Other opioids | 3 | 0 (0) | 62 | 33 (53) | 68 | 35 (51) |
| Other drugs | 3 | 0 (0) | 36 | 19 (53) | 44 | 20 (45) |
| p value | | 0.058 | | <0.000 | | <0.000 |
| 2022 | | | | | | |
| Heroin | 1 | 0 (0) | 61 | 25 (41) | 63 | 26 (41) |
| Methamphetamine | 9 | 2 (22) | 249 | 81 (33) | 277 | 90 (32) |
| Other opioids | 1 | 0 (0) | 31 | 22 (71) | 36 | 25 (69) |
| Other drugs | 5 | 0 (0) | 26 | 7 (27) | 34 | 7 (21) |
| p value | | 0.620 | | <0.001 | | <0.001 |

Table 9.3.7 HCV antibody prevalence by frequency of drug injection last month, years since first injection and survey year

| Western Australia | <3 years | | 3+ years | | Total | |
|-----------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Frequency of injection last month | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Less than daily | 11 | 2 (18) | 143 | 53 (37) | 157 | 57 (36) |
| Daily or more | 16 | 1 (6) | 280 | 112 (40) | 308 | 120 (39) |
| Not last month | 5 | 0 (0) | 20 | 5 (25) | 27 | 6 (22) |
| p value | | 0.734 | | 0.381 | | 0.217 |
| 2019 | | | | | | |
| Less than daily | 13 | 2 (15) | 170 | 81 (48) | 192 | 88 (46) |
| Daily or more | 15 | 0 (0) | 255 | 113 (44) | 292 | 122 (42) |
| Not last month | 0 | 0 (0) | 26 | 11 (42) | 26 | 11 (42) |
| p value | | 0.206 | | 0.753 | | 0.675 |
| 2020 | | | | | | |
| Less than daily | 7 | 0 (0) | 161 | 70 (43) | 173 | 74 (43) |
| Daily or more | 8 | 0 (0) | 171 | 86 (50) | 183 | 89 (49) |
| Not last month | 1 | 0 (0) | 30 | 11 (37) | 32 | 12 (38) |
| p value | | -- | | 0.256 | | 0.359 |
| 2021 | | | | | | |
| Less than daily | 9 | 0 (0) | 201 | 74 (37) | 217 | 77 (35) |
| Daily or more | 3 | 1 (33) | 201 | 96 (48) | 214 | 101 (47) |
| Not last month | 4 | 0 (0) | 35 | 9 (26) | 40 | 9 (23) |
| p value | | 0.187 | | 0.013 | | 0.003 |
| 2022 | | | | | | |
| Less than daily | 8 | 0 (0) | 140 | 46 (33) | 154 | 49 (32) |
| Daily or more | 5 | 2 (40) | 205 | 83 (40) | 229 | 92 (40) |
| Not last month | 3 | 0 (0) | 24 | 7 (29) | 29 | 8 (28) |
| p value | | 0.108 | | 0.255 | | 0.151 |

Table 9.3.8 HCV antibody prevalence by imprisonment last year, years since first injection and survey year

| Western Australia | <3 years | | 3+ years | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Imprisonment last year | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| No imprisonment | 26 | 3 (12) | 357 | 137 (38) | 392 | 146 (37) |
| Imprisonment | 4 | 0 (0) | 69 | 28 (41) | 79 | 31 (39) |
| p value | | 1.000 | | 0.731 | | 0.738 |
| 2019 | | | | | | |
| No imprisonment | 24 | 1 (4) | 373 | 163 (44) | 420 | 175 (42) |
| Imprisonment | 3 | 0 (0) | 58 | 30 (52) | 69 | 34 (49) |
| p value | | 1.000 | | 0.253 | | 0.236 |
| 2020 | | | | | | |
| No imprisonment | 14 | 0 (0) | 299 | 134 (45) | 321 | 140 (44) |
| Imprisonment | 2 | 0 (0) | 49 | 26 (53) | 54 | 29 (54) |
| p value | | -- | | 0.283 | | 0.168 |
| 2021 | | | | | | |
| No imprisonment | 15 | 1 (7) | 364 | 147 (40) | 392 | 151 (39) |
| Imprisonment | 2 | 0 (0) | 55 | 27 (49) | 61 | 30 (49) |
| p value | | 1.000 | | 0.242 | | 0.114 |
| 2022 | | | | | | |
| No imprisonment | 15 | 1 (7) | 310 | 106 (34) | 343 | 115 (34) |
| Imprisonment | 1 | 1 (100) | 50 | 25 (50) | 57 | 29 (51) |
| p value | | 0.125 | | 0.031 | | 0.012 |

Table 9.3.9 HCV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Western Australia Aboriginal and Torres Strait Islander origin | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| Non Indigenous | 224 | 76 (34) | 136 | 42 (31) | 360 | 118 (33) |
| Indigenous | 66 | 36 (55) | 64 | 28 (44) | 130 | 64 (49) |
| p value | | 0.002 | | 0.075 | | 0.001 |
| 2019 | | | | | | |
| Non Indigenous | 238 | 107 (45) | 131 | 49 (37) | 369 | 156 (42) |
| Indigenous | 76 | 40 (53) | 65 | 27 (42) | 143 | 67 (47) |
| p value | | 0.243 | | 0.576 | | 0.349 |
| 2020 | | | | | | |
| Non Indigenous | 165 | 66 (40) | 86 | 32 (37) | 251 | 98 (39) |
| Indigenous | 65 | 40 (62) | 57 | 26 (46) | 122 | 66 (54) |
| p value | | 0.003 | | 0.316 | | 0.006 |
| 2021 | | | | | | |
| Non Indigenous | 185 | 63 (34) | 115 | 37 (32) | 303 | 101 (33) |
| Indigenous | 80 | 43 (54) | 68 | 30 (44) | 150 | 75 (50) |
| p value | | 0.003 | | 0.105 | | 0.001 |
| 2022 | | | | | | |
| Non Indigenous | 190 | 70 (37) | 87 | 25 (29) | 279 | 95 (34) |
| Indigenous | 73 | 34 (47) | 63 | 19 (30) | 139 | 55 (40) |
| p value | | 0.148 | | 0.850 | | 0.268 |

Table 9.3.10 HCV antibody prevalence by main language spoken at home by parents, gender and survey year

| Western Australia Main language spoken at home by parents | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018 | | | | | | |
| English speaking | 280 | 107 (38) | 194 | 68 (35) | 474 | 175 (37) |
| Non-English speaking | 15 | 7 (47) | 9 | 4 (44) | 25 | 12 (48) |
| p value | | 0.512 | | 0.723 | | 0.265 |
| 2019 | | | | | | |
| English speaking | 306 | 140 (46) | 193 | 75 (39) | 501 | 215 (43) |
| Non-English speaking | 13 | 9 (69) | 3 | 1 (33) | 16 | 10 (63) |
| p value | | 0.154 | | 1.000 | | 0.120 |
| 2020 | | | | | | |
| English speaking | 227 | 110 (48) | 146 | 60 (41) | 374 | 170 (45) |
| Non-English speaking | 11 | 4 (36) | 4 | 3 (75) | 15 | 7 (47) |
| p value | | 0.543 | | 0.310 | | 0.926 |
| 2021 | | | | | | |
| English speaking | 274 | 113 (41) | 179 | 64 (36) | 459 | 181 (39) |
| Non-English speaking | 11 | 4 (36) | 6 | 3 (50) | 17 | 7 (41) |
| p value | | 1.000 | | 0.669 | | 0.885 |
| 2022 | | | | | | |
| English speaking | 248 | 96 (39) | 143 | 42 (29) | 396 | 140 (35) |
| Non-English speaking | 16 | 8 (50) | 6 | 2 (33) | 22 | 10 (45) |
| p value | | 0.370 | | 1.000 | | 0.336 |

Table 9.3.11 HCV antibody prevalence by region/country of birth, gender and survey year

| Western Australia | | Male | | Female | | Total | |
|--------------------------------|------------------|------------------------|------------------|------------------------|------------------|------------------------|--|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | |
| 2018 | | | | | | | |
| Australia | 236 | 84 (36) | 178 | 64 (36) | 414 | 148 (36) | |
| Other Oceania | 12 | 5 (42) | 8 | 0 (0) | 20 | 5 (25) | |
| Asia | 7 | 4 (57) | 4 | 4 (100) | 11 | 8 (73) | |
| UK & Ireland | 26 | 12 (46) | 6 | 1 (17) | 32 | 13 (41) | |
| Other | 14 | 9 (64) | 7 | 3 (43) | 22 | 13 (59) | |
| p value | | 0.158 | | 0.008 | | 0.018 | |
| 2019 | | | | | | | |
| Australia | 273 | 126 (46) | 174 | 66 (38) | 449 | 192 (43) | |
| Other Oceania | 7 | 2 (29) | 7 | 4 (57) | 14 | 6 (43) | |
| Asia | 4 | 1 (25) | 0 | 0 (0) | 4 | 1 (25) | |
| UK & Ireland | 19 | 10 (53) | 11 | 5 (45) | 30 | 15 (50) | |
| Other | 17 | 10 (59) | 4 | 0 (0) | 21 | 10 (48) | |
| p value | | 0.587 | | 0.318 | | 0.878 | |
| 2020 | | | | | | | |
| Australia | 190 | 93 (49) | 139 | 60 (43) | 330 | 153 (46) | |
| Other Oceania | 10 | 4 (40) | 4 | 1 (25) | 14 | 5 (36) | |
| Asia | 4 | 1 (25) | 0 | 0 (0) | 4 | 1 (25) | |
| UK & Ireland | 18 | 8 (44) | 5 | 1 (20) | 23 | 9 (39) | |
| Other | 12 | 6 (50) | 2 | 1 (50) | 14 | 7 (50) | |
| p value | | 0.909 | | 0.711 | | 0.809 | |
| 2021 | | | | | | | |
| Australia | 235 | 98 (42) | 158 | 61 (39) | 399 | 163 (41) | |
| Other Oceania | 11 | 6 (55) | 11 | 3 (27) | 22 | 9 (41) | |
| Asia | 2 | 0 (0) | 2 | 0 (0) | 4 | 0 (0) | |
| UK & Ireland | 19 | 6 (32) | 7 | 0 (0) | 26 | 6 (23) | |
| Other | 9 | 4 (44) | 5 | 2 (40) | 14 | 6 (43) | |
| p value | | 0.654 | | 0.194 | | 0.219 | |
| 2022 | | | | | | | |
| Australia | 228 | 85 (37) | 133 | 39 (29) | 365 | 126 (35) | |
| Other Oceania | 9 | 6 (67) | 3 | 1 (33) | 12 | 7 (58) | |
| Asia | 1 | 0 (0) | 4 | 1 (25) | 5 | 1 (20) | |
| UK & Ireland | 8 | 3 (38) | 7 | 1 (14) | 16 | 4 (25) | |
| Other | 15 | 10 (67) | 3 | 1 (33) | 18 | 11 (61) | |
| p value | | 0.059 | | 0.973 | | 0.054 | |

HCV RNA prevalence

Table 9.4.1 HCV RNA prevalence by gender and survey year

| Western Australia | Male | | Female | | Total | |
|------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Survey year | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | 67 | 17 (25) | 48 | 10 (21) | 115 | 27 (23) |
| 2019* | 313 | 63 (20) | 138 | 23 (17) | 453 | 86 (19) |
| 2020 | 219 | 59 (27) | 132 | 31 (23) | 351 | 90 (26) |
| 2021 | 286 | 75 (26) | 177 | 27 (15) | 469 | 103 (22) |
| 2022 | 260 | 56 (22) | 150 | 17 (11) | 415 | 74 (18) |
| X ² p trend | | 0.766 | | 0.057 | | 0.386 |

* 2018, 2019: Weighted for gender and HCV antibody status

Totals include respondents where gender was reported as other or not reported

Table 9.4.2 HCV RNA prevalence by sexual identity, gender and survey year

| Western Australia | Male | | Female | | Total | |
|-------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| Sexual identity | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heterosexual | 56 | 15 (27) | 35 | 8 (23) | 91 | 24 (26) |
| Bisexual | 2 | 1 (50) | 6 | 0 (0) | 8 | 1 (13) |
| Homosexual | 4 | 0 (0) | 1 | 0 (0) | 5 | 0 (0) |
| p value | | 0.311 | | 0.355 | | 0.362 |
| 2019* | | | | | | |
| Heterosexual | 261 | 50 (19) | 102 | 16 (16) | 363 | 65 (18) |
| Bisexual | 15 | 2 (13) | 18 | 3 (17) | 34 | 4 (12) |
| Homosexual | 11 | 1 (9) | 1 | 0 (0) | 12 | 1 (8) |
| p value | | 0.479 | | 0.909 | | 0.443 |
| 2020 | | | | | | |
| Heterosexual | 177 | 49 (28) | 95 | 27 (28) | 272 | 76 (28) |
| Bisexual | 16 | 5 (31) | 15 | 1 (7) | 31 | 6 (19) |
| Homosexual | 9 | 2 (22) | 3 | 0 (0) | 12 | 2 (17) |
| p value | | 0.889 | | 0.116 | | 0.430 |
| 2021 | | | | | | |
| Heterosexual | 255 | 70 (27) | 128 | 18 (14) | 384 | 88 (23) |
| Bisexual | 12 | 2 (17) | 35 | 5 (14) | 48 | 7 (15) |
| Homosexual | 9 | 1 (11) | 5 | 2 (40) | 15 | 3 (20) |
| p value | | 0.404 | | 0.277 | | 0.414 |
| 2022 | | | | | | |
| Heterosexual | 228 | 48 (21) | 109 | 13 (12) | 338 | 61 (18) |
| Bisexual | 12 | 2 (17) | 21 | 1 (5) | 34 | 3 (9) |
| Homosexual | 7 | 2 (29) | 5 | 0 (0) | 12 | 2 (17) |
| p value | | 0.720 | | 0.227 | | 0.183 |

Table 9.4.3 HCV RNA prevalence by age group, gender and survey year

| Western Australia Age group | Male | | Female | | Total | |
|--------------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <25 years | 1 | 0 (0) | 2 | 1 (50) | 3 | 1 (33) |
| 25-34 years | 7 | 0 (0) | 15 | 5 (33) | 22 | 5 (23) |
| 35-44 years | 27 | 5 (19) | 21 | 2 (10) | 48 | 7 (15) |
| 45+ years | 32 | 12 (38) | 10 | 2 (20) | 42 | 14 (33) |
| p value | | 0.134 | | 0.461 | | 0.294 |
| 2019* | | | | | | |
| <25 years | 13 | 1 (8) | 10 | 2 (20) | 24 | 3 (13) |
| 25-34 years | 65 | 8 (12) | 35 | 5 (14) | 99 | 13 (13) |
| 35-44 years | 117 | 21 (18) | 51 | 7 (14) | 168 | 27 (16) |
| 45+ years | 117 | 33 (28) | 42 | 10 (24) | 160 | 43 (27) |
| p value | | 0.017 | | 0.617 | | 0.009 |
| 2020 | | | | | | |
| <25 years | 3 | 0 (0) | 9 | 3 (33) | 12 | 3 (25) |
| 25-34 years | 32 | 7 (22) | 25 | 3 (12) | 57 | 10 (18) |
| 35-44 years | 74 | 27 (36) | 53 | 15 (28) | 127 | 42 (33) |
| 45+ years | 109 | 25 (23) | 45 | 10 (22) | 154 | 35 (23) |
| p value | | 0.121 | | 0.385 | | 0.097 |
| 2021 | | | | | | |
| <25 years | 4 | 1 (25) | 3 | 1 (33) | 7 | 2 (29) |
| 25-34 years | 30 | 5 (17) | 26 | 5 (19) | 58 | 11 (19) |
| 35-44 years | 101 | 27 (27) | 76 | 8 (11) | 180 | 35 (19) |
| 45+ years | 147 | 42 (29) | 70 | 13 (19) | 218 | 55 (25) |
| p value | | 0.612 | | 0.399 | | 0.482 |
| 2022 | | | | | | |
| <25 years | 8 | 2 (25) | 3 | 0 (0) | 11 | 2 (18) |
| 25-34 years | 42 | 7 (17) | 19 | 1 (5) | 62 | 8 (13) |
| 35-44 years | 96 | 19 (20) | 66 | 7 (11) | 164 | 26 (16) |
| 45+ years | 111 | 28 (25) | 59 | 9 (15) | 172 | 38 (22) |
| p value | | 0.641 | | 0.694 | | 0.311 |

Table 9.4.4 HCV RNA prevalence by years since first injection, gender and survey year

| Western Australia Years of injection | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| <3 years | 2 | 0 (0) | 4 | 2 (50) | 6 | 2 (33) |
| 3 to 10 years | 9 | 1 (11) | 10 | 3 (30) | 19 | 4 (21) |
| 11+ years | 55 | 14 (25) | 34 | 5 (15) | 89 | 20 (22) |
| p value | | 0.519 | | 0.291 | | 0.795 |
| 2019* | | | | | | |
| <3 years | 24 | 1 (4) | 6 | 0 (0) | 30 | 2 (7) |
| 3 to 10 years | 42 | 7 (17) | 36 | 9 (25) | 80 | 16 (20) |
| 11+ years | 229 | 50 (22) | 86 | 11 (13) | 315 | 61 (19) |
| p value | | 0.219 | | 0.140 | | 0.156 |
| 2020 | | | | | | |
| <3 years | 6 | 0 (0) | 6 | 0 (0) | 12 | 0 (0) |
| 3 to 10 years | 36 | 8 (22) | 26 | 5 (19) | 62 | 13 (21) |
| 11+ years | 172 | 48 (28) | 95 | 25 (26) | 267 | 73 (27) |
| p value | | 0.261 | | 0.284 | | 0.071 |
| 2021 | | | | | | |
| <3 years | 13 | 0 (0) | 3 | 0 (0) | 17 | 0 (0) |
| 3 to 10 years | 36 | 11 (31) | 31 | 6 (19) | 67 | 17 (25) |
| 11+ years | 223 | 62 (28) | 138 | 18 (13) | 365 | 80 (22) |
| p value | | 0.077 | | 0.513 | | 0.072 |
| 2022 | | | | | | |
| <3 years | 11 | 1 (9) | 5 | 0 (0) | 16 | 1 (6) |
| 3 to 10 years | 36 | 5 (14) | 16 | 2 (13) | 53 | 7 (13) |
| 11+ years | 195 | 44 (23) | 120 | 14 (12) | 319 | 59 (19) |
| p value | | 0.421 | | 1.000 | | 0.377 |

Table 9.4.5 HCV RNA prevalence by re-use of someone else's used needle and syringe last month, gender and survey year

| Western Australia Receptively shared syringe last month | Male | | Female | | Total | |
|--|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| No receptive sharing | 44 | 7 (16) | 36 | 7 (19) | 80 | 14 (18) |
| Receptive sharing | 17 | 8 (47) | 4 | 2 (50) | 21 | 10 (48) |
| p value | | 0.018 | | 0.178 | | 0.005 |
| 2019* | | | | | | |
| No receptive sharing | 199 | 48 (24) | 94 | 14 (15) | 294 | 62 (21) |
| Receptive sharing | 89 | 12 (13) | 33 | 8 (24) | 122 | 20 (16) |
| p value | | 0.040 | | 0.207 | | 0.261 |
| 2020 | | | | | | |
| No receptive sharing | 161 | 40 (25) | 77 | 19 (25) | 238 | 59 (25) |
| Receptive sharing | 39 | 16 (41) | 40 | 9 (23) | 79 | 25 (32) |
| p value | | 0.043 | | 0.794 | | 0.232 |
| 2021 | | | | | | |
| No receptive sharing | 182 | 46 (25) | 125 | 16 (13) | 308 | 62 (20) |
| Receptive sharing | 67 | 23 (34) | 36 | 9 (25) | 106 | 32 (30) |
| p value | | 0.157 | | 0.075 | | 0.033 |
| 2022 | | | | | | |
| No receptive sharing | 163 | 32 (20) | 97 | 7 (7) | 262 | 39 (15) |
| Receptive sharing | 71 | 22 (31) | 41 | 9 (22) | 115 | 32 (28) |
| p value | | 0.058 | | 0.013 | | 0.003 |

Table 9.4.6 HCV RNA prevalence by last drug injected, gender and survey year

| Western Australia Last drug injected | Male | | Female | | Total | |
|---|-----------|-----------------|-----------|-----------------|-----------|-----------------|
| | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) |
| 2018* | | | | | | |
| Heroin | 17 | 3 (18) | 9 | 2 (22) | 26 | 5 (19) |
| Methamphetamine | 38 | 11 (29) | 28 | 4 (14) | 66 | 15 (23) |
| Other opioids | 7 | 2 (29) | 4 | 1 (25) | 11 | 3 (27) |
| Other drugs | 3 | 0 (0) | 5 | 3 (60) | 8 | 3 (38) |
| p value | | 0.657 | | 0.251 | | 0.849 |
| 2019* | | | | | | |
| Heroin | 50 | 19 (38) | 22 | 8 (36) | 71 | 27 (38) |
| Methamphetamine | 209 | 37 (18) | 100 | 13 (13) | 311 | 50 (16) |
| Other opioids | 27 | 6 (22) | 7 | 1 (14) | 34 | 7 (21) |
| Other drugs | 23 | 1 (4) | 8 | 2 (25) | 30 | 3 (10) |
| p value | | 0.002 | | 0.048 | | <0.001 |
| 2020 | | | | | | |
| Heroin | 41 | 13 (32) | 26 | 10 (38) | 67 | 23 (34) |
| Methamphetamine | 137 | 27 (20) | 81 | 14 (17) | 218 | 41 (19) |
| Other opioids | 23 | 12 (52) | 14 | 3 (21) | 37 | 15 (41) |
| Other drugs | 15 | 5 (33) | 9 | 4 (44) | 24 | 9 (38) |
| p value | | 0.007 | | 0.068 | | 0.003 |
| 2021 | | | | | | |
| Heroin | 48 | 14 (29) | 33 | 7 (21) | 83 | 21 (25) |
| Methamphetamine | 159 | 44 (28) | 104 | 13 (13) | 264 | 57 (22) |
| Other opioids | 43 | 13 (30) | 23 | 3 (13) | 67 | 16 (24) |
| Other drugs | 29 | 1 (3) | 15 | 4 (27) | 44 | 5 (11) |
| p value | | 0.035 | | 0.379 | | 0.308 |
| 2022 | | | | | | |
| Heroin | 40 | 4 (10) | 21 | 2 (10) | 62 | 6 (10) |
| Methamphetamine | 160 | 39 (24) | 110 | 13 (12) | 272 | 53 (19) |
| Other opioids | 26 | 8 (31) | 8 | 1 (13) | 36 | 9 (25) |
| Other drugs | 29 | 4 (14) | 5 | 0 (0) | 34 | 4 (12) |
| p value | | 0.100 | | 0.861 | | 0.142 |

Table 9.4.7 HCV RNA prevalence by frequency of drug injection last month, gender and survey year

| Western Australia | | Male | | Female | | Total | |
|-----------------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|--|
| Frequency of injection last month | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | |
| 2018* | | | | | | | |
| Less than daily | 15 | 5 (33) | 16 | 4 (25) | 31 | 9 (29) | |
| Daily or more | 46 | 11 (24) | 24 | 3 (13) | 70 | 14 (20) | |
| Not last month | 5 | 1 (20) | 6 | 2 (33) | 11 | 3 (27) | |
| p value | | 0.820 | | 0.554 | | 0.579 | |
| 2019* | | | | | | | |
| Less than daily | 130 | 25 (19) | 44 | 11 (25) | 175 | 36 (21) | |
| Daily or more | 164 | 36 (22) | 85 | 11 (13) | 249 | 47 (19) | |
| Not last month | 14 | 2 (14) | 8 | 1 (13) | 22 | 3 (14) | |
| p value | | 0.591 | | 0.155 | | 0.571 | |
| 2020 | | | | | | | |
| Less than daily | 105 | 25 (24) | 54 | 13 (24) | 159 | 38 (24) | |
| Daily or more | 95 | 31 (33) | 63 | 15 (24) | 158 | 46 (29) | |
| Not last month | 18 | 2 (11) | 13 | 3 (23) | 31 | 5 (16) | |
| p value | | 0.111 | | 0.997 | | 0.256 | |
| 2021 | | | | | | | |
| Less than daily | 131 | 36 (27) | 82 | 11 (13) | 215 | 47 (22) | |
| Daily or more | 120 | 33 (28) | 81 | 15 (19) | 203 | 48 (24) | |
| Not last month | 28 | 3 (11) | 12 | 1 (8) | 40 | 4 (10) | |
| p value | | 0.157 | | 0.519 | | 0.158 | |
| 2022 | | | | | | | |
| Less than daily | 91 | 21 (23) | 60 | 7 (12) | 154 | 29 (19) | |
| Daily or more | 143 | 33 (23) | 78 | 9 (12) | 223 | 42 (19) | |
| Not last month | 21 | 1 (5) | 8 | 0 (0) | 29 | 1 (3) | |
| p value | | 0.134 | | 0.832 | | 0.093 | |

Table 9.4.8 HCV RNA prevalence by imprisonment last year, gender and survey year

| Western Australia | | Male | | Female | | Total | |
|------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|--|
| Imprisonment last year | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | |
| 2018* | | | | | | | |
| No imprisonment | 52 | 13 (25) | 41 | 8 (20) | 93 | 21 (23) | |
| Imprisonment | 11 | 4 (36) | 4 | 2 (50) | 15 | 6 (40) | |
| p value | | 0.696 | | 0.181 | | 0.289 | |
| 2019* | | | | | | | |
| No imprisonment | 236 | 47 (20) | 114 | 19 (17) | 351 | 66 (19) | |
| Imprisonment | 55 | 12 (22) | 16 | 3 (19) | 71 | 14 (20) | |
| p value | | 0.816 | | 1.000 | | 0.765 | |
| 2020 | | | | | | | |
| No imprisonment | 174 | 46 (26) | 113 | 26 (23) | 287 | 72 (25) | |
| Imprisonment | 38 | 10 (26) | 12 | 4 (33) | 50 | 14 (28) | |
| p value | | 0.988 | | 0.426 | | 0.663 | |
| 2021 | | | | | | | |
| No imprisonment | 220 | 58 (26) | 156 | 21 (13) | 380 | 80 (21) | |
| Imprisonment | 47 | 12 (26) | 13 | 4 (31) | 61 | 16 (26) | |
| p value | | 0.906 | | 0.091 | | 0.363 | |
| 2022 | | | | | | | |
| No imprisonment | 199 | 40 (20) | 134 | 15 (11) | 338 | 56 (17) | |
| Imprisonment | 46 | 12 (26) | 10 | 2 (20) | 56 | 14 (25) | |
| p value | | 0.371 | | 0.335 | | 0.126 | |

Table 9.4.9 HCV RNA prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year

| Western Australia | | Male | | Female | | Total | |
|--|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|--|
| Aboriginal and Torres Strait Islander origin | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | |
| 2018* | | | | | | | |
| Non Indigenous | 53 | 11 (21) | 30 | 6 (20) | 83 | 17 (20) | |
| Indigenous | 13 | 6 (46) | 16 | 3 (19) | 29 | 9 (31) | |
| p value | | 0.097 | | 0.760 | | 0.334 | |
| 2019* | | | | | | | |
| Non Indigenous | 234 | 46 (20) | 86 | 16 (19) | 319 | 62 (19) | |
| Indigenous | 73 | 16 (22) | 51 | 7 (14) | 126 | 23 (18) | |
| p value | | 0.610 | | 0.366 | | 0.794 | |
| 2020 | | | | | | | |
| Non Indigenous | 148 | 35 (24) | 71 | 18 (25) | 219 | 53 (24) | |
| Indigenous | 62 | 20 (32) | 53 | 11 (21) | 115 | 31 (27) | |
| p value | | 0.196 | | 0.550 | | 0.581 | |
| 2021 | | | | | | | |
| Non Indigenous | 182 | 39 (21) | 108 | 12 (11) | 293 | 51 (17) | |
| Indigenous | 80 | 26 (33) | 65 | 15 (23) | 147 | 42 (29) | |
| p value | | 0.056 | | 0.036 | | 0.007 | |
| 2022 | | | | | | | |
| Non Indigenous | 187 | 40 (21) | 86 | 10 (12) | 275 | 50 (18) | |
| Indigenous | 71 | 16 (23) | 63 | 7 (11) | 137 | 24 (18) | |
| p value | | 0.842 | | 0.922 | 412 | 0.869 | |

Table 9.4.10 HCV RNA prevalence by main language spoken at home by parents, gender and survey year

| Western Australia | | Male | | Female | | Total | |
|---|-----------------------|-----------------------------|-----------------------|-----------------------------|-----------------------|-----------------------------|--|
| Main language spoken at home by parents | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | N ^o tested | N ^o with HCV (%) | |
| 2018* | | | | | | | |
| English speaking | 64 | 15 (23) | 46 | 10 (22) | 110 | 26 (24) | |
| Non-English speaking | 3 | 1 (33) | 1 | 0 (0) | 4 | 1 (25) | |
| p value | | 0.567 | | 0.597 | | 0.798 | |
| 2019* | | | | | | | |
| English speaking | 299 | 58 (19) | 134 | 22 (16) | 434 | 81 (19) | |
| Non-English speaking | 14 | 4 (29) | 3 | 1 (33) | 16 | 5 (31) | |
| p value | | 0.280 | | 0.453 | | 0.178 | |
| 2020 | | | | | | | |
| English speaking | 208 | 57 (27) | 128 | 29 (23) | 336 | 86 (26) | |
| Non-English speaking | 10 | 2 (20) | 4 | 2 (50) | 14 | 4 (29) | |
| p value | | 0.607 | | 0.204 | | 0.803 | |
| 2021 | | | | | | | |
| English speaking | 271 | 70 (26) | 170 | 24 (14) | 447 | 95 (21) | |
| Non-English speaking | 11 | 4 (36) | 6 | 2 (33) | 17 | 6 (35) | |
| p value | | 0.436 | | 0.192 | | 0.169 | |
| 2022 | | | | | | | |
| English speaking | 243 | 52 (21) | 142 | 15 (11) | 390 | 68 (17) | |
| Non-English speaking | 16 | 3 (19) | 6 | 1 (17) | 22 | 4 (18) | |
| p value | | 1.000 | | 0.503 | | 1.000 | |

Table 9.4.11 HCV RNA prevalence by region/country of birth, gender and survey year

| Western Australia | | Male | | Female | | Total | |
|-------------------------|-----------|-----------------|-----------|-----------------|-----------|-----------------|--|
| Region/Country of birth | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | N° tested | N° with HCV (%) | |
| 2018* | | | | | | | |
| Australia | 54 | 12 (22) | 41 | 9 (22) | 95 | 21 (22) | |
| Other Oceania | 1 | 0 (0) | 2 | 0 (0) | 3 | 0 (0) | |
| Asia | 0 | 0 (0) | 1 | 0 (0) | 1 | 0 (0) | |
| UK & Ireland | 9 | 2 (22) | 1 | 1 (100) | 10 | 3 (30) | |
| Other | 3 | 2 (67) | 2 | 0 (0) | 5 | 2 (40) | |
| p value | | 0.227 | | 0.306 | | 0.565 | |
| 2019* | | | | | | | |
| Australia | 266 | 53 (20) | 119 | 20 (17) | 388 | 73 (19) | |
| Other Oceania | 8 | 2 (25) | 5 | 1 (20) | 13 | 3 (23) | |
| Asia | 5 | 1 (20) | 0 | 0 (0) | 5 | 1 (20) | |
| UK & Ireland | 19 | 3 (16) | 9 | 2 (22) | 27 | 5 (19) | |
| Other | 15 | 3 (20) | 3 | 0 (0) | 18 | 3 (17) | |
| p value | | 0.995 | | 0.829 | | 1.000 | |
| 2020 | | | | | | | |
| Australia | 174 | 44 (25) | 123 | 29 (24) | 297 | 73 (25) | |
| Other Oceania | 10 | 2 (20) | 4 | 1 (25) | 14 | 3 (21) | |
| Asia | 4 | 0 (0) | 0 | 0 (0) | 4 | 0 (0) | |
| UK & Ireland | 15 | 6 (40) | 4 | 1 (25) | 19 | 7 (37) | |
| Other | 12 | 5 (42) | 1 | 0 (0) | 13 | 5 (38) | |
| p value | | 0.340 | | 0.957 | | 0.380 | |
| 2021 | | | | | | | |
| Australia | 234 | 61 (26) | 152 | 23 (15) | 392 | 85 (22) | |
| Other Oceania | 11 | 6 (55) | 9 | 2 (22) | 20 | 8 (40) | |
| Asia | 2 | 0 (0) | 2 | 1 (50) | 4 | 1 (25) | |
| UK & Ireland | 18 | 3 (17) | 6 | 0 (0) | 24 | 3 (13) | |
| Other | 8 | 2 (25) | 5 | 0 (0) | 13 | 2 (15) | |
| p value | | 0.196 | | 0.374 | | 0.250 | |
| 2022 | | | | | | | |
| Australia | 223 | 44 (20) | 132 | 14 (11) | 359 | 59 (16) | |
| Other Oceania | 9 | 4 (44) | 3 | 0 (0) | 12 | 4 (33) | |
| Asia | 1 | 0 (0) | 4 | 1 (25) | 5 | 1 (20) | |
| UK & Ireland | 8 | 2 (25) | 7 | 1 (14) | 16 | 3 (19) | |
| Other | 15 | 6 (40) | 3 | 1 (33) | 18 | 7 (39) | |
| p value | | 0.112 | | 0.349 | | 0.072 | |

Appendix A: Participating NSP services

| Year | '18 | '19 | '20 | '21 | '22 |
|--|-----------|-----------|-----------|-----------|-----------|
| Number of NSP sites | 53 | 54 | 38 | 37 | 53 |
| Australian Capital Territory | 1 | 1 | 1 | 1 | 1 |
| Directions Health, Canberra | ✓ | ✓ | ✓ | ✓ | ✓ |
| New South Wales | 19 | 18 | 12 | 3 | 18 |
| ACON Hunter | ✓ | ✓ | ✓ | | ✓ |
| Ballina Harm Reduction Services | ✓ | ✓ | | | ✓ |
| Blacktown HIV/Hepatitis C Prevention Service | ✓ | ✓ | ✓ | | ✓ |
| Byron/Tweed Harm Reduction Services | ✓ | ✓ | | | ✓ |
| Canterbury Harm Minimisation Program | ✓ | ✓ | ✓ | | ✓ |
| Central Coast NSP | | | | | ✓ |
| Clinic 180 (formerly K2), Kings Cross | ✓ | ✓ | | | |
| Coffs Harbour/Grafton Harm Reduction Services | ✓ | ✓ | | | ✓ |
| First Step Program, Port Kembla | ✓ | ✓ | | | |
| Health ConneXions, Liverpool | ✓ | | ✓ | | ✓ |
| Hunter Harm Reduction Services | ✓ | ✓ | ✓ | | ✓ |
| Kelly Close Needle Syringe Program, Mt Druitt | ✓ | ✓ | ✓ | | ✓ |
| Kirketon Road Centre | ✓ | ✓ | ✓ | ✓ | ✓ |
| KRC Kellett St | | | | | ✓ |
| KRC South, Sutherland | ✓ | ✓ | ✓ | ✓ | ✓ |
| Nimbin/Lismore Harm Reduction Services | ✓ | ✓ | | | ✓ |
| NSW Users and AIDS Association (NUAA) | ✓ | ✓ | ✓ | ✓ | ✓ |
| Parramatta HIV/Hepatitis C Prevention Service | ✓ | ✓ | ✓ | | ✓ |
| Redfern Harm Minimisation Program | ✓ | ✓ | ✓ | | ✓ |
| Royal Prince Alfred Hospital, Camperdown | ✓ | ✓ | | | |
| South Court Primary Care NSP | ✓ | ✓ | ✓ | | ✓ |
| Northern Territory | 3 | 3 | 3 | 3 | 3 |
| Northern Territory AIDS & Hepatitis Council, Alice Springs | ✓ | ✓ | ✓ | ✓ | ✓ |
| Northern Territory AIDS & Hepatitis Council, Darwin | ✓ | ✓ | ✓ | ✓ | ✓ |
| Northern Territory AIDS & Hepatitis Council, Palmerston | ✓ | ✓ | ✓ | ✓ | ✓ |
| Queensland | 9 | 10 | 7 | 11 | 10 |
| Biala Community Alcohol and Drug Services, Brisbane | ✓ | ✓ | ✓ | ✓ | ✓ |
| Cairns ATODS NSP | ✓ | ✓ | | ✓ | ✓ |
| Inala Alcohol and Drugs Service | | | | ✓ | ✓ |
| Kobi House, Toowoomba | ✓ | ✓ | ✓ | ✓ | |
| Queensland Injectors Health Network, Brisbane | ✓ | ✓ | ✓ | ✓ | ✓ |
| Queensland Injectors Health Network, Gold Coast | ✓ | ✓ | ✓ | ✓ | ✓ |
| Queensland Injectors Health Network, Sunshine Coast | ✓ | ✓ | ✓ | ✓ | ✓ |
| Rockhampton AODS NSP | | ✓ | | ✓ | ✓ |
| Townsville ATODS NSP | ✓ | ✓ | ✓ | ✓ | ✓ |
| West Moreton Sexual Health Service | ✓ | ✓ | | ✓ | ✓ |
| Youth Link, Cairns | ✓ | ✓ | ✓ | ✓ | ✓ |

| Year | '18 | '19 | '20 | '21 | '22 |
|---|------------|------------|------------|------------|------------|
| Number of NSP sites | 53 | 54 | 38 | 37 | 53 |
| South Australia | 7 | 8 | 7 | 8 | 8 |
| Anglicare Salisbury (formerly SAVIVE – Shopfront) | ✓ | | | | |
| Community Access & Services SA (CASSA) | ✓ | ✓ | ✓ | ✓ | ✓ |
| Drug Arm Australasia | ✓ | ✓ | | | |
| Kurlana Tampawardli, Hendon | | ✓ | ✓ | ✓ | ✓ |
| Noarlunga Primary Health (formerly SAVIVE Noarlunga) | ✓ | ✓ | ✓ | ✓ | ✓ |
| Northern DASSA Elizabeth | | | | ✓ | ✓ |
| Nunkuwarrin Yunti Community Health Centre | ✓ | ✓ | ✓ | ✓ | ✓ |
| Streetlink Youth Services | ✓ | ✓ | ✓ | ✓ | ✓ |
| Watto Purrinna, Port Adelaide (formerly SAVIVE Port Adelaide) | ✓ | ✓ | ✓ | ✓ | ✓ |
| West Coast Youth, Port Lincoln | | ✓ | ✓ | ✓ | ✓ |
| Tasmania | 4 | 4 | 4 | 4 | 4 |
| Clarence Integrated Care Centre | ✓ | ✓ | ✓ | ✓ | ✓ |
| Glenorchy NSP Service | ✓ | ✓ | ✓ | ✓ | ✓ |
| Hobart NSP Service | ✓ | ✓ | ✓ | ✓ | ✓ |
| Salvation Army Launceston | ✓ | ✓ | ✓ | ✓ | ✓ |
| Victoria | 6 | 6 | 0 | 3 | 5 |
| Barwon Health Drug and Alcohol Services | ✓ | ✓ | | ✓ | ✓ |
| Innerspace, Collingwood | ✓ | ✓ | | | ✓ |
| Health Works, Footscray | ✓ | ✓ | | ✓ | ✓ |
| SHARPS, Frankston | ✓ | ✓ | | | |
| North Richmond NSP Services | ✓ | ✓ | | | ✓ |
| Health Information Exchange, St Kilda | ✓ | ✓ | | ✓ | ✓ |
| Western Australia | 4 | 4 | 4 | 4 | 4 |
| HepatitisWA | ✓ | ✓ | ✓ | ✓ | ✓ |
| Peer Based Harm Reduction WA (formerly WASUA) - Perth | ✓ | ✓ | ✓ | ✓ | ✓ |
| Peer Based Harm Reduction WA (formerly WASUA) - Bunbury | ✓ | ✓ | ✓ | ✓ | ✓ |
| WAAC | ✓ | ✓ | ✓ | ✓ | ✓ |

Appendix B: Methodological notes

Capillary blood was obtained by finger-prick using disposable lancets and cotton-fibre blotting paper. Dried blood samples were kept at room temperature at the survey sites and couriered to a central laboratory every second or third day. HIV antibody was detected using the Murex 1.2.0 ELISA (Diasorin). Repeatedly reactive specimens were subjected to Western blot confirmatory testing (Bio-Rad New LAV blot 1, France). A modified third generation enzyme immunoassay (Monolisa anti-HCV Plus Version 3 EIA, Biorad, France) was used to test for HCV antibody. A modified cut off value for optical density was calculated to capture greater than 95% of the seronegative population. Specimens were considered positive for HCV antibody if the optical density to cut off ratio was greater than or equal to one on initial and subsequent testing.

Between 2018 and 2022, HCV RNA was detected and quantified on the Hologic Panther™ using the Aptima™ HCV Quant Dx assay (Hologic, Inc. San Diego, Calif) which has a limit of quantitation (LOQ) 10 IU/mL and an upper limit of quantitation of 10^8 IU/mL in plasma. An offboard DBS elution protocol was applied prior to sampling. To correct for haematocrit, all post run DBS VL results on the Aptima assay were multiplied by a plasma conversion factor of 36.36. This calculation was based on the following assumptions: 45% haematocrit average per DBS, 50uL DBS volume and 1000ul Aptima transport media volume. A qualitative result of <10IU/mL detected (trace/equivocal) was applied to samples <400IU/mL (plasma equivalency).

Data presented in this report were analysed using Stata, Version 14 (Stata Corporation, College Station TX). Temporal changes in HIV/HCV antibody and HCV RNA prevalence were assessed using the χ^2 test for linear trend. Associations between demographic and behavioural variables and HIV/HCV antibody and HCV RNA prevalence were assessed using the χ^2 test or Fisher's exact test where frequencies were less than five. It should be noted that in 2018 and 2019 the sub-sample of respondents with sufficient DBS sample for HCV RNA testing was <90% and HCV RNA tables were weighted to account for potential sample bias with respect to HCV antibody status and gender. Weights were applied to national data and to each individual jurisdiction, such that jurisdictional HCV RNA results will not necessarily add to data presented at the national level. Results with small sample size, particularly at the jurisdictional level, should be interpreted with caution.

Percentage values in tables may not add to 100 due to rounding. Totals in HIV and HCV antibody prevalence tables stratified by gender include people whose gender was not reported or reported as other. Time since first injection was calculated by subtracting age at first injection from age at survey completion. Totals in HIV and HCV antibody prevalence tables stratified by time since first injection include people who did not report age at survey completion or age at first injection. Tables reporting HIV antibody prevalence stratified by demographic and drug use characteristics are not included in this report for states and territories or survey years where the number of HIV antibody positive participants was less than 5 in all years.

Where data were collected but no participants endorsed a particular response, results are presented as "0 (0)". The symbol "--" shown in some tables denotes missing data; and is used to identify circumstances where data were not collected in the reported format. The vast majority of survey questions were consistent over all survey years.

NSP staff recorded sex, age group and survey participation for all clients at the NSP during the survey period and this data was used to assess response rates. Some clients visited the site more than once during the survey week, and only one attendance per person was included in the response rate denominator. Ethical approvals for the study were obtained from Human Research Ethics Committees associated with the investigators and participating sites. Verbal rather than written informed consent was obtained to assure the anonymity of participants.

Appendix C: List of tables

1. Demographic characteristics and drug use

| | |
|-------------|---|
| Table 1.1.1 | Number (percentage) of respondents by demographic characteristics and survey year |
| Table 1.1.2 | Number (percentage) of respondents by last drug injected, frequency of injecting and survey year |
| Table 1.1.3 | Number (percentage) of respondents by injecting behaviour in the last month and survey year |
| Table 1.1.4 | Number (percentage) of respondents by treatment for drug use and survey year |
| Table 1.1.5 | Number (percentage) of respondents by testing for HIV and HCV, HCV treatment by survey year |
| Table 1.1.6 | Number (percentage) of respondents by sexual behaviour in the month prior to survey and survey year |
| Table 1.1.7 | Number (percentage) of respondents by COVID-19 related behaviours in 2022 |

2. HIV antibody prevalence

| | |
|--------------|--|
| Table 1.2.1 | HIV antibody prevalence by gender and survey year |
| Table 1.2.2 | HIV antibody prevalence by sexual identity, gender and survey year |
| Table 1.2.3 | HIV antibody prevalence by age group, gender and survey year |
| Table 1.2.4 | HIV antibody prevalence by last drug injected, gender and survey year |
| Table 1.2.5 | HIV antibody prevalence by years of injection, gender and survey year |
| Table 1.2.6 | HIV antibody prevalence by re-use of someone else's used needle and syringe last month, gender and survey year |
| Table 1.2.7 | HIV antibody prevalence by frequency of drug injection last month, gender and survey year |
| Table 1.2.8 | HIV antibody prevalence by imprisonment last year, gender and survey year |
| Table 1.2.9 | HIV antibody prevalence by condom use at last sex with regular partner, gender and survey year |
| Table 1.2.10 | HIV antibody prevalence by sex work last month, gender and survey year |
| Table 1.2.11 | HIV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year |
| Table 1.2.12 | HIV antibody prevalence by main language spoken at home by parents, gender and survey year |
| Table 1.2.13 | HIV antibody prevalence by region/country of birth, gender and survey year |

3. HCV antibody prevalence

| | |
|--------------|---|
| Table 1.3.1 | HCV antibody prevalence by gender and survey year |
| Table 1.3.2 | HCV antibody prevalence by sexual identity, gender and survey year |
| Table 1.3.3 | HCV antibody prevalence by age group, gender and survey year |
| Table 1.3.4 | HCV antibody prevalence by years since first injection, gender and survey year |
| Table 1.3.5 | HCV antibody prevalence by re-use of someone else's used needle and syringe last month, years since first injection and survey year |
| Table 1.3.6 | HCV antibody prevalence by last drug injected, years since first injection and survey year |
| Table 1.3.7 | HCV antibody prevalence by frequency of drug injection last month, years since first injection and survey year |
| Table 1.3.8 | HCV antibody prevalence by imprisonment last year, years since first injection and survey year |
| Table 1.3.9 | HCV antibody prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year |
| Table 1.3.10 | HCV antibody prevalence by main language spoken at home by parents, gender and survey year |
| Table 1.3.11 | HCV antibody prevalence by region/country of birth, gender and survey year |

4. HCV RNA prevalence

- Table 1.4.1 HCV RNA prevalence by gender and survey year
- Table 1.4.2 HCV RNA prevalence by sexual identity, gender and survey year
- Table 1.4.3 HCV RNA prevalence by age group, gender and survey year
- Table 1.4.4 HCV RNA prevalence by years since first injection, gender and survey year
- Table 1.4.5 HCV RNA prevalence by re-use of someone else's used needle and syringe last month, gender and survey year
- Table 1.4.6 HCV RNA prevalence by last drug injected, gender and survey year
- Table 1.4.7 HCV RNA prevalence by frequency of drug injection last month, gender and survey year
- Table 1.4.8 HCV RNA prevalence by imprisonment last year, gender and survey year
- Table 1.4.9 HCV RNA prevalence by Aboriginal and Torres Strait Islander origin status, gender and survey year
- Table 1.4.10 HCV RNA prevalence by main language spoken at home by parents, gender and survey year
- Table 1.4.11 HCV RNA prevalence by region/country of birth, gender and survey year

